



TAXANES FROM *TAXUS YUNNANENSIS*

HONGJIE ZHANG, YOSHIO TADEDA*† and HANDONG SUN*†

Laboratory of Phytochemistry, Kunming Institute of Botany, Academia Sinica, Kunming 650 204, Yunnan, China; *Faculty of Integrated Arts and Sciences, The University of Tokushima, Tokushima 770, Japan

(Received in revised form 4 January 1995)

Key Word Index—*Taxus yunnanensis*; Taxaceae; roots; diterpenoids; taxanes; taxuyunnanine E and F.

Abstract—Two new taxane diterpenoids as well as a known one, taxayunnansin A, were isolated from the roots of *Taxus yunnanensis*. One of the new compounds was identified as a unique taxane with a 2,20-oxetane ring. Their structures were elucidated by spectroscopic techniques, especially by 2D NMR spectra.


INTRODUCTION

Although taxol, a unique diterpenoid alkaloid, has shown unusual anticancer activity and has been recently approved for treating the sufferers of advanced ovarian cancers [1], its scarcity has limited its widespread use and this has stimulated a search for new sources of taxol. Semi-synthesis from natural taxoids has been successful in producing taxol and related compounds [2]. The total synthesis to taxol has recently been achieved [3], but it has an uncertain future in affording large-scale production of taxol. Cell culture is hoped to provide a new resource for producing taxol or taxol precursors. The search has resulted in the isolation of a large number of new taxoids from *Taxus* plants. Among these new taxoids, a new kind of taxane diterpenoid with an A-5 ring rearranged skeleton has been discovered [4, 5]. As a continuation of our investigation on *T. yunnanensis* [6–8], A-5 ring-rearranged taxoids were also obtained. We now report three such taxanes: taxayunnansin A (2) [5] and two new ones, named taxuyunnanine E (1) and F (3), respectively. These three compounds were isolated from the ethereal extracts of the roots of *T. yunnanensis* as minor constituents. One of them (1) proved to have a 2,20-oxetane ring and H- α configuration on C-9, which has never been reported in *Taxus* plants before. The structures of the two novel taxanes were determined by spectroscopic means, including 2D NMR spectra.

RESULTS AND DISCUSSION

Compound 1, $[\alpha]_D^{26} + 3.3^\circ$ (CHCl₃; c 0.6), produced a $[M - 1]^-$ ion at m/z 629 in the FAB-mass spectrum. The combination of FAB-mass and ¹³C NMR spectra (including DEPT technique) suggested a formula

C₃₃H₄₂O₁₂, which was confirmed by high-resolution FAB-mass spectrum (m/z 629.25797 $[M - H]^-$). Compound 1 was shown to have four methyl groups, two methylene groups, one oxymethylene group, one methine group, six oxymethine groups, two quaternary carbons, two oxyquaternary carbons, two olefinic quaternary carbons, three acetyl groups and one benzoyl group in its ¹H and ¹³C NMR spectra (Tables 1 and 2). Based on the comparison of NMR spectra of 1 with those of known taxanes, as well as consideration of the structures of taxanes from the genus *Taxus*, we presumed 1 to have a taxane diterpenoid skeleton. By comparing the ¹H and ¹³C NMR spectra of 1 with those of taxanes from the genus *Taxus*, we could easily assign the only methine group as C-3 [δ_H 2.57 (1H, *dd*, $J = 7.7$ and 1.6 Hz) and δ_C 51.1 (*d*)], the only oxymethylene group as C-20 [δ_H 4.37 (1H, *dd*, $J = 11.5$ and 1.8 Hz) and δ_C 3.69 (1H, *d*, $J = 11.5$ Hz) and δ_C 72.8 (*t*)], one quaternary carbon as C-8 (δ_C 43.2), the two olefinic quaternary carbons as C-11 (δ_C 139.5) and C-12 (δ_C 141.7), the four methyls as Me-16 [δ_H 1.10 (3H, *s*) and δ_C 27.8 (*q*)], Me-17 [δ_H 1.50 (3H, *s*) and



† Authors to whom correspondence should be addressed.

Table 1. ^1H NMR spectral data of compounds 1–3 [in CDCl_3 , 400 MHz, δ in ppm with reference to the signal of CDCl_3 ($\delta 7.24$ ppm)]

H	1	2	3
2	4.93 <i>d</i> (7.7)	6.06 <i>d</i> (7.8)	5.87 <i>d</i> (7.4)
3	2.57 <i>dd</i> (7.7, 1.6)	3.09 <i>d</i> (7.8)	2.84 <i>d</i> (7.4)
5	4.30 <i>ddd</i> (11.4, 7.8, 2.6)	4.91 <i>d</i> (7.3)	4.91 <i>dd</i> (8.2, 1.6)
6 α	2.00 <i>m</i>	2.48 <i>dt</i> (16.1, 8.2)	2.44 <i>dt</i> (16.0, 8.2)
6 β	1.84 <i>m</i>	1.85 <i>dd</i> (16.0, 8.7)	1.18 <i>m</i>
7	4.82 <i>dd</i> (12.6, 3.2)	5.50 <i>t</i> (8.3)	5.34 <i>t</i> (8.4)
9	5.20 <i>d</i> (4.0)	6.10 <i>d</i> (10.8)	5.73 <i>d</i> (10.3)
10	4.90 <i>d</i> (4.0)	6.49 <i>d</i> (10.8)	4.53 <i>t</i> (10.1)
13	5.62 <i>br t</i> (6.9)	4.45 <i>m</i>	5.55 <i>t</i> (7.2)
14 α	2.12 <i>dd</i> (14.9, 7.4)	1.59 <i>dd</i> (15.1, 7.2)	1.57 <i>m</i>
14 β	2.28 <i>dd</i> (14.9, 7.3)	2.20 <i>dd</i> (15.2, 7.5)	2.06 <i>m</i>
Me-16	1.10 <i>s</i>	1.02 <i>s</i>	1.05 <i>s</i>
Me-17	1.50 <i>s</i>	1.12 <i>s</i>	1.12 <i>s</i>
Me-18	1.35 <i>s</i>	1.97 <i>s</i>	1.76 <i>s</i>
Me-19	1.52 <i>s</i>	1.63 <i>s</i>	1.57 <i>s</i>
20a	4.37 <i>dd</i> (11.5, 1.8)	4.50 <i>AB d</i> (7.6)	4.38 <i>AB d</i> (7.5)
20b	3.69 <i>d</i> (11.5)	4.36 <i>AB d</i> (7.6)	4.32 <i>AB d</i> (7.5)
H ₂ -3''	7.96 <i>dd</i> (8.0, 1.6)	7.80 <i>dd</i> (7.3, 1.2)	
H ₂ -4''	7.40 <i>tt</i> (7.6, 1.6)	7.37 <i>t</i> (7.6)	
5''	7.55 <i>tt</i> (7.8, 1.6)	7.50 <i>tt</i> (7.6, 1.1)	
OH-5	4.40 <i>d</i> (2.6)		
OH-10			3.84 <i>d</i> (10.0)
OH-15		2.72 <i>br s</i>	2.29 <i>br s</i>
OAc-Me	2.08 <i>s</i>	2.11 <i>s</i>	2.06 <i>s</i>
	2.05 <i>s</i>	2.03 <i>s</i> (2-)	2.06 <i>s</i>
	1.93 <i>s</i> (7-)	1.98 <i>s</i>	2.04 <i>s</i>
		1.69 <i>s</i> (9-)	2.01 <i>s</i>
			1.94 <i>s</i>

$\delta_{\text{C}} 27.9$ (*q*), Me-18 [$\delta_{\text{H}} 1.35$ (3H, *s*) and $\delta_{\text{C}} 12.9$ (*q*)] and Me-19 [$\delta_{\text{H}} 1.52$ (3H, *s*) and $\delta_{\text{C}} 15.0$ (*q*)], respectively. From these NMR signals, we could assign all other proton and carbon NMR signals by using ^1H - ^1H COSY, ^1H - ^{13}C HETCOR, and COLOC techniques (Tables 3 and 4). The long-range correlation of two quaternary carbons ($\delta_{\text{C}} 65.0$ and 76.3) to Me-16 and Me-17 revealed one of them was C-1 and the other was C-15, and the signal of $\delta_{\text{C}} 65.0$ was further judged as C-1 owing to its long-range correlation to H-3. Compared with that of normal taxanes, the unusual shift downfield of the C-1 signal and upfield of the C-15 signal in ^{13}C NMR spectrum suggested **1** to have an A-ring rearranged carbon skeleton [4], which was supported by the following. The signal at $\delta_{\text{H}} 5.20$ (1H, *d*, $J = 4.0$ Hz) was assigned to H-9 from observation of long-range couplings to C-3 and C-11, from which we established the signal at $\delta_{\text{H}} 4.90$ (1H, *d*, $J = 4.0$ Hz) as H-10 owing to the presence of direct coupling to H-9 in the COSY spectrum. H-10 showed three-bond coupling with C-1 in the COLOC spectrum, which thus confirmed **1** to have the rearranged carbon skeleton of taxayunnansin A (**2**).

The ^{13}C NMR and DEPT spectra (Table 2) showed the presence of two methylenes in **1**. HETCOR experiment proved one methylene carbon ($\delta_{\text{C}} 29.9$) as attached by the $\delta_{\text{H}} 2.00$ (1H, *m*) and 1.84 (1H, *m*) protons, and the other one ($\delta_{\text{C}} 35.6$) was attached by the $\delta_{\text{H}} 2.12$ (1H, *dd*,

$J = 14.9$ and 7.4 Hz) and 2.28 (1H, *dd*, $J = 14.9$ and 7.3 Hz) protons. In addition, the COLOC spectrum revealed three-bond couplings of one methylene proton ($\delta_{\text{H}} 2.00$ and 1.84) to C-8 and another methylene proton ($\delta_{\text{H}} 2.12$ and 2.28) to C-15, which indicated the former methylene was C-6 and the latter was C-14. Since the determination of the two methylenes, the six oxymethines in **1** could only be assigned to C-2, C-5, C-7, C-9, C-10 and C-13, respectively. The remaining oxyquaternary carbon ($\delta_{\text{C}} 94.6$) on the skeleton of **1** was determined to be C-4 by considering its long-range correlations to H-3, H-20 and H-6. Examining the ^{13}C NMR data between **1** and taxayunnansin A (**2**), we noticed that the ^{13}C NMR data changed strikingly at C-2, C-3, C-4 and C-5 [$\delta_{\text{C}} 79.2$ (*d*, C-2), 51.1 (*d*, C-3), 94.6 (*s*, C-4) and 71.7 (*d*, C-5)] in **1**. The 13 ppm upfield shift of C-5 revealed the scission of a 5,20-epoxy group. The observation of three-bond coupling of C-2 and H-20a [$\delta_{\text{H}} 4.37$ (*dd*, $J = 11.5$ and 1.8 Hz)] strongly suggested the connection between C-2 and C-20 through an oxygen atom. The severely deshielded ^{13}C NMR shift of C-4 [5] was explained by the hydrogen bond of OH-5 [$\delta_{\text{H}} 4.40$ (*d*, $J = 2.6$ Hz)] formed to the oxygen atom which connected to C-4. The same phenomenon has been observed in taxayunnansin A and related compounds, in which the hydrogen bond of OH-10 to the carbonyl oxygen atom of C-9 led to a 7 ppm downfield shift of C-9 in the ^{13}C NMR spectra [7].

Table 2. ^{13}C NMR spectral data of compounds 1–3 [in CDCl_3 , 100 MHz, δ in ppm from the signal of CDCl_3 ($\delta 77.0$ ppm)]

C	1	2	3	DEPT
1	65.0	67.7	67.0	C
2	79.2	68.1	68.7	CH
3	51.1	44.1	44.7	CH
4	94.6	79.9	79.4	C
5	71.7	85.0	84.6	CH
6	29.9	34.8	34.7	CH_2
7	70.5	70.4	70.6	CH
8	43.2	43.8	43.3	C
9	75.2	76.6	79.2	CH
10	69.1	69.2	66.1	CH
11	139.5	134.0	139.9	C
12	141.7	151.5	142.8	C
13	81.8	77.0	79.4	CH
14	35.6	39.6	36.5	CH_2
15	76.3	75.5	76.2	C
16	27.8	27.5	27.8	CH_3
17	27.9	25.4	25.5	CH_3
18	12.9	11.8	11.2	CH_3
19	15.0	12.6	12.6	CH_3
20	72.8	74.9	74.5	CH_2
1''	164.8	164.1		C
2''	129.7	129.2		C
3''	129.7	129.5		$\text{CH} \times 2$
4''	128.4	128.7		$\text{CH} \times 2$
5''	133.3	133.3		CH
OAc-Me	21.9	22.2	21.9	CH_3
	20.9 (7-)	21.5	21.4	CH_3
	20.9	21.3 (2-)	21.4	CH_3
		20.5 (9-)	21.1	CH_3
			21.1	CH_3
OAc-C=O	172.0	171.0	170.7	C
	170.3	170.3 (2-)	170.4	C
	170.1 (7-)	169.8 (9-)	170.4	C
		169.8	169.9	C
			169.0	C

Similarly to most taxane compounds, **1** also possessed acyl groups. Its NMR spectra showed the presence of three acetyl and one benzoyl group, which could be assigned by the following considerations. The long-range COLOC spectrum revealed the correlations of the proton signals at $\delta_{\text{H}} 4.82$ (1H, *dd*, $J = 12.6$ and 3.2 Hz, H-7) and 5.20 (1H, *d*, $J = 4.0$ Hz, H-9) with the carbon signals at $\delta_{\text{C}} 170.1$ (*s*, the carbonyl carbon of an acetyl) and 164.8 (*s*, the carbonyl carbon of benzoyl) respectively, indicating that an acetoxy group was attached to C-7 [$\delta_{\text{C}} 70.5$ (*d*)] and the benzyloxy group to C-9 [$\delta_{\text{C}} 75.2$ (*d*)]. An acetoxy substituent was assigned to C-13 [$\delta_{\text{C}} 81.8$ (*d*)] owing to the deshielding of H-13 in the ^1H NMR spectrum [$\delta_{\text{H}} 5.62$ (1H, *br t*, $J = 6.9$ Hz)]. The ^1H NMR data of H-10 and H-5 appeared at usual chemical shifts for hydroxylated positions [$\delta_{\text{H}} 4.90$ (1H, *d*, $J = 4.0$ Hz, H-10) and $\delta_{\text{H}} 4.30$ (1H, *ddd*, $J = 11.4$, 7.8 and 2.6 Hz, H-5)]. In the EI-mass spectrum of **1**, the prominent peaks at m/z 512 [$\text{M} - \text{MeCO}_2\text{H} - \text{Me}_2\text{CO}]^+$ (7%), 452 [$\text{M} - \text{MeCO}_2\text{H} \times 2 - \text{Me}_2\text{CO}]^+$ (83%), 392 [$\text{M} - \text{MeCO}_2\text{H} \times 3 - \text{Me}_2\text{CO}]^+$ (12%) and 390 [$\text{M} - \text{C}_6\text{H}_5\text{CO}_2\text{H} -$

Table 3. 2D ^1H - ^1H COSY and NOESY data for compounds 1 and 3 (in CDCl_3)

H	COSY		NOESY
	1	3	1
2	H-3	H-3	H-3, 19
3	H-2, 20a	H-2, 20b	H-2
5	H-6	H-6	
H ₂ -6	H-5, 7		
6 α		H-5, 6 β , 7	
6 β		H-6 α , 7	
7	H-6	H-6	H-10
9	H-10	H-10	H-10, 19
10	H-9	H-9, 10-OH	H-7, 9, 18
13	H-14	H-14, 18	
14 α	H-13, 14 β	H-13, 14 β	
14 β	H-13, 14 α	H-13, 14 α	
Me-16	H-17	H-17	
Me-17	H-16	H-16	
Me-18		H-13	
Me-19			
20a	H-3, 20b	H-20b	H-20b
20b	H-20a	H-3, 20a	H-20a
H ₂ -3''	H-4''		H-17, 19, 4''
H ₂ -4''	H-3'', 5''		H-3'', 5''
5''	H-4''		H-4''
OH-5			H-6
OH-10		H-10	

$\text{MeCO}_2\text{H} - \text{Me}_2\text{CO}]^+$ (11%) confirmed the presence of a $\text{Me}_2(\text{OH})$ group in **1**. We could, therefore, conclude that the last acetoxy group was connected to C-4.

The orientation of those secondary oxy groups could be obtained by analysing the ^1H - ^1H NOESY spectrum (Table 3). The observation of NOEs between H-2 and Me-19 established the oxy substituents of C-2 as α -oriented. Both H-7 and H-10 were determined to be α owing to the presence of a NOESY correlation between the two protons. H-9 was also assigned to α -configuration owing to its small coupling constant to H-10 ($J = 4.0$ Hz) and the observation of NOEs between the benzoyl protons and Me-17. The last secondary oxy group at C-13 was judged to be α from comparison of the coupling pattern of H-13 with those similar congeners. Consequently, we established the structure of taxuyunnanin E as **1**.

Compound **3**, [α] $^{27} - 22.6^\circ$ (CHCl_3 ; c 0.9), had a molecular formula of $\text{C}_{30}\text{H}_{42}\text{O}_{13}$, which was determined by the high-resolution FAB-mass spectrum (609.25131 [$\text{M} - 1$] $^-$). The ^1H and ^{13}C NMR spectra revealed metabolite **3** closely resembled that of taxchinin B [9] except that **3** contained five acetate groups but no other kind of acyloxy groups. ^1H - ^1H COSY, ^1H - ^{13}C HETCOR and COLOC (Tables 3 and 4) were used to assign proton and carbon signals. The chemical shifts at $\delta_{\text{H}} 5.87$ (1H, *d*, $J = 7.4$ Hz), 5.34 (1H, *t*, $J = 8.4$ Hz), 5.73 (1H, *d*, $J = 10.3$ Hz), 4.53 (1H, *t*, $J = 10.1$ Hz) and 5.55 (1H, *t*, $J = 7.2$ Hz) were assigned as H-2, H-7, H-9,

Table 4. COLOC data for compounds **1** and **3** (in CDCl₃)

C	Correlated proton	
	1 (6/8 Hz)	3 (8 Hz)
1	H-3,10,14 α ,16,17	H-16,17
2	H-20a	H-2
3	H-3,9,19,20a	H-19
4	H-3,6 α ,20	H-20b
5	H-6,5-OH	H-20b
6	5-OH	
7	H-6,7,19	
8	H-3,6,10,19	H-19
9	H-9,10,19	H-3,19
10	H-9	
11	H-9,10,18	H-18
12	H-10,14 β ,18	H-18
13	H-14 α ,18	H-18
14	H-14	
15	H-14,16,17	H-16,17
16	H-17	H-16,17
17	H-16	H-16,17
18	H-18	
19	H-19	
20	H-20	H-20
1''	H-9,3''	
2''		
3''	H-3'',4'',5''	
4''	H-4'',5''	
5''	H-3''	
OAc-Me-2		H-2-OAc
OAc-C = O-2		H-2-OAc
OAc-Me-4	H-4-OAc	H-4-OAc
OAc-C = O-4	H-4-OAc	H-4-OAc
OAc-Me-7	H-7-OAc	H-7-OAc
OAc-C = O-7	H-7,7-OAc	H-7-OAc
OAc-Me-9		H-9-OAc
OAc-C = O-9		H-9-OAc
OAc-Me-13	H-13-OAc	H-13-OAc
OAc-C = O-13	H-13-OAc	H-13-OAc

H-10 and H-13, respectively according to the results of 2D ¹H-¹H COSY, ¹H-¹³CHETCOR and COLOC experiments (Tables 3 and 4). Comparing the ¹H NMR shifts of **3** with those of taxchinin B, the deshielded H-2, H-7, H-9 and H-13 signals established four acetate groups attached to C-2, C-7, C-9 and C-13 respectively, but the usual chemical shift of H-10 indicated no acetate group attached to C-10. Additionally, the stronger ion peak at *m/z* 432 in the EI-mass spectrum of **3**, being produced by the fragment pathway of [M - MeCO₂H × 2 - Me₂CO], suggested the remaining acetate group attached to C-4 but not to C-15. Thus the structure of taxuyunnanine F is **3**.

Compounds **1-3** all belonged to A-5 ring rearranged taxoids, and **1** was the first example having a 2,20-oxetane ring and the H- α configuration on C-9 in the taxane diterpenoid family. The stereo-conformation of

2 [**5**] was similar to that of usual taxoids, which implied that the A-5 ring rearranged taxoids with a similar taxol 13-side chain would possess similar bioactivity to that of taxol.

EXPERIMENTAL

Prep. HPLC: 10 C₁₈ Cosmosil Packed Column, 20 × 250 mm. NMR: Bruker AM-400 Fourier transform spectrometer operating at 400.134 and 100.614 MHz for ¹H and ¹³C. The samples were made up in CDCl₃. All chemical shifts are expressed in ppm with reference to the solvent signals: 7.24 ppm/77.0 ppm for CDCl₃.

Plant material. The roots were collected in the suburb of Kunming, Yunnan, China and identified as *Taxus yunnanensis*. A voucher specimen is kept in the Yunnan Academy of Forestry, Kunming, Yunnan, China.

Isolation of taxuyunnanine E and F. Dried, powdered roots (40 kg) of *T. yunnanensis* were extracted with Et₂O (6 × 100 l) at room temp. during 3 weeks to give 650 g crude extract. The ethereal extract was chromatographed on a silical gel column (2 kg), which was eluted with CHCl₃-Me₂CO. The CHCl₃-Me₂CO (4:1) eluting part (500 mg) was subjected to prep. HPLC (10 C₁₈ Cosmosil Packed Column, 20 × 250 mm) eluting with MeOH-H₂O (3:1) to give taxuyunnanine E (**1**, 14.1 mg), F (**3**, 35.6 mg), and taxayunnansin A (**2**, 27.7 mg), respectively.

Taxuyunnanine E (1). C₃₃H₄₂O₁₂, powders, [α]_D²⁶ + 3.3° (CHCl₃; *c* 0.6); IR ν_{\max}^{KBr} cm⁻¹: 3400, 3250, 2960, 2910, 1715, 1595, 1450, 1357, 1320, 1260, 1175, 1105, 1070, 1040, 1015, 945, 828, 810, 755, and 715; EIMS *m/z*: 612 [M - H₂O]⁺ (10%), 597 [M - Ac]⁺ (6%), 570 [M - MeCO₂H]⁺ (4%), 552 [M - MeCO₂H - H₂O]⁺ (2%), 512 [M - MeCO₂H - Me₂CO]⁺ (7%), 481 [M - C₆H₅CHO - Ac]⁺ (4%), 452 [M - MeCO₂H × 2 - Me₂CO]⁺ (83%), 434 [M - MeCO₂H × 2 - Me₂CO - H₂O]⁺ (15%), 392 [M - MeCO₂H × 3 - Me₂CO]⁺ (12%), 390 [M - C₆H₅CO₂H - MeCO₂H - Me₂CO]⁺ (11%), 373 [M - MeCO₂H × 3 - C₆H₅]⁺ (27%), 356 [M - MeCO₂H × 3 - Me₂CO - H₂O × 2]⁺ (13%), 330 [M - C₆H₅CO₂H - MeCO₂H × 2 - Me₂CO]⁺ (84%), 313 [M - C₆H₅CO₂H - MeCO₂H × 2 - Me₂CO - OH]⁺ (58%), 297 [M - MeCO₂H × 3 - Me₂CO - C₆H₅ - H₂O]⁺ (22%), 270 [M - C₆H₅CO₂H - MeCO₂H × 3 - Me₂CO]⁺ (41%), 252 [M - C₆H₅CO₂H - MeCO₂H × 3 - Me₂CO - H₂O]⁺ (34%), 179 (28%), 148 (53%), 133 (71%), 122 [C₆H₅CO₂H]⁺ (43%), 105 [C₆H₅CO]⁺ (64%), 77 [C₆H₅]⁺ (96%), 60 [MeCO₂H]⁺ (37%), and 43 [Ac]⁺ (100%); FAB-MS *m/z*: 629 [M - 1]⁻ (100%); HRFAB-MS *m/z* 629.25797 (calcd 629.25980 for C₃₃H₄₁O₁₂); ¹H and ¹³C NMR data: see Tables 1 and 2, respectively.

Taxayunnansin A (2). C₃₃H₄₄O₁₃, powders; IR ν_{\max}^{KBr} cm⁻¹: 3530, 2960, 2875, 1725, 1435, 1365, 1260, 1235, 1180, 1085, 1065, 1025, 982, 915, 845, 712, and 602; FAB-MS *m/z*: 711 [M + K]⁺ (12%), and 695 [M + Na]⁺ (13%); ¹H and ¹³C NMR data: see Tables 1 and 2, respectively.

Taxuyunnanine F (3). $C_{30}H_{42}O_{13}$, powders, $[\alpha]_D^{27}$ -22.6° ($CHCl_3$, c 0.9); IR $\nu_{max}^{KBr} cm^{-1}$: 3400, 2970, 2925, 2880, 1730, 1435, 1369, 1235, 1025, 985, 950, 898, 845, 780, 755, and 605; EIMS m/z : 432 $[M - MeCO_2H \times 2 - Me_2CO]^+$ (44%), 372 $[M - MeCO_2H \times 3 - Me_2CO]^+$ (38%), 312 $[M - MeCO_2H \times 4 - Me_2CO]^+$ (21%), 297 (72%), 58 $[Me_2CO]^+$ (59%), and 43 $[Ac]^+$ (100%); FAB-MS m/z : 609 $[M - 1]^-$ (100%); HRFAB-MS m/z 609.25131 (calcd 609.25472 for $C_{30}H_{41}O_{13}$); 1H and ^{13}C NMR data: see Tables 1 and 2, respectively.

Acknowledgements—The authors are grateful to Mr Xiang Wei (Yunnan Academy of Forestry) for providing the plant material, and to the members of the analytical group in the Phytochemistry Laboratory of Kunming Institute of Botany, Academia Sinica for measuring the mass spectra, IR and optical rotation data. We also wish to thank the NMR Laboratory of Institute of Botany Academia Sinica for opportunities to record NMR spectra. This work was supported in part by Goho Life Science International Fund of Japan.

REFERENCES

- Joyce, C. (1993) *BioScience* **43**, 133.
- Kingston, D. G. I. (1991) *Pharmac. Ther.* **52**, 1.
- Nicolaou, K. C., Yang, Z., Liu, J. J., Ueno, H., Nantmet, P. G., Guy, R. K., Claibome, C. F., Renaud, J., Couladouros, E. A., Paulvannan, K. and Sorenson, E. J. (1994) *Nature* **367**, 630.
- Fuji, K., Tanaka, K., Li, B., Shingu, T., Sun, H. D. and Taga, T. (1992) *Tetrahedron Letters* **33**, 7915.
- Liu, X. K., Wu, D. G. and Wang, Z. Y. (1992) *Kexue Tongbao* **23**, 2186.
- Zhang, H. J., Takeda, Y., Minami, Y., Yoshida, K., Unemi, N., Mu, Q., Xiang, W., Matsumoto, T. and Sun, H. D. (1993) *Acta Botan. Yunnan.* **15**, 424.
- Zhang, H. J., Takeda, Y., Matsumoto, T., Minami, Y., Yoshida, K., Xiang, W., Mu, Q. and Sun, H. D. (1994) *Heterocycles* **38**, 975.
- Zhang, H. J., Takeda, Y., Minami, Y., Yoshida, K., Matsumoto, T., Xiang, W., Mu, Q. and Sun, H. D. (1994) *Chem. Letters* 957.
- Wahl, A., Gueritte-Voegelein, F., Guenard, D., Goff, T. and Potier, P. (1992) *Tetrahedron* **6965**.