

MEGASTIGMANE GLUCOSIDES FROM *STACHYS BYZANTINA*

YOSHIO TAKEDA,* HONGJIE ZHANG,† TOSHIYA MASUDA, GISHO HONDA,‡ HIDEAKI OTSUKA,§
 EKREM SEZIK,|| ERDEM YESILADA|| and HANDONG SUN¶

Faculty of Integrated Arts and Sciences, The University of Tokushima, Tokushima 770, Japan; ‡Faculty of
 Pharmaceutical Sciences, Kyoto University, Sakyo-ku, Kyoto 606, Japan; §Institute of Pharmaceutical Sciences,
 Hiroshima University School of Medicine, Hiroshima 734, Japan; ||Faculty of Pharmacy, Gazi University,
 Ankara 06330, Turkey; ¶Kunming Institute of Botany, Academia Sinica, Kunming, Yunnan 650204, China

(Received 27 August 1996)

Key Word Index—*Stachys byzantina*; Labiatae; byzantionosides A and B; megastigmene
 glucoside.

Abstract—From the aerial parts of *Stachys byzantina*, two new megastigmene glucosides, byzantionosides A and B, were isolated, together with the known glycosides, icaraside B₂, blumeol C glucoside, (6*R*, 9*R*)- and (6*R*, 9*S*)-3-oxo- α -ionol glucosides and verbascoside. The structures of the new compounds have been elucidated by spectroscopic means. Copyright © 1997 Elsevier Science Ltd

INTRODUCTION

To the best of our knowledge, no reports have appeared on the constituents of *Stachys byzantina* C. Koch. This paper deals with the isolation of two new megastigmene glucosides, byzantionosides A (1) and B (3), together with the known compounds, icaraside B₂ [1], blumeol C glucoside [2], (6*R*, 9*R*)- and (6*R*, 9*S*)-3-oxo- α -ionol glucosides [3] and verbascoside [4] from the aerial parts of the title plant collected in northern Anatolia.

RESULTS AND DISCUSSION

Compound 1, $[\alpha]_D + 74.9^\circ$ (methanol) was obtained as an amorphous powder, and the molecular formula was determined to be C₁₉H₃₀O₇ based on its high resolution negative ion FAB mass spectrum. The ¹H and ¹³C NMR spectra showed the presence of a methyl ketone group conjugated with a *trans*-double bond [δ_H 2.26 (3H, *s*), δ_C 27.1 (*q*) and 201.0 (*s*); δ_H 6.12 (1H, *d*, *J* = 15.6 Hz) and 6.67 (1H, *dd*, *J* = 15.6 and 9.8 Hz), δ_C 134.4 and 149.7 (each *d*)], a methine group [δ_H 2.56 (1H, *d*, *J* = 9.8 Hz), δ_C 55.7 (*d*)] which is connected with the *trans*-double bond, a trisubstituted double bond [δ_H 5.74 (1H, *d*, *J* = 1.5 Hz), δ_C 125.9 (*d*) and 136.7 (*s*)] and a secondary oxymethine group [δ_H 4.36 (1H, *m*), δ_C 73.9 (*d*)] in addition to the signals

Table 1. ¹³C NMR data* for byzantionosides A (1) and B (3) and blumeol C glucoside (4)

C	1	3	4
1	34.6	37.2	37.3
2	40.7	48.0	48.0
3	73.9	202.3	202.2
4	125.9	125.3	125.4
5	136.7	170.0	169.7
6	55.7	52.3	52.4
7	149.7	26.7	26.6
8	134.4	37.7	37.3
9	201.0	75.0	77.5
10	27.1	19.8	21.9
11	29.6	27.5	27.5
12	25.4	29.0	29.0
13	22.9	25.0	25.0
1'	103.1	102.0	103.9
2'	75.1	75.4	75.2
3'	77.9	78.0	78.1
4'	71.6	71.7	71.5
5'	78.2	77.7	77.7
6'	62.8	62.8	62.7

*Measured at 100 MHz for CD₃OD solution.

due to a methylene group, a methyl group on the latter double bond, two tertiary methyl groups and a β -glucopyranosyl moiety (Table 1) in the structure of 1. These spectral data, coupled with the UV data, suggested that 1 has an ionol glucoside structure having the same side chain structure as that of alan-gionoside L [5]. The orientation of the glucosyloxy group was inferred as pseudo- α -equatorial judged

*Author to whom correspondence should be addressed.

†On leave from Kunming Institute of Botany, Academia Sinica.




Fig. 1. NOE correlations for byzantionoside A tetraacetate (2) detected by differential NOE experiments.

from the coupling pattern of H₂-2 [δ_{H} 1.57 (1H, *dd*, $J = 13.7$ and 5.9 Hz) and 1.83 (1H, *dd*, $J = 13.7$ and 6.4 Hz)] [6]. The relative stereochemistry in the aglycone portion was elucidated on the basis of the results of differential NOE experiments for the tetraacetate (2) as shown in Fig. 1. The absolute stereochemistry at C-6 was determined to be *6R* by comparing the circular dichroic (CD) spectrum ($\Delta \epsilon_{239} + 8.04$) to that reported for (3*R*, 6*R*)-3-methoxy- α -ionone [6].

Compound 3, $[\alpha]_{\text{D}}^{25} + 48.0^{\circ}$ (methanol), was also obtained as an amorphous powder, and the elemental composition was determined as C₁₉H₃₂O₇ based on its high resolution negative ion FAB mass spectrum, which is the same as that of blumeol C glucoside (4). Usual acetylation gave the tetraacetate (5). The ¹H and ¹³C NMR (Table 1) spectra of 3 were very similar to those of 4 except for the resonance due to C-9 (δ_{C} 75.0). The corresponding signals were observed at δ_{C} 77.5 and 76.8 in 4 and dihydroalanganoside A [7], respectively, which have the *R*-configuration at C-9, whereas the C-9 signal was observed at δ_{C} 74.6 in salvionoside C [8], which has the *S*-configuration at C-9. These observations indicated that 3 has the same structure as that of 4 except for the stereochemistry at C-9, i.e. *9S*. The stereochemistry at C-6 was confirmed as *6R* by the fact that the CD spectrum showed a positive extreme at 237 nm [$\Delta \epsilon + 3.28$], which was essentially the same as that [$\Delta \epsilon_{236} + 3.02$] of 4. Thus, the structure of byzantionoside B was elucidated as 3, an epimer of 4 at C-9.

EXPERIMENTAL


General. ¹H (400 MHz) and ¹³C (100 MHz) NMR, TMS as int. standard. FABMS: matrix PEG-400. CC: silica gel 60 (230–400 mesh, Merck). TLC and prep. TLC: precoated silica gel plates 60 F₂₅₄ (0.25 and 0.5

mm). HPLC: column, ODS (Cosmosil 10C₁₈); solvent, H₂O–MeOH (5 ml min⁻¹); detection, 230 nm.

Plant materials. Plant material, *S. byzantina*, was collected in the suburbs of Tokat, Turkey, on 4 July 1991 and identified by the authors (G. H. and E. S.). Voucher specimens (91D 106) are deposited in the Herbaria of the Faculty of Pharmaceutical Sciences, Kyoto University, and Faculty of Pharmacy, Gazi University.

Isolation. Dried aerial parts (4.8 kg) of *S. byzantina* were extracted with MeOH (47 l) for 2 weeks at room temp. The plant was extracted again with the same vol. of MeOH for 3 weeks. The combined MeOH extracts were evapd *in vacuo*. The residue was dissolved in 90% MeOH (1.1 l), and the soln was washed with *n*-hexane (1 × 3). The aq. MeOH soln was concd *in vacuo*. The residue was suspended in H₂O (1 l) and partitioned with EtOAc (1 × 3). The aq. layer was extracted with *n*-BuOH (1 × 3). After being washed with H₂O, the *n*-BuOH extract was evapd *in vacuo* to give a residue (32.4 g). The residue was subjected to CC over silica gel (1.2 kg) with CHCl₃–MeOH as eluent with increasing MeOH content. 2 l each of CHCl₃, CHCl₃–MeOH (19:1), CHCl₃–MeOH (93:7), CHCl₃–MeOH (9:1), CHCl₃–MeOH (17:3), CHCl₃–MeOH (4:1) and CHCl₃–MeOH (7:3) were passed successively through the column and 500-ml frs were collected. Fr. 9 gave a residue (1.027 g) on evapn which was repeatedly sepd by HPLC (MeOH–H₂O, 9:11, then 7:13) to give icaraside B₂ [1] (15.4 mg), 3 (35.8 mg) and 4 [2] (25.8 mg). Fr. 10 gave a residue (860 mg) on evapn which was sepd by HPLC (MeOH–H₂O, 7:13, then 1:3) to give 1 (9.9 mg), (6*R*, 9*S*)-3-oxo- α -ionol glucoside (4.6 mg) [3] and (6*R*, 9*R*)-3-oxo- α -ionol glucoside (10.5 mg) [3]. Fr. 16 gave a residue (1.85 g) which was repeatedly sepd by HPLC (MeOH–H₂O, 9:11, then 7:13) to give verbascoside (31.2 mg) [4]. The known compounds were identified by comparisons of spectral data with those reported.

Byzantionoside A (1). Amorphous powder, $[\alpha]_{\text{D}}^{24} + 74.9^{\circ}$ (MeOH, c 0.44), UV $\lambda_{\text{max}}^{\text{MeOH}}$ nm (ϵ): 226.5 (8830). IR $\nu_{\text{max}}^{\text{KBr}}$ cm⁻¹: 3392, 1670, 1076. ¹H NMR (CD₃OD): δ 0.91 (3H, *s*, H₃-11), 1.04 (3H, *s*, H₃-12), 1.57 (1H, *dd*, $J = 13.7, 5.9$ Hz, H₂-2), 1.63 (3H, *br s*, H₃-13), 1.83 (1H, *dd*, $J = 13.7, 6.4$ Hz, H₂-2), 2.26 (3H, *s*, H₃-10), 2.56 (1H, *d*, $J = 9.8$ Hz, H-6), 3.15 (1H, *dd*, $J = 9.5, 8.5$ Hz, H-2'), 3.67 (1H, *m*, H₁-6'), 3.87 (1H, *br d*, $J = 11.7$ Hz, H₁-6'), 4.36 (1H, *m*, H-



1 R=H
2 R=Ac

3 9*S* ; R=H
4 9*R* ; R=H
5 9*S* ; R=Ac

3), 4.39 (1H, *d*, *J* = 7.8 Hz, H-1'), 5.74 (1H, *d*, *J* = 1.5 Hz, H-4), 6.12 (1H, *d*, *J* = 15.6 Hz, H-8), 6.67 (1H, *dd*, *J* = 15.6, 9.8 Hz, H-7). CD $\Delta \epsilon_{239} + 8.04$ (MeOH, 2.67×10^{-5} M). Negative ion FABMS *m/z* 369.1907 [M-H]⁻ (C₁₉H₂₉O₇ requires 369.1913).

Byzantionoside B (3). Amorphous powder, $[\alpha]_D^{24} + 48.0^\circ$ (MeOH, *c* 1.53). UV $\lambda_{\max}^{\text{MeOH}}$ nm (ϵ): 240 (12500). IR ν_{\max}^{KBr} cm⁻¹: 3368, 1648. ¹H NMR (CD₃OD): δ 1.01, 1.09 (each 3H, *s*, H₃-11, H₃-12), 1.18 (3H, *d*, *J* = 6.4 Hz, H₃-10), 1.50 (1H, *m*), 1.64 (2H, *m*), 1.96 (2H, *m*), 1.97 (1H, *d*, *J* = 17.6 Hz, H₁-2), 2.05 (3H, *d*, *J* = 1.5 Hz), 2.46 (1H, *d*, *J* = 17.6 Hz, H₁-2), 3.14 (1H, *dd*, *J* = 8.8, 7.8 Hz, H-2'), 3.64 (1H, *dd*, *J* = 11.7, 5.4 Hz, H₁-6'), 3.87 (2H, H-9, H₁-6'), 4.32 (1H, *d*, *J* = 7.8 Hz, H-1'), 5.80 (1H, *br s*, H-4). ¹³C NMR: Table 1. CD $\Delta \epsilon_{237} + 3.28$ (MeOH, 2.49×10^{-3} M). Negative ion FABMS *m/z*: 371.2068 [M-H]⁻ (C₁₉H₃₁O₇ requires 371.2070).

Byzantionoside A tetraacetate (2). Compound **1** (2.2 mg) was acetylated with a mixt. of Ac₂O (0.5 ml) and pyridine (0.5 ml) for 18 h at room temp. Excess of MeOH was added to the mixt. and the soln was concd *in vacuo* to give a residue which was purified by prep. TLC (Et₂O) to give **2** (2.3 mg). ¹H NMR (CDCl₃): δ 0.88 (3H, *s*, H₃-11), 0.97 (3H, *s*, H₃-12), 1.44 (1H, *dd*, *J* = 14.2, 4.9 Hz, H₂-2), 1.62 (1H, *br s*, H₃-13), 1.71 (1H, *dd*, *J* = 14.2, 6.1 Hz, H₂-2), 2.00 (3H), 2.03 (6H), 2.10 (3H) (each *s*, 4 × OAc), 2.25 (3H, *s*, H₃-10), 2.45 (1H, *d*, *J* = 9.8 Hz, H-6), 3.69 (1H, *m*, H-5'), 4.16 (1H, *dd*, *J* = 12.2, 2.7 Hz, H₁-6'), 4.22 (1H, *m*, H-3), 4.26 (1H, *dd*, *J* = 12.2, 5.4 Hz, H₁-6'), 4.62 (1H, *d*, *J* = 8.1 Hz, H-1'), 4.97 (1H, *dd*, *J* = 9.8, 8.1 Hz, H-2'), 5.08 (1H, *dd*, *J* = 9.8 and 9.8 Hz, H-4'), 5.21 (1H, *dd*, *J* = 9.8, 9.8 Hz, H-3'), 5.66 (1H, *m*, H-4), 6.08 (1H, *d*, *J* = 15.9 Hz, H-8), 6.52 (1H, *dd*, *J* = 15.9, 9.8 Hz, H-7). Negative ion FABMS *m/z*: 537.2298 [M-H]⁻ (C₂₇H₃₇O₁₁ requires 537.2336).

Byzantionoside B tetraacetate (5). Compound **3** (2.5 mg) was acetylated and purified as before to give **5** (2.3 mg) as an amorphous powder. ¹H NMR (CDCl₃): δ 1.01, 1.04 (each 3H, *s*, H₃-11 and 12), 1.11 (3H, *d*, *J* = 6.4 Hz, H₃-10), 1.40 (1H, *m*), 1.55 (1H, *m*), 1.86 (1H, *m*), 1.98 (3H, *d*, *J* = 1.0 Hz, H₃-13), 2.01 (3H),

2.03 (6H), 2.07 (3H) (each *s*, 4 × OAc), 2.36 (1H, *d*, *J* = 17.1 Hz, H₁-2), 3.36 (1H, *m*, H-5'), 3.69 (1H, *m*, H-9), 4.12 (1H, *dd*, *J* = 12.2, 2.4 Hz, H₁-6'), 4.23 (1H, *dd*, *J* = 12.2, 4.9 Hz, H₁-6'), 4.53 (1H, *d*, *J* = 7.8 Hz, H-1'), 4.94 (1H, *dd*, *J* = 9.8, 7.8 Hz, H-2'), 5.08 (1H, *dd*, *J* = 9.8, 9.8 Hz, H-4'), 5.20 (1H, *dd*, *J* = 9.8, 9.8 Hz, H-3'), 5.82 (1H, *br s*, H-4). Negative ion FABMS *m/z*: 539.2506 [M-H]⁻ (C₂₇H₃₉O₁₁ requires 539.2493).

Acknowledgements—The authors thank the Cooperative Centre of the University of Tokushima for opportunities to record NMR spectra. One of the authors (H. Z.) is grateful to the Gohou Life Sciences International Fund for financial support. This work was supported in part by a grant-in-aid from the Ministry of Education, Science, Sports and Culture, Japan (No. 02041048).

REFERENCES

- Miyase, T., Ueno, A., Takizawa, N., Kobayashi, H. and Karasawa, H., *Chemical and Pharmaceutical Bulletin*, 1987, **35**, 1109.
- Miyase, T., Ueno, A., Takizawa, N., Kobayashi, H. and Oguchi, H., *Chemical and Pharmaceutical Bulletin*, 1988, **36**, 2475.
- Pabst, A., Barron, D., Semon, E. and Schreier, P., *Phytochemistry*, 1992, **31**, 1649.
- Birkofer, L., Kaiser, C. and Thomas, U., *Zeitschrift Naturforschung*, 1968, **23B**, 1051.
- Otsuka, H., Yao, M., Kamada, K. and Takeda, Y., *Chemical and Pharmaceutical Bulletin*, 1995, **43**, 754.
- Buchecker, R., Hamm, P. and Eugster C. H., *Helvetica Chimica Acta*, 1974, **57**, 631.
- Otsuka, H., Kamada, K., Ogimi, C., Hirata, E., Takushi, A. and Takeda, Y., *Phytochemistry*, 1994, **35**, 1331.
- Takeda, Y., Zhang, H., Matsumoto, T., Otsuka, H., Oiso, Y., Honda, G., Tabata, M., Fujita, T., Sun, H., Sezik, E. and Yesilada, E., *Phytochemistry*, 1997, **44**, 117.