

Development of 11 Polymorphic Microsatellite Loci from *Primula amethystina* Franchet. (Primulaceae)

Yuan Huang

Key Laboratory of Biodiversity and Biogeography, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650204, Yunnan, China

Xue-qin Wang and Chun-yan Yang

Key Laboratory of Biodiversity and Biogeography, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650204, Yunnan, China; and Graduate School, Chinese Academy of Sciences, Beijing 100049, China

Chun-lin Long¹

Key Laboratory of Biodiversity and Biogeography, Kunming Institute of Botany, Chinese Academy of Sciences, Lanhei Road 134, Heilongtan, Kunming 650204, Yunnan, China; and College of Life and Environmental Sciences, Minzu University of China, Beijing 100081, China

Additional index words: *Primula amethystina*, microsatellite, genetic diversity

Abstract. *Primula amethystina* Franchet, is a beautiful perennial herbaceous plant locally endemic to the alpine area in southwest China. We isolated and characterized 11 polymorphic microsatellite primer pairs from this species. The number of alleles ranged from two to five. The observed and expected heterozygosities (H_o and H_e) were 0.25 to 0.875 and 0.223 to 0.691, respectively. Six loci were significantly deviated from Hardy-Weinberg equilibrium as a result of the heterozygote deficiency. These markers will have great potential to reveal the genetic population structure and genetic diversity of *P. amethystina*.

Primula amethystina Franchet, is a beautiful perennial herbaceous plant locally endemic to the alpine area in southwest China. It is widely distributed in alpine meadows at altitudes of 3400 to 5000 m in west and northwest Yunnan, southwest Sichuan, over the border into Tibet (Hu and Kebo, 1996; Richards, 2002). *P. amethystina* subsp. *amethystina* is confined to the Cangshan Mountains, Dali, west Yunnan, whereas *P. amethystina* subsp. *brevifolia* widely spreads through the range. Frequently, *P. amethystina* subsp. *brevifolia* is more robust than *P. amethystina* subsp. *amethystina* with up to 20 flowers borne on stems to 25 cm. To classify these two species and study the speciation mode of them, their genetic structures need to be understood. Here, 11 polymorphic micro-

satellite loci of *P. amethystina* were developed as potential tools to investigate the genetic structures of these species.

Genomic DNA was extracted from leaf tissues using the cetyltrimethyl ammonium bromide (CTAB) method (Mulligan, 1993). The isolation of microsatellite loci was performed according to the fast isolation of microsatellite by amplified fragment length polymorphism of sequences containing repeats (FASCO) (Zane et al., 2002). Approximately 300 ng genomic DNA was completely digested with *Msp*I restriction enzyme (Fermentas). The digested DNA was ligated to *Msp*I adapter pair (Vos et al., 1995), then 5 μl of the adapter-ligated fragments acted as templates to perform polymerase chain reaction (PCR) in a volume of 20 μl, using *Msp*-N (5'-GATGGAOTCTTGAGTAAN-3') as a primer following the program: 95 °C for 3 min, 30 cycles of 94 °C for 30 s, 53 °C for 60 s, 72 °C for 60 s followed by an elongation step of 5 min at 72 °C.

For enrichment, the PCR products were denatured at 95 °C for 5 min and then hybridized with a 5'-biotinylated probe (A01) in 250 μl hybridization solution (20 × SSC, 10% SDS, 100 pmol/μl probe) at 48 °C for 2 h. The DNA hybridized to the probe was separated and captured by streptavidin-coated magnetic beads at room temperature for 20 min followed by two washing steps, including three times in TEN_{low} for 15 min and three times in TEN_{low} for 24 min. The

separated single-stranded DNA was subjected to a second round of PCR according to the same procedure as the first round of PCR. The PCR products, after being purified with the E.Z.N.A Gel Extraction Kit (Oncor, Bio-Tek, Atlanta, GA), were ligated into pCR-TOPO-T vector (Takara, Otsu, Shiga, Japan) according to the manufacturer's instructions and then transformed into *Escherichia coli* strain JM109 (Sangon, Shanghai, China). The positive clones were picked out by blue-white screening and tested by PCR using (A/G) and M13⁺/M13⁻ as primers, respectively. Of the 91 clones sequenced, 79 contained potential microsatellite motifs and 42 with unique microsatellites were selected for designing primers using Primer 5.0 (Clarke and Gorley, 2001).

A total of 12 primer sets were abandoned as a result of amplification of multiple bands or unsuccessful amplification of target fragments, and the remaining 30 primer pairs were tested for polymorphisms across 24 individuals from a population of *P. amethystina* subsp. *amethystina* and three populations of *P. amethystina* subsp. *brevifolia* (Table 1). PCR reaction was done in a 20-μl volume using a PTC220 thermal cycler (MJ Research, Ashland, OR). Each reaction was performed using 20 ng of genomic DNA, 1 μM of each dNTP, 1 μM of each primer, 1× Taq buffer [100 mM Tris-HCl, pH 8.8, 2.0 mM MgCl₂, 200 mM (NH₄)₂SO₄, 0.1% Tween 20], and 1 U of Taq polymerase (Takara). The PCR programs are as follows: initial denaturing step at 95 °C for 3 min, 30 cycles of 94 °C for 30 s, primer-specific annealing temperature 55 to 62 °C for 30 s, 72 °C for 30 s, and a final extension step at 72 °C for 8 min. The PCR products were electrophoresed in denaturing 6% polyacrylamide gels using a 25-μl DNA ladder molecular size standard (Fermentas, Ontario, Canada) to estimate allele sizes by silver staining.

Of the 30 new primers, 11 of them displayed polymorphisms across populations. The number of alleles per locus, observed (H_o) and expected heterozygosity (H_e), and deviation from Hardy-Weinberg equilibrium (HWE) were assessed using GENEPOP Version 3.4 (<http://www1.zoo.ucl.ac.be/genepop/>) (Raymond and Rousset, 1995). The number of alleles per locus ranged from two to five with an average of 2.8 (Table 2). The observed and expected heterozygosities (H_o and H_e) ranged from 0.25 to 0.875 and from 0.223 to 0.691 with averages of 0.368 and 0.442, respectively (Table 2). Among the 11 microsatellite markers, six loci showed significant deviation from HWE ($P < 0.01$) (Table 2), which was the result of deficiency of heterozygotes or the limitation of sample size. Tests for linkage disequilibrium were run in FSTAT Version 2.9.3.2 (Goudet, 1995). Significance levels were adjusted using sequential Bonferroni corrections (Rice, 1989). No loci provided significant linkage disequilibrium after Bonferroni correction. These polymorphic microsatellite loci presented here would provide a useful tool for studying the pop-

Received for publication 14 Sept. 2009. Accepted for publication 21 Oct. 2009.

This work was supported by a project of "Light of Western China" (20080338137W), the National Natural Science Foundation (30870170), the Ministry of Science and Technology of China (2008FY110402-2-2 and 2008DFA21008), the Ministry of Education of China (20080444 and MUC 985-1-3), and the Japan Society for the Promotion of Science (JSPS-AP1099000).

Dr. Zhiyan Wu made contributions in collecting samples from the field.

To whom reprint requests should be addressed; e-mail: longlin@mail.kib.ac.cn.

10.1. Search interface of *P. aeruginosa* phage particles and *P. aeruginosa* ciliop. lysoph.

Station	Species	Location	Position	Altitude (m)	Accession no.
D	<i>P. amabilis</i> subsp. <i>brevifolia</i>	Laijia Shan, Liping, Yunnan province	N $26^{\circ}54'40.5''$ E $105^{\circ}17'45.6''$	3500	Wu20090617
C	<i>P. amabilis</i> subsp. <i>brevifolia</i>	Tianshi, Xanggong, Yunnan province	N $27^{\circ}37'44.3''$ E $105^{\circ}37'10.5''$	3800	Wu20090620
B	<i>P. amabilis</i> subsp. <i>amabilis</i>	Chang dan, Dali, Yunnan province	N $25^{\circ}46'17.4''$ E $106^{\circ}03'33.8''$	3900	Wu20090621
A	<i>P. amabilis</i> subsp. <i>brevifolia</i>	Changhui, Muli, Sichuan province	N $27^{\circ}56'12.4''$ E $101^{\circ}11'13.3''$	3400	Wu20090609

Table 2. Characteristics of 111 endocrinologic interventions for *Pomacanthus semicirculatus* females.

Locs (GenBank no.)	Repeat motif	Primer sequence (5'-3')	Allele size range (bp)	A	F _{ST} (%)	R ₀	N ₀	HWE (P value)
0010487	(AG) _n	R: CTCAATACGAACTTAAAGG F: ATCTTCAGTTGGAAAGAAA	291	3	33	0.75	0.606	0.0001**
0010488	(AT) _n (AC) _n (AC) _n	R: CGGAAACCGGAAACTTCTAC F: CCACATACCTTGACCTGACCT	285	2	33	0.292	0.254	1
0010489	(GA) _n (GA) _n (AG) _n	R: TAGAAGGTGTTAGGTATTAGG F: ATATGGCTCTTCAGTAGAAA	265	2	33	0.875	0.500	0.0002**
0010490	(AG) _n (AC) _n (AG) _n	R: ATATTCACCCGAACTTCTTGG F: ACACGTTTACTTTTCAGAG	253	3	34	0.75	0.494	0.0007**
0010491	(CT) _n (CT) _n (CT) _n (CT) _n	R: CTCTGGGAACTATTCATTTT F: ACACGCTCTTCTCATAAGTC	251	3	34	0.792	0.554	0.0189
0010492	(CT) _n	R: ACCCTAGAGTCATTTCAGGC F: GTTACGCCCTCTATCACCTC	236	3	68	0.333	0.439	0.0001**
0010493	(CTT) _n (CT) _n (CT) _n (CT) _n	R: TACAAACAACCTTCTTCCTC F: AGGAAAGTGTACCTTCCCAAAT	223	3	53	0.333	0.264	1
0010494	(GA) _n	R: TTGAGTTGTTAACTCTTCTGG F: GGTAAAGGTTAAAGGAGGTAA	198	2	56	0.25	0.223	1
0010495	(AG) _n	R: AAACGTGATACCCATAGAAC F: CATAGATGATGTTACGCGTGAD	190	4	53	0.625	0.491	0.0072*
0010496	(CT) _n (CT) _n	R: TTGTGTTGTTGGAAAGAGTA F: TGGGAACTGGTCAAGTTTATT	186	2	58	0.25	0.223	1
0010497	(GA) _n (AG) _n (GA) _n (GA) _n	R: AACAAACAAACAAACCGGAACT F: TGGGAACTGGTCAAGTTTATT	186	3	53	0.333	0.390	0.00001**

The annealing temperature of primer pair A = number of alleles; H_o = observed heterozygosity; H_e = expected heterozygosity. Statistically significant deviation from Hardy-Weinberg proportions ($P < 0.05$) is indicated by * or **.

avian genetic structure and genetic diversity

Literature Review