


A Pair of Novel Heptentriol Stereoisomers from the Ascomycete *Daldinia concentrica*

by Fei Wang and Ji-Kai Liu*

Kunming Institute of Botany, the Chinese Academy of Sciences, Kunming 650204, P. R. China
(e-mail: jkliu@mail.kib.ac.cn)

A pair of novel heptentriol stereoisomers, hept-6-ene-2,4,5-triols **2** and **3**, were isolated from the culture broth of the ascomycete *Daldinia concentrica* (BOLTON : FRIES) CESATI & DE NOTARIS, besides three known compounds, *i.e.*, 2,3-dihydro-5-hydroxy-2-methyl-4*H*-1-benzopyran-4-one (**1**), 3,5-dihydroxy-2-(1-oxobutyl)-cyclohex-2-en-1-one (**4**), and pyroglutamic acid (= 5-oxo-L-proline; **5**). Their structures were determined by spectroscopic means, including 2D-NMR (HMQC, HMBC, ¹H,¹H-COSY).

Introduction. – Fungi of the ascomycete genus *Daldinia* (Xylariaceae) have been shown to be a good source of bioactive secondary metabolites. Several decades ago, [1,1'-binaphthalene]-4,4',5,5'-tetrol and dihydroxyperylenequinone were reported from the stromata of *Daldinia* sp. [1], while the same authors later reported 2,6-dihydroxybutyrophenone (= 1-(2,6-dihydroxyphenyl)butan-1-one), 8-methoxynaphthalen-1-ol, and 5-hydroxy-2-methylchromone (= 5-hydroxy-2-methyl-4*H*-1-benzopyran-4-one) from its mycelia [2]. Some of these compounds were later found to show antimicrobial and nematocidal activities [3]. More recently, the squalene-type triterpenoids concentricols A and B–D were isolated from the fruiting bodies of the fungus by *Stadler* and co-workers [4] and *Asakawa* and co-workers [5], respectively. As part of our search for bioactive metabolites from higher fungi in Yunnan Province, China [6–10], the culture broth of *D. concentrica* was investigated. This report deals with the isolation and structure elucidation of a pair of novel heptentriol stereoisomers, hept-6-ene-2,4,5-triols **2/3**, from the AcOEt extract of the culture broth of *D. concentrica*.



Results and Discussion. – Compound **2** was obtained as an oil. The molecular formula of **2** was determined to be C₇H₁₄O₃ by the ¹³C-NMR (DEPT; 1 Me, 1 CH₂, 3 CH–O, 1 CH=, 1 CH₂=) and FAB-MS (pos.) data ([*M* + 1]⁺ at *m/z* = 147). The IR (neat) absorptions at 3418 (br.), 3082, 3021, 1645, 992, and 923 cm⁻¹ indicated the presence of an OH and CH=CH₂ group, and the ¹H-NMR (*d* at δ 1.13 (*J* = 6.0, Me(1);

see Table) and the HMBC data (cross-peak $H-C(1)/C(3)$, see Fig.) that of a $\text{MeCH}(\text{OH})\text{CH}_2$ moiety. From these evidences and detailed $^1\text{H}, ^1\text{H}$ -COSY analysis, the structure of **2** was elucidated as hept-6-ene-2,4,5-triol.




Figure. Significant HMBC correlations of **2/3**

Table. ^1H - (500 MHz) and ^{13}C -NMR (125 MHz) Data of **2** and **3**. Solvent $\text{CDCl}_3/\text{CD}_3\text{OD}$ 2 : 1; δ in ppm, J in Hz.

2			3		
	$\delta(\text{C})(\text{DEPT})$	$\delta(\text{H})$	$\delta(\text{C})(\text{DEPT})$	$\delta(\text{H})$	
Me(1)	21.0 (Me)	1.13 (<i>d</i> , $J = 6.1$)	20.1 (Me)	1.01 (<i>d</i> , $J = 6.2$)	
H-C(2)	73.6 (CH)	3.79 (<i>m</i>)	74.2 (CH)	4.03 (<i>m</i>)	
$\text{CH}_2(3)$	42.0 (CH_2)	1.35, 2.20 (<i>2m</i>)	41.1 (CH_2)	1.39 (<i>m</i>), 1.66 (<i>ddd</i> , $J = 13.0, 5.4, 2.3$)	
H-C(4)	73.2 (CH)	4.08 (<i>m</i>)	76.0 (CH)	3.80 (<i>m</i>)	
H-C(5)	83.9 (CH)	3.91 (<i>m</i>)	87.2 (CH)	3.90 (<i>m</i>)	
H-C(6)	133.6 (CH)	5.76 (<i>m</i>)	136.7 (CH)	5.56 (<i>ddd</i> , $J = 17.1, 10.4, 6.3$)	
$\text{CH}_2(7)$	117.9 (CH_2)	5.09 (<i>dt</i> , $J = 10.1, 1.6$), 5.16 (<i>dt</i> , $J = 16.6, 1.6$)	115.4 (CH_2)	4.87 (<i>dt</i> , $J = 10.4, 1.4$), 5.04 (<i>dt</i> , $J = 17.1, 1.4$)	
OH		3.99 (<i>br. s</i>)		4.20 (<i>br. s</i>)	

Compound **3** was also obtained as an oil. It possessed the same FAB-MS as **2**, and its ^1H - and ^{13}C -NMR spectra (see Table) were also very similar to those of **2**. In view of the perfect accordance of the IR spectrum and the key HMBC correlations with those of **2**, the structure of **3** was deduced to have the same chemical structure as that of **2**, *i.e.*, that of a hept-6-ene-2,4,5-triol. Considering the obvious differences of **2** and **3** with respect to their optical rotation ($[\alpha]_{\text{D}} = +21.7$ and -52.2 , resp. ($c = 0.75$, CHCl_3)) and behavior on TLC (silica gel, $\text{CHCl}_3/\text{MeOH}$ 150 : 1; R_{f} 0.75 and 0.50, resp.), they were established as being a pair of stereoisomers. The determination of their absolute configuration at C(2), C(4), and C(5) is in progress and will be reported in the future.

Comparison of the physicochemical properties with reported data allowed to identify the compounds **1**, **4**, and **5**, isolated from the same fungus, as 2,3-dihydro-5-hydroxy-2-methyl-4*H*-1-benzopyran-4-one [2][11], 3,5-dihydroxy-2-(1-oxobutyl)cyclohex-2-en-1-one [12], and pyroglutamic acid (= 5-oxo-L-proline) [13], respectively. Compound **1** was reported as a metabolite from the rice culture solution of the fungus *Phialophora gregata* and shown to have biological activity against soybean cells [11]. Compound **4** has also previously been isolated from the culture broth of the fungus *Nodulisporium* sp. and found to have chlorosis activity, which was stronger against monocotyledons than against dicotyledons [12].

This project was supported by the National Natural Science Foundation of China (30225048).

Experiment Part

General. CC = Column chromatography. TLC: visualization by heating silica-gel plates sprayed with 10% H₂SO₄ in EtOH. Optical rotations: *Horiba SEPA-300* digital polarimeter. IR Spectra: *Bruker Tensor-27* spectrometer; $\tilde{\nu}$ in cm⁻¹. ¹H- and ¹³C-NMR Spectra: *Bruker DXR-500* spectrometer; δ in ppm, J in Hz. MS: *VG Autospec-3000* spectrometer; m/z (rel. int.).

Mushroom Material and Culture. The fungus *D. concentrica* was collected at the Botanic Garden of the Kunming Institute of Botany, the Chinese Academy of Sciences (CAS), P. R. China, in July 2003, and identified by Prof. *Mu Zang*, Kunming Institute of Botany. The voucher specimen was deposited in the Herbarium of the Kunming Institute of Botany. Culture medium: potato (peeled) 200 g, glucose 20 g, KH₂PO₄ 3 g, MgSO₄ 1.5 g, citric acid 0.1 g, and thiamin hydrochloride 10 mg in 1 l of deionized H₂O (pH 6.5 before autoclaving). The culture soln. was fermented at 22° for 10 days on a rotary shaker (150 rpm).

Extraction and Isolation. The culture broth (15 l) was filtered and then successively extracted with AcOEt. The AcOEt extract was evaporated and the oily residue (2.8 g) subjected to CC (silica gel, gradient CHCl₃/MeOH). *Fr. A* (eluted with pure CHCl₃) was further purified by CC (silica gel): **1** (23 mg). *Fr. B* (eluted with CHCl₃/MeOH 99 : 1) was further purified by CC (silica gel): pure **2** (28 mg), a crude oil (*Fr. B.1*) containing **3**, and a crude oil (*Fr. B.2*) containing **4**. *Fr. B.1* was further purified by repeated CC (silica gel, CHCl₃/MeOH 120 : 1): pure **3** (23 mg). *Fr. B.2* was further purified by prep. TLC (CHCl₃/MeOH 98 : 2; detection by UV light (254 nm)): pure **4** (6.7 mg). *Fr. C* (eluted with CHCl₃/MeOH 85 : 15; see above) was subjected to CC (silica gel): **5** (54 mg).

(2*E*)-2,3-Dihydro-5-hydroxy-2-methyl-4H-1-benzopyran-4-one (**1**): Yellow needles. M.p. 30–35° (hexane). ¹H-NMR (CDCl₃): 11.67 (*s*, OH); 7.33 (*t*, $J = 8.3$, H–C(7)); 6.48 (*dd*, $J = 8.3, 0.7$, H–C(6)); 6.41 (*dd*, $J = 8.3, 0.7$, H–C(8)); 4.56 (*m*, H–C(2)); 2.73 (*dd*, $J = 17.1, 12.0$, H _{α} –C(3)); 2.66 (*dd*, $J = 17.1, 4.0$, H _{β} –C(3)); 1.50 (*d*, $J = 6.0$, Me–C(2)). ¹³C-NMR (CDCl₃): 198.4 (*s*, C=O); 162.2 (*s*, C(5)); 161.7 (*s*, C(8a)); 138.1 (*d*, C(7)); 109.2 (*d*, C(6)); 108.1 (*s*, C(4a)); 107.3 (*d*, C(8)); 73.8 (*d*, C(2)); 43.9 (*t*, C(3)); 20.8 (*q*, Me–C(2)). EI-MS: 178 (53, *M*⁺, C₁₀H₁₀O₃⁺), 163 (24), 136 (52), 108 (39), 80 (8).

Hept-6-ene-2,4,5-triol (**2**): Oil. *R*_f (CHCl₃/MeOH 150 : 1) 0.75. $[\alpha]_{\text{D}}^{25} = +21.7$ ($c = 0.75$, CHCl₃). IR (neat): 3418, 3082, 3021, 2967, 2927, 1645, 1445, 1384, 1078, 992, 923. ¹H- and ¹³C-NMR: *Table*. FAB-MS (pos.): 147 ([*M* + 1]⁺, C₇H₁₅O₃⁺).

Hept-6-ene-2,4,5-triol (**3**): Oil. *R*_f (CHCl₃/MeOH 150 : 1) 0.50. $[\alpha]_{\text{D}}^{26} = -52.2$ ($c = 0.75$, CHCl₃). IR (neat): 3418, 3093, 3009, 2973, 2930, 1644, 1443, 1384, 1106, 990, 926. ¹H- and ¹³C-NMR: *Table*. FAB-MS (pos.): 147 ([*M* + 1]⁺, C₇H₁₅O₃⁺).

3,5-Dihydroxy-2-(1-oxobutyl)cyclohex-2-en-1-one (**4**): Oily solid. ¹H-NMR (CDCl₃): 4.38 (*m*, H–C(5)); 2.98 (*t*, $J = 7.4$, CH₂(2'')); 2.92 (*dd*, $J = 17.9, 4.0$, H _{α} –C(6)); 2.77 (*ddd*, $J = 17.9, 5.8, 1.3$, H _{β} –C(6)); 2.74 (*dd*, $J = 16.5, 3.6$, H _{α} –C(4)); 2.62 (*ddd*, $J = 16.5, 6.5, 1.3$, H _{β} –C(4)); 1.63 (*sext.*, $J = 7.4$, CH₂(3'')); 0.96 (*t*, $J = 7.4$, Me(4')). ¹³C-NMR (CDCl₃): 205.4 (*s*, C(1'')); 196.0 (*s*, C(1)); 193.0 (*s*, C(3)); 112.9 (*s*, C(2)); 63.4 (*d*, C(5)); 47.1 (*t*, C(4)); 42.2 (*t*, C(2'')); 41.5 (*t*, C(6)); 18.0 (*t*, C(3'')); 13.9 (*q*, C(4')). FAB-MS (neg.): 197 (100, [*M* – 1]⁻, C₁₀H₁₃O₄⁻).

(2*S*)-Pyroglutamic acid (= (2*S*)-5-Oxopyrrolidine-2-carboxylic Acid = 5-Oxo-L-proline; **5**): Yellowish needles. M.p. 155–157° (MeOH). ¹H-NMR (CD₃OD): 7.88 (*s*, NH); 4.24 (*dd*, $J = 9.0, 4.6$, H–C(2)); 2.48 (*m*, H _{α} –C(3)); 2.32 (*m*, CH₂(4)); 2.16 (*m*, H _{β} –C(3)). ¹³C-NMR (CD₃OD): 181.1 (*s*, CONH); 175.8 (*s*, COOH); 57.0 (*d*, C(2)); 30.4 (*t*, C(4)); 26.0 (*t*, C(3)). EI-MS: 129 (9, *M*⁺, C₅H₇NO₃⁺), 100 (4), 84 (100), 73 (5), 56 (12). FAB-MS (neg.): 128 (100, [*M* – 1]⁻).

REFERENCES

- [1] D. C. Allport, J. D. Bu'lock, *J. Chem. Soc.* **1958**, 4090.
- [2] D. C. Allport, J. D. Bu'lock, *J. Chem. Soc.* **1960**, 654.
- [3] H. Anke, M. Stadler, A. Mayer, O. Sterner, *Can. J. Bot.* **1995**, 73, 802.
- [4] M. Stadler, M. Baumgartner, T. Grothe, A. Mühlbauer, S. Seip, H. Wollweber, *Phytochemistry* **2001**, 56, 787.
- [5] D. N. Quang, T. Hashimoto, M. Tanaka, M. Baumgartner, M. Stadler, Y. Asakawa, *Phytochemistry* **2002**, 61, 345.
- [6] J. M. Gao, L. Hu, J. K. Liu, *Steroids* **2001**, 66, 771.
- [7] J. M. Gao, L. Hu, Z. J. Dong, J. K. Liu, *Lipids* **2001**, 36, 521.
- [8] J. K. Liu, *Heterocycles* **2002**, 57, 157.

- [9] J. K. Liu, J. W. Tan, Z. J. Dong, X. H. Wang, P. G. Liu, *Helv. Chim. Acta* **2002**, *85*, 1439.
- [10] J. W. Tan, Z. J. Dong, L. Hu, J. K. Liu, *Helv. Chim. Acta* **2003**, *86*, 307.
- [11] L. E. Gray, H. W. Gardner, D. Weisleder, M. Lieb, *Phytochemistry* **1999**, *50*, 1337.
- [12] M. Igarashi, Y. Tetsuka, Y. Mimura, A. Takahashi, T. Tamamura, K. Sato, *J. Antibiot.* **1993**, *46*, 1843.
- [13] D. A. Parrish, L. J. Mathias, *J. Org. Chem.* **2002**, *67*, 1820.

Received April 13, 2004