

Mulberrofuran G and Isomulberrofuran G from *Morus alba* L.: Anti-hepatitis B Virus Activity and Mass Spectrometric Fragmentation

Chang-An Geng,[†] Yun-Bao Ma,[†] Xue-Mei Zhang,[†] Shu-Ying Yao,[†] Duo-Qing Xue,[†] Rong-Ping Zhang,[‡] and Ji-Jun Chen^{*,†}

[†]State Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, Yunnan 650201, People's Republic of China

[‡]Kunming Medical University, Kunming, Yunnan 650500, People's Republic of China

S Supporting Information

ABSTRACT: Mulberrofuran G (**1**) and isomulberrofuran G (**2**), a pair of isomeric Diels–Alder-type adducts, were isolated from the root bark of *Morus alba* L. Isomulberrofuran G (**2**) as a new IIB-type Diels–Alder-type adduct, was elucidated by extensive ¹H, ¹³C, and two-dimensional (2D) nuclear magnetic resonance (NMR) and mass spectrometry (MS) spectroscopic analyses. A fragmentation study on compounds **1** and **2** was performed by high-resolution electrospray ionization (ESI) multistage tandem mass spectrometry linked with ion-trap (IT) and time-of-flight (TOF) mass analyzers (ESI–MSⁿ/IT–TOF) in negative mode, which resulted in obviously different fragmentations. In the MS² experiments, the characteristic ions at *m/z* 451 and 439 could be revealed as their respective diagnostic ions. Mulberrofuran G (**1**) showed moderate activity, inhibiting hepatitis B virus (HBV) DNA replication with the IC₅₀ value of 3.99 μM, according to the anti-HBV assay on the HepG 2.2.15 cell line *in vitro*.

KEYWORDS: *Morus alba*, isomulberrofuran G, Diels–Alder-type adducts, MSⁿ fragmentation study, LCMS–IT–TOF, anti-HBV activity

■ INTRODUCTION

Morus alba L. is a useful plant widely cultivated in China for thousands of years, whose leaves are indispensable food for silkworms and fruits (Sang-Shen in Chinese) are largely consumed in normal diets.¹ Its root bark (Mori Cortex), leaves, and twigs are all well-known traditional Chinese medicines (TCMs) documented in *Chinese Pharmacopoeia*.² In particular, Mori Cortex is famous with the Chinese name of “Sang-Bai-Pi” and widely used for anti-inflammation, anti-hypertension, hypoglycemic, and diuretic purposes.³ The earliest usage of Mori Cortex can be traced back to “Shen-Nong-Ben-Cao-Jing”, the first Chinese dispensatory, which probably appeared at the end of the third century.⁴ Mori Cortex is rich in phenolic compounds, especially for isoprenylated flavonoids and Diels–Alder-type adducts, which have aroused much interest of scientists for their complicated structures and diverse activities.^{5–8} Up to now, more than 50 Diels–Alder-type adducts with hypotensive, hypoglycemic, antioxidative, and anti-human immunodeficiency virus (HIV) activities have been isolated from Moraceous plants.^{9–14} Diels–Alder-type adducts derived from [4 + 2]cyclo-adducts of chalcones as dienophiles and dehydroprenylphenols as dienes are the first examples of natural products biosynthesized by the enzyme-controlled intermolecular Diels–Alder reaction.¹⁵ The regioselectivity of the Diels–Alder reaction, including *exo* and *endo* addition, results in all-*trans* and *cis-trans* types of Diels–Alder-type adducts. Structurally, Diels–Alder-type adducts can be classified into two types based on the phenol nuclei: (I) adducts of one chalcone and one dehydroprenylflavone and (II) adducts of one chalcone and one dehydroprenylstilbene. In the case of

type II, *para*- or *meta*- dehydroprenylstilbenes will produce two subgroups IIA and IIB. The keto carbonyl group in the chalcone part can be further cyclized with the intramolecular hydroxy groups by the ketol reaction.^{16,17} To search for more Diels–Alder-type adducts from Mori Cortex and characterize their bioactivities and mass spectrometry (MS) fragmentation regularities, this investigation was designed to isolate and determine the structure and multistage tandem mass spectrometry (MSⁿ) fragmentation of Diels–Alder-type adducts from *M. alba* and study their anti-hepatitis B virus (HBV) properties.

■ MATERIALS AND METHODS

General Experimental Procedures. One-dimensional (1D) and two-dimensional (2D) nuclear magnetic resonance (NMR) spectra were recorded on Bruker DRX-500 and AVANCE III-600 spectrometers (Bruker, Bremerhaven, Germany) with tetramethylsilane (TMS) as the internal standard. MS data were collected on Shimadzu liquid chromatography–mass spectrometry (LCMS)–ion-trap (IT)–time of flight (TOF) (Shimadzu, Kyoto, Japan). Infrared (IR) (KBr) spectra were recorded on a Bio-Rad FTS-135 spectrometer (Bio-Rad, Hercules, CA). Ultraviolet (UV) data were collected on a Shimadzu UV-2401A spectrophotometer (Shimadzu, Kyoto, Japan). Optical rotations were collected on a Jasco model 1020 polarimeter (Horiba, Tokyo, Japan). Silica gel (Makall, Qingdao, China) and Toyopearl HW-40C gel (Toson, Tokyo, Japan) were used for column chromatography.

Received: June 19, 2012

Revised: July 25, 2012

Accepted: July 27, 2012

Published: July 27, 2012




Figure 1. Structures of mulberrofuran G (1) and isomulberrofuran G (2).

Table 1. ^1H and ^{13}C NMR Data of Isomulberrofuran G (2)^a

number	δ_{H}	δ_{C}	number	δ_{H}	δ_{C}
2		156.4, s	6''a	2.05, 1H, dd, 16.5, 11.0	36.4, t
3	7.03, 1H, s	102.1, d	6''b	2.71, 1H, dd, 16.5, 5.5	
4		133.2, s	7''	1.80, 3H, brs	23.8, q
5	6.86, 1H, d, 2.0	103.9, d	8''		102.4, s
6		159.9, s	9''		117.5, s
7	6.37, 1H, t, 2.0	103.8, d	10''		157.4, s
8		159.9, s	11''	6.43, 1H, d, 2.4	104.7, d
9	6.86, 1H, d, 2.0	103.9, d	12''		160.0, s
1'		155.1, s	13''	6.21, 1H, dd, 8.8, 2.4	107.1, d
2'	7.11, 1H, s	99.3, d	14''	7.26, 1H, d, 8.8	130.5, s
3'		150.8, s	15''		117.3, s
4'		122.4, s	16''		153.5, s
5'	7.41, 1H, s	121.1, d	17''	6.37, 1H, d, 2.5	104.1, d
6'		124.6, s	18''		157.8, s
1''		135.0, s	19''	6.48, 1H, dd, 8.3, 2.5	109.9, d
2''	6.12, 1H, brd, 5.0	123.5, d	20''	7.09, 1H, d, 8.3	127.9, d
3''	3.44, 1H, dd, 5.5, 5.0	35.3, d	OH-6/OH-8	8.54, 2H, s	
4''	3.36, 1H, dd, 11.5, 5.5	36.5, d	OH-10''	8.64, 1H, s	
5''	2.88, 1H, ddd, 11.5, 11.5, 5.5	27.8, d	OH-12''	8.51, 1H, s	
			OH-18''	8.40, 1H, s	

^aData were measured in acetone-*d*₆ with δ in ppm and *J* in Hz.

Electrospray ionization (ESI)–MSⁿ spectra were acquired on a Shimadzu LCMS–IT–TOF mass spectrometer equipped with an ESI source in negative mode. The mass resolution was about 10 000 full width at half maximum (fwhm). Accurate masses were corrected by calibration using the sodium trifluoroacetate (CF₃CO₂Na) clusters. Unless specified otherwise, analytical conditions were as follows: spray voltage, –3.50 kV; detector voltage, 1.70 kV; pressure of the TOF region, 1.6×10^{-4} Pa; pressure of the IT, 1.5×10^{-2} Pa; rotary pump (RP) area vacuum, 65.5 Pa; dry gas pressure, 100.0 kPa; nebulizing gas (N₂) flow, 0.5 L/min; curved desolvation line (CDL) temperature, 200 °C; heat block temperature, 200 °C; equipment temperature, 40 °C; precursor-ion-selected width, *m/z* 3.0, selected time, 20 ms; collision-induced dissociation (CID) collision time, 20 ms; collision energy, 50%; collision gas, 50%; and *q*, 0.251. The Shimadzu composition formula predictor was used to speculate the molecular formula. Sample solutions of mulberrofuran G (1) and isomulberrofuran G (2) were prepared by dissolving each sample in a solution of acetonitrile (0.5% Et₃NH) to a final concentration of 0.1 mg/mL. The sample was introduced into the source via a syringe pump at a flow rate of 1 $\mu\text{L}/\text{min}$.

Plant Material. The root bark of *M. alba* L. (Sang-Bai-Pi) was purchased from Jü-Hua-Cun Medicinal Herb Market, Kunming, Yunnan, China, and was identified by Prof. Li-Gong Lei (Kunming Institute of Botany, Chinese Academy of Sciences). A voucher specimen (2011-01-02) was deposited in the Laboratory of Antiviral and Natural Medicine Chemistry, Kunming Institute of Botany, Chinese Academy of Sciences.

Extraction and Isolation. The powdered root bark of *M. alba* (2.0 kg) was soaked in EtOH (95%) and extracted under reflux 2 times, at

2 h for each time. The combined EtOH extracts (150.0 g) were condensed and partitioned between H₂O and EtOAc. The EtOAc part was separated by silica gel column chromatography (2.0 kg, 50.0 \times 11.0 cm inner diameter, 200–300 mesh), eluted with the CHCl₃–MeOH system (each 20 L, from 100:0 to 50:50, v/v). The CHCl₃–MeOH (90:10) eluted portion (10.0 g) was further subjected on silica gel column chromatography (100.0 g, 50.0 \times 2.5 cm inner diameter, 200–300 mesh) using a CHCl₃–Me₂CO gradient (each 1.0 L), from which the CHCl₃–Me₂CO (80:20) fraction was obtained. This fraction was purified by repeated column chromatography (silica gel and Toyopearl HW-40C gel) to yield mulberrofuran G (1, 100 mg) and isomulberrofuran G (2, 20 mg), with the purity higher than 95% (Figure 1).

Isomulberrofuran G (2). Brownish yellow powder. $[\alpha]_{\text{D}}^{24} +514.8$ (*c* 0.2, MeOH). UV (MeOH) λ_{max} (log ϵ): 335 (4.46), 322 (4.52), 296 (4.16), 284 (4.21) nm. IR (KBr) ν_{max} : 3375, 2970, 1623, 1510, 1455, 1353, 1284, 1258, 1104, 1020, 967, 841 cm⁻¹. ^1H and ^{13}C NMR data (Table 1). HRESIMS *m/z*: 561.1570 [*M* – H][–] (calcd for C₃₄H₂₅O₈, 561.1555).

Anti-HBV Assays. The procedure for the anti-HBV screen was performed in accordance with our previous reports.¹⁸ Anti-HBV activity and cytotoxicity of mulberrofuran G (1) and isomulberrofuran G (2) were assayed on the HepG 2.2.15 cell line *in vitro*, which was stably transfected with the HBV genome using the Lipofectamine 2000 reagent (Invitrogen, Carlsbad, CA). The activity inhibiting HBV antigen [hepatitis B surface antigen (HBsAg) and hepatitis B e antigen (HBeAg)] secretion and HBV DNA replication was analyzed by enzyme-linked immunosorbent assay (ELISA) and real-time polymerase chain reaction (PCR) methods, and cytotoxicity was measured by a




Figure 2. Selected 2D NMR correlations for isomulberrofuran G (2).

Table 2. Accurate Masses and Elemental Compositions of Mulberrofuran G (1) and Isomulberrofuran G (2) from Negative ESI–IT–TOF MSⁿ Experiments

MS ⁿ	precursor ion	product ion	calculated <i>m/z</i>	measured <i>m/z</i>	error <i>m/z</i>	error ppm	elemental composition
MS	mulberrofuran G (1)	561	561.1555	561.1536	-1.9	-3.39	C ₃₄ H ₂₅ O ₈
MS ²	561 ([M - H] ⁻)	451	451.1187	451.1153	-3.4	-7.54	C ₂₈ H ₁₉ O ₆
MS ³	561-451	436	436.0952	436.0949	-0.3	-0.69	C ₂₇ H ₁₆ O ₆
MS ³	561-451	437	437.1031	437.0991	-4.0	-9.15	C ₂₇ H ₁₇ O ₆
MS ⁴	561-451-437	408	408.1003	408.1006	0.3	0.74	C ₂₆ H ₁₆ O ₅
MS ⁵	561-451-437-408	381	381.1132	381.1119	-1.3	-3.41	C ₂₅ H ₁₇ O ₄
MS ⁶	561-451-437-408-381	352	352.1105	352.1144	3.9	11.08	C ₂₄ H ₁₆ O ₃
MS ⁷	561-451-437-408-381-352	324	324.1156	324.1124	-3.2	-9.87	C ₂₃ H ₁₆ O ₂
MS	isomulberrofuran G (2)	561	561.1555	561.1570	1.5	2.67	C ₃₄ H ₂₅ O ₈
MS ²	561 ([M - H] ⁻)	439	439.1187	439.1160	-2.7	-6.15	C ₂₇ H ₁₉ O ₆
MS ³	561-439	422	422.1160	422.1136	-2.4	-5.69	C ₂₇ H ₁₈ O ₅
MS ⁴	561-439-422	379	379.0976	379.0965	-1.1	-2.90	C ₂₅ H ₁₅ O ₄
MS ⁵	561-439-422-379	337	337.0870	337.0885	1.5	4.45	C ₂₃ H ₁₃ O ₃
MS ³	561-439	396	396.1367	396.1338	-2.9	-7.32	C ₂₆ H ₂₀ O ₄
MS ⁴	561-439-396	355	355.0976	355.0980	0.4	1.13	C ₂₃ H ₁₅ O ₄
MS ⁴	561-439-396	311	311.1078	311.1057	-2.1	-6.75	C ₂₂ H ₁₅ O ₂
MS ⁵	561-439-396-355	311	311.1078	311.1037	-4.1	-13.18	C ₂₂ H ₁₅ O ₂
MS ⁵	561-439-396-311	282	282.1050	282.1026	-2.4	-8.51	C ₂₁ H ₁₄ O
MS ⁶	561-439-396-311-282	254	254.0737	254.0689	-4.8	-18.89	C ₁₉ H ₁₀ O

modified (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) method. Samples were dissolved in dimethylsulfoxide (DMSO) for anti-HBV activity and cytotoxicity assays, with the concentration below 2.5 $\mu\text{L}/\text{mL}$ not affecting the growth of cells.

RESULTS AND DISCUSSION

Structural Identification. Isomulberrofuran G (2) was isolated as a brownish yellow powder, with the molecular formula of C₃₄H₂₆O₈ identical to mulberrofuran G (1), which was determined by negative HRESIMS ([M - H]⁻). The ¹³C [distortionless enhancement by polarization transfer (DEPT)] spectrum displayed 34 carbons, involving 1 methyl, 1 methylene, 16 methines, and 16 quaternary carbons. In ¹H NMR spectrum, five hydroxy groups were recognized at the downfield and confirmed by the heteronuclear singular-quantum correlation (HSQC) experiment. Two hydroxy groups overlapped at δ_{H} 8.54 (OH-6 and OH-8) in combination with the aromatic protons at δ_{H} 6.86 (2H, d, *J* = 2.0 Hz, H-5, H-9) and 6.37 (1H, d, *J* = 2.0 Hz, H-7), indicating a symmetric 3,5-dihydroxyphenyl moiety. Two characteristic ABX spin systems were denoted by the resonances at δ_{H} 6.43 (1H, d, *J* = 2.4 Hz, H-11''), 6.21 (1H, dd, *J* = 8.8, 2.4 Hz, H-13''), and 7.26 (1H, d, *J* = 8.8 Hz, H-14'') and at δ_{H} 6.37 (1H, d, *J* = 2.5 Hz, H-17''), 6.48 (1H, dd, *J* = 8.3, 2.5 Hz, H-19''), and 7.09 (1H, d, *J* = 8.3 Hz, H-20''), which

were determined as two 2,4-dihydroxyphenyl moieties. Three singlet peaks at δ_{H} 7.03 (1H, s, H-3), 7.11 (1H, s, H-2'), and 7.41 (1H, s, H-5') were displayed in the ¹H NMR spectrum, indicating three isolated aromatic protons. Isomulberrofuran G (2) was proposed to have the same ring E portion with mulberrofuran G (1) from ¹H-¹H (homonuclear chemical shift) correlation spectroscopy (COSY) (H-2''/H-3''/H-4''/H-5''/H-6''), heteronuclear multiple-bond correlation (HMBC) (H-7''/C-2'' and C-6'' and H-3'' and H-5''/C-1''), and rotating frame Overhauser effect spectroscopy (ROESY) (H-4''/H-3'', H-4''/6'' α , and H-5''/H-6'' β) experiments, as well as the almost identical NMR data and coupling constants (*J*) of H-3'', H-4'', H-5'', and H-6'' as those of mulberrofuran G (1). Similarly, the identical rings D, F, G, and H were constructed on the basis of extensive 1D and 2D NMR spectrometric analyses (Table 1 and Figure 2). A 2-(3,5-dihydroxyphenyl)benzofuran moiety (rings A-C) was completed by HMBC correlations of H-5 with C-2, H-3 with C-4 and C-1', H-5' with C-3 and C-1', and H-2' with C-4' and C-6', together with ROESY cross-peaks of OH-6/H-5 and H-7, OH-8/H-7 and H-9, H-9/H-3, and H-3/H-5'. The detected ROESY correlation of H-5' with H-2'' and HMBC correlations of H-2' with C-4' and C-6' were important for the determination of the linear coupled mode for rings B, C, and D. The above analyses were further confirmed by comparing its NMR data to those of australisin C,¹⁹ indicating

Table 3. Anti-HBV Activity and Cytotoxicity of Mulberrofurin G (1) and Isomulberrofurin G (2)^a

compounds	MTT		HBsAg		HBeAg		HBV DNA	
	CC ₅₀	IC ₅₀	SI	IC ₅₀	SI	IC ₅₀	SI	
mulberrofurin G (1)	8.04 ± 1.00	14.34 ± 0.31	<1	61.55 ± 3.57	<1	3.99 ± 2.28	2.0	
isomulberrofurin G (2)	6.79 ± 0.24	33.77 ± 4.59	<1	183.19 ± 26.11	<1	9.10 ± 2.37	<1	
tenofovir ^b	>1740.89	>1740.89	^c	>1740.89	^c	2.54 ± 0.14	>685.4	

^aResults are expressed as the mean (μM) \pm standard deviation (SD) from three independent experiments. ^bTenofovir was used as the positive control. ^cCC₅₀ and IC₅₀ were not reached at the highest tested concentration.




Figure 3. Cytotoxicity and anti-HBV activity of mulberrofurin G (1) and isomulberrofurin G (2) on HepG 2.2.15 cells: (A) cytotoxicity, (B) anti-HBsAg activity, (C) anti-HBeAg activity, and (D) anti-HBV DNA activity. Results are expressed as the mean \pm SD from three independent experiments. Significant difference: $p < 0.05$.

the same rings A–C. Thus, the structure of isomulberrofurin G (2) was constructed and confirmed by the HMBC, ¹H–¹H COSY, and ROESY analyses (Figure 2). It should be noted that the correlations between the hydroxy groups and adjacent aromatic protons in the ROESY experiment played important roles in structural determination, which benefited from the hydroxy signals that appeared in the ¹H NMR experiment (acetone-*d*₆).

MSⁿ Fragmentation Study. Mulberrofurin G (1) and isomulberrofurin G (2) are a pair of isomers with identical molecular formula of C₃₄H₂₆O₈ and similar spectroscopic properties. Structurally, they occupy a complicated system consisting of eight rings with five hydroxy groups, and the main difference is located at rings A–C. The similar structural features and spectroscopic properties present challenges for rapid and accurate differentiation of them. A mass spectrometer equipped with an ESI source and IT analyzer coupled to a TOF mass analyzer (ESI–IT–TOF) is effective for the fragmentation study on complicated compounds, which enables multistage tandem spectra (MS^{1–10}) with high accuracy and resolution in both MS and MSⁿ modes.^{20,21} Up to date, no extensive MS fragmentation study has been conducted on Diels–Alder-type adducts. Therefore, the first-time characterization of the fragmentation patterns of mulberrofurin G (1) and isomulberrofurin G (2) will provide valuable information

for identification and differentiation of Diels–Alder-type adducts.

The accurate mass spectrometric data of compounds 1 and 2 were obtained from 16 multistage tandem MS experiments in negative ESI mode. Generally, the most abundant ion (base peak) or the product ion with the highest mass (excluding adducts with a larger mass than that of the precursor ion) in each spectrum was subjected to the next stage analysis. The accurate masses and corresponding assigned elemental compositions of the product ions from the tandem MS analyses were summarized in Table 2. The accuracy of obtained masses relative to the assigned structures was always within ± 5 mDa or 10 ppm error, which made the calculated elemental compositions reliable.

In the single-stage mass experiments, using full-scan acquisition (m/z 100–1000) in negative-ion mode, both the deprotonated molecular ions $[M - H]^-$ and adduct ions $[M + Cl]^-$ were readily observed for mulberrofurin G (1) and isomulberrofurin G (2). When their respective $[M - H]^-$ ion was selected as the precursor ion to perform MS² analysis, different product ions at m/z 451.1153 for compound 1 and m/z 439.1160 for compound 2 were detected. This distinction resulted in obviously different fragmentations in the subsequent multistage tandem MSⁿ analyses (Table 2). Therefore, the product ions at m/z 451.1153 and 439.1160 in MS² spectra could be recognized as the diagnostic ions for mulberrofurin G

(1) and isomulberrofuran G (2), from which they can be rapidly and accurately distinguished.

Anti-HBV Investigation. To evaluate the anti-HBV properties of the isolates, mulberrofuran G (1) and isomulberrofuran G (2) were assayed on the HepG 2.2.15 cell line *in vitro*. From the data presented in Table 3 and Figure 3, compound 1 exhibited moderate activity against HBV DNA replication, with the IC_{50} value of $3.99 \mu M$ [selective index (SI) = 2.0]. Although mulberrofuran G (1) and isomulberrofuran G (2) showed potent anti-HBV DNA activities with low IC_{50} values equivalent to that of tenofovir, their obvious cytotoxicity (CC_{50} value is 8.04 and $6.79 \mu M$, respectively) resulted in low selectivity. At low concentrations ($<3.56 \mu M$), compounds 1 and 2 exhibited weak cytotoxicity; however, the damage ratio to HepG 2.2.15 cells increased rapidly with the concentration up to $14.20 \mu M$. Since then, the cytotoxicity was not increased with the concentration increasing (from 14.20 to $909.42 \mu M$). The anti-HBV activities of compounds 1 and 2 showed similar trends to their cytotoxicity. Thus, it will be very interesting to perform an extensive structure–modification study on mulberrofuran G (1) and isomulberrofuran G (2) to decrease their cytotoxicity for developing novel anti-HBV agents.

Briefly, mulberrofuran G (1), belonging to the IIA-type Diels–Alder-type adduct, was initially isolated from *Morus lhou*,²² whereas its isomer isomulberrofuran G (2) is the first example of the IIB-type Diels–Alder-type adduct with ketolization. The tandem mass spectrometric study on mulberrofuran G (1) and isomulberrofuran G (2) resulted in obviously different MS fragmentation regularities and their respective diagnostic ions at m/z 451.1153 and 439.1160. Mulberrofuran G (1) showed potent activity, inhibiting HBV DNA replication with the IC_{50} value of $3.99 \mu M$ based on the anti-HBV assay on the HepG 2.2.15 cell line *in vitro*. To our knowledge, this is the first report on the anti-HBV constituents of Mori Cortex and MSⁿ fragmentation rules of Diels–Alder-type adducts, which will provide valuable information for the further understanding of *M. alba*.

■ ASSOCIATED CONTENT

Supporting Information

¹H, ¹³C (DEPT), and 2D NMR (HSQC, HMBC, ¹H–¹H COSY, and ROESY), HRESIMS, IR, UV, and [α]_D spectra for isomulberrofuran G (2), as well as the MSⁿ spectra for mulberrofuran G (1) and isomulberrofuran G (2). This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*Telephone: (+86) 0871-5223265. Fax: (+86) 0871-5227197. E-mail: chenjj@mail.kib.ac.cn.

Funding

This work was funded by the National Science Foundation of China for Distinguished Young Scholars (81025023).

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We thank the staff of the analytical group of the State Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, Chinese Academy of Sciences.

■ ABBREVIATIONS USED

CID, collision-induced dissociation; COSY, (homonuclear chemical shift) correlation spectroscopy; DEPT, distortionless enhancement by polarization transfer; ESI, electrospray ionization; fwhm, full width at half maximum; HBeAg, hepatitis B e antigen; HBsAg, hepatitis B surface antigen; HBV, hepatitis B virus; HMBC, heteronuclear multiple-bond correlation; HSQC, heteronuclear singular-quantum correlation; ROESY, rotating frame Overhauser effect spectroscopy; SI, selective index; TOF, time of flight

■ REFERENCES

- (1) Mei, M.; Ruan, J. Q.; Wu, W. J.; Zhou, R. N.; Lei, J. P. C.; Zhao, H. Y.; Yan, R.; Wang, Y. T. *In vitro* pharmacokinetic characterization of mulberroside A, the main polyhydroxylated stilbene in mulberry (*Morus alba* L.), and its bacterial metabolite oxyresveratrol in traditional oral use. *J. Agric. Food Chem.* **2012**, *60*, 2299–2308.
- (2) State Pharmacopoeia Committee. *Chinese Pharmacopoeia*; China Medical Pharmaceutical Science and Technology Publishing House: Beijing, China, 2010; p 182.
- (3) Asano, N.; Yamashita, T.; Yasuda, K.; Ikeda, K.; Kizu, H.; Kameda, Y.; Kato, A.; Nash, R. J.; Lee, H. S.; Ryu, K. S. Polyhydroxylated alkaloids isolated from mulberry trees (*Morus alba* L.) and silkworms (*Bombyx mori* L.). *J. Agric. Food Chem.* **2001**, *49*, 4208–4213.
- (4) Nomura, T.; Hano, Y.; Fukai, T. Chemistry and biosynthesis of isoprenylated flavonoids from Japanese mulberry tree. *Proc. Jpn. Acad.* **2009**, *B85*, 391–408.
- (5) Geng, C. A.; Yao, S. Y.; Xue, D. Q.; Zuo, A. X.; Zhang, X. M.; Jiang, Z. Y.; Ma, Y. B.; Chen, J. J. New isoprenylated flavonoid from *Morus alba*. *Chin. J. Chin. Mater. Med.* **2010**, *35*, 1560–1565.
- (6) Pawlowska, A. M.; Oleszek, W.; Braca, A. Quali-quantitative analyses of flavonoids of *Morus nigra* L. and *Morus alba* L. (Moraceae) fruits. *J. Agric. Food Chem.* **2008**, *56*, 3377–3380.
- (7) Fozing, C. D. A.; Ali, Z.; Ngadjue, B. T.; Choudhary, M. I.; Kapche, G. D. W. F.; Abegaz, B.; Khan, I. A. Phosphodiesterase I-inhibiting Diels–Alder adducts from the leaves of *Morus mesozygia*. *Planta Med.* **2012**, *78*, 154–159.
- (8) Hano, Y.; Kohno, H.; Itoh, M.; Nomura, T. Structures of three new 2-arylbenzofuran derivatives from the Chinese crude drug “Sang-Bai-Pi” (*Morus* root bark). *Chem. Pharm. Bull.* **1985**, *33*, 5294–5300.
- (9) Luo, S. D.; Nemeč, J.; Ning, B. M. Anti-HIV flavonoids from *Morus alba*. *Acta Bot. Yunnanica* **1995**, *17*, 89–95.
- (10) Dai, S. J.; Chen, R. Y.; Yu, D. Q. Diels–Alder-type adducts of *Morus macroura* Miq. *J. Chin. Pharm. Univ.* **2006**, *37*, 119–122.
- (11) Hano, Y.; Takizawa, S.; Mizuno, E.; Nomura, T. Structure of kuwanon P, a new Diels–Alder type adduct from the root bark of the cultivated mulberry tree (*Morus lhou* (Ser.) Koidz. *Chem. Pharm. Bull.* **1983**, *31*, 2936–2939.
- (12) Nikaido, T.; Ohmoto, T.; Nomura, T.; Fukai, T.; Sankawa, U. Inhibition of adenosine 3',5'-cyclic monophosphate phosphodiesterase by phenolic constituents of mulberry tree. *Chem. Pharm. Bull.* **1984**, *32*, 4929–4934.
- (13) Basnet, P.; Kadota, S.; Terashma, S.; Shimizu, M.; Namba, T. Two new 2-arylbenzofuran derivatives from hypoglycemic activity-bearing fraction of *Morus insignis*. *Chem. Pharm. Bull.* **1993**, *41*, 1238–1243.
- (14) Tan, Y. X.; Liu, C.; Zhang, T.; Chen, R. Y.; Yu, D. Q. Bioactive constituents of *Morus wittiorum*. *Phytochem. Lett.* **2010**, *3*, 57–61.
- (15) Gunawan, C.; Rizzacasa, M. A. Mulberry Diels–Alder adducts: Synthesis of chalconoracin and mulberrofuran C methyl ethers. *Org. Lett.* **2010**, *12*, 1388–1391.
- (16) Hirakura, K.; Hano, Y.; Fukai, T.; Nomura, T.; Uzawa, J.; Fukushima, K. Structures of three new natural Diels–Alder type adducts, kuwanons P and X, and mulberrofuran J, from the cultivated mulberry tree (*Morus lhou* Koidz.). *Chem. Pharm. Bull.* **1985**, *33*, 1088–1096.

(17) Ikuta, J.; Fukai, T.; Nomura, T.; Ueda, S. Constituents of *Morus alba* L. cell cultures. (1). Structures of four new natural Diels–Alder type adducts, kuwanons J, Q, R, and V. *Chem. Pharm. Bull.* **1986**, *34*, 2471–2478.

(18) Geng, C. A.; Wang, L. J.; Zhang, X. M.; Ma, Y. B.; Huang, X. Y.; Luo, J.; Guo, R. H.; Zhou, J.; Shen, Y.; Zuo, A. X.; Jiang, Z. Y.; Chen, J. J. Anti-hepatitis B virus active lactones from the traditional Chinese herb: *Swertia mileensis*. *Chem.—Eur. J.* **2011**, *17*, 3893–3903.

(19) Zhang, Q. J.; Tang, Y. B.; Chen, R. Y.; Yu, D. Q. Three new cytotoxic Diels–Alder type adducts from *Morus australis*. *Chem. Biodiversity* **2007**, *4*, 1533–1540.

(20) Li, Q. R.; Yan, G. Q.; Ge, T. T. A fragmentation study of podophyllotoxin and its 4'-demethyl-4 β -substituted derivatives by electrospray ionization ion-trap time-of-flight tandem mass spectrometry. *Rapid Commun. Mass Spectrom.* **2007**, *21*, 2843–2852.

(21) Shibano, M.; Lin, A. S.; Itokawa, H.; Lee, K. H. Separation and characterization of active flavonolignans of *Silybum marianum* by liquid chromatography connected with hybrid ion-trap and time-of-flight mass spectrometry (LC–MS/IT–TOF). *J. Nat. Prod.* **2007**, *70*, 1424–1428.

(22) Fukai, T.; Hano, Y.; Hirakura, K.; Nomura, T.; Uzawa, J.; Fukushima, K. Structures of two natural hypotensive Diels–Alder type adducts, mulberrofurans F and G, from the cultivated mulberry tree (*Morus lhou* Koidz.). *Chem. Pharm. Bull.* **1985**, *33*, 3195–3204.