This article was downloaded by: *[CAS Consortium]* On: *13 May 2009* Access details: *Access Details: [subscription number 909168890]* Publisher *Taylor & Francis* Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Asian Natural Products Research

Publication details, including instructions for authors and subscription information: http://www.informaworld.com/smpp/title~content=t713454007

Cytotoxic prenylated xanthones from Calophyllum inophyllum

Qi Xiao ^a; Yan-Bo Zeng ^a; Wen-Li Mei ^a; You-Xing Zhao ^b; Yuan-Yuan Deng ^a; Hao-Fu Dai ^a ^a Institute of Tropical Bioscience and Biotechnology, Chinese Academy of Tropical Agricultural Sciences, Haikou, China ^b State Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, China

Online Publication Date: 01 October 2008

To cite this Article Xiao, Qi, Zeng, Yan-Bo, Mei, Wen-Li, Zhao, You-Xing, Deng, Yuan-Yuan and Dai, Hao-Fu(2008)'Cytotoxic prenylated xanthones from Calophyllum inophyllum', Journal of Asian Natural Products Research, 10:10,993 — 997 To link to this Article: DOI: 10.1080/10519990802240387

URL: http://dx.doi.org/10.1080/10519990802240387

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: http://www.informaworld.com/terms-and-conditions-of-access.pdf

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Journal of Asian Natural Products Research Vol. 10, No. 10, October 2008, 993–997



Cytotoxic prenylated xanthones from Calophyllum inophyllum

Qi Xiao^a, Yan-Bo Zeng^a, Wen-Li Mei^a*, You-Xing Zhao^b, Yuan-Yuan Deng^a and Hao-Fu Dai^a*

^aInstitute of Tropical Bioscience and Biotechnology, Chinese Academy of Tropical Agricultural Sciences, Haikou 571101, China; ^bState Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650204, China

(Received 29 February 2008; final version received 5 May 2008)

A new prenylated xanthone (1), named caloxanthone N, together with two known constituents, gerontoxanthone C (2) and 2-hydroxyxanthone (3), was isolated from the ethanolic extract of the twigs of *Calophyllum inophyllum*. Their structures were completely elucidated using a combination of 1D, 2D NMR techniques (COSY, HMQC, HMBC, and ROESY) and HR-ESI-MS analyses. Compounds 1 and 2 exhibited cytotoxicity against chronic myelogenous leukemia cell line (K562) with IC₅₀ values of 7.2 and 6.3 μ g ml⁻¹, respectively.

Keywords: Calophyllum inophyllum; cytotoxic activity; prenylated xanthone; caloxanthone N

1. Introduction

Calophyllum inophyllum Linn. (Clusiaceae) is an evergreen shrub widely distributed in tropical areas. In China, C. inophyllum is distributed in Hainan and Taiwan provinces, and is used in traditional Chinese folk medicine for the treatment of eye diseases, wounds, rheumatisms, and inflammations [1,2]. Since inophyllums B and P isolated by Ptail [3] showed strong activity against human immunodeficiency virus type 1, much attention has been paid to studies on the chemical components of C. inophyllum. These studies have revealed that, besides pyranocoumarins [4-9], C. inophyllum is also a rich source of xanthones [4,5,8], triterpenes [9,10], and flavonoids [11]. In our on-going search for cytotoxic agents from tropical medicinal plants in Hainan, the ethanolic extract of the twigs of C. inophyllum showed inhibitory activity toward human chronic myelogenous leukemia cell line (K562). Bioassay-guided fractionation led to the isolation of a new and two known xanthones, and their structures were unambiguously elucidated as caloxanthone N (1), gerontoxanthone C (2), and 2-hydroxyanthone (3) (Figure 1) by extensive spectroscopic analysis. Compounds 1 and 2 showed significant cytotoxicity against chronic myelogenous leukemia cell line (K562) *in vitro* by MTT method. In this paper, we report the isolation and identification of the cytotoxic principles from *C. inophyllum*.

2. Results and discussion

Compound 1 was obtained as yellow powder and reacted positively to the Gibbs and FeCl₃ reagent indicating the presence of a phenolic group. The $[M + Na]^+$ ion peak at m/z449.1576 in HR-ESI-MS spectrum corresponded to the molecular formula C₂₄H₂₆O₇. This formula can also be validated through ¹H-NMR, ¹³C-NMR, and DEPT spectra. The IR spectrum displayed free hydroxyl

^{*}Corresponding authors. Email: meiwenli@yahoo.com.cn; hfdai@yahoo.cn

Q. Xiao et al.



Figure 1. Structures of compounds 1 and 2.

(3473 cm⁻¹), chelated hydroxyl (3244 cm⁻¹), conjugated carbonyl (1648 cm⁻¹), and aromatic ring (1605, 1565, 1505, 1479 cm⁻¹) absorptions. These data, together with those obtained from the UV spectrum [λ (MeOH) 238, 249, 255, 280 sh, and 331 nm] were

consistent with the presence of a xanthone skeleton [4,5]. In the ¹H-NMR spectrum (Table 1), a chelated hydroxyl group (δ 13.66), two aromatic proton singlets (δ 7.02 and 6.12), and one methoxyl (δ 3.98) were observed. The ¹H-NMR spectrum of **1** also

Table 1. 1 H (400 MHz) and 13 C (100 MHz) NMR spectral data of 1 (acetone- d_6) and 2 (CD₃OD).

No.	1		2	
	$\delta_{\rm C}$	$\delta_{ m H}$ (Hz)	$\delta_{ m C}$	δ_{H} (Hz)
1	165.5 (s)	13.66 (1H, s)	161.9 (s)	
2	94.2 (d)	6.12 (1H, s)	113.2 (s)	
3	166.7 (s)		165.4 (s)	
4	113.2 (s)		107.7 (s)	
5	135.2 (s)		134.1 (s)	
6	156.3 (s)		153.0 (s)	
7	112.8 (d)	7.02 (1H, s)	113.7 (d)	6.85 (1H, d, 8.7)
8	138.2 (s)		117.6 (d)	7.56 (1H, d, 8.7)
9	183.1 (s)		182.0 (s)	
4a	152.9 (s)		152.5 (s)	
8a	112.0 (s)		115.2 (s)	
9a	104.4 (s)		103.8 (s)	
10a	152.4 (s)		147.9 (s)	
11	44.5 (s)		45.3 (s)	
12	21.8 (q)	1.33 (3H, s)	21.5 (q)	1.76 (3H, s)
13	25.9 (q)	1.62 (3H, s)	25.9 (q)	1.32 (3H, s)
14	91.7 (d)	4.56 (1H, q, 6.5)	92.0 (d)	4.50 (1H, q, 6.5)
15	14.5 (q)	1.40 (3H, d, 6.4)	14.7 (q)	1.40 (3H, d, 6.6)
16	126.8 (d)	7.89 (1H, d, 15.9)	22.5 (t)	3.23 (2H, d, 7.3)
17	142.3 (d)	6.20 (1H, d, 15.9)	123.1 (d)	5.23 (1H, t, 7.3)
18	70.8 (s)		132.4 (s)	
19	30.6 (q)	1.40 (3H, s)	26.1 (q)	1.65 (3H, s)
20	30.6 (q)	1.40 (3H, s)	17.9 (q)	1.61 (3H, s)
5-OMe	62.0 (q)	3.98 (3H, s)		

showed two methyl singlets (δ 1.40, 6H) and two *trans*-olefinic protons (δ 7.89 and 6.20, each 1H, d, J = 15.9 Hz), which suggested the presence of a γ , γ -dimethylallyl chain. Two methyl singlets (δ 1.33 and 1.62, each 3H), a methyl doublet (δ 1.40, 3H, d, J = 6.4 Hz), and one-proton quartet (δ 4.56, 1H, q, J = 6.5 Hz) suggested the presence of an α, α, β -trimethyldihydrofuran ring. A combination of the ¹H-¹H COSY and HSQC experiments permitted the assignment of all of the protonated carbons (Table 1). It remained to establish the positions of the substituents on the xanthone skeleton. In the HMBC spectrum (Figure 2), the proton of chelated hydroxyl group (δ 13.66) was correlated to three carbons C-9a (δ 104.4), C-2 (894.2), and C-1 (8165.5), which suggested that the chelated hydroxyl group was located at C-1. Aromatic carbons with an oxygen function were observed at C-1 (δ 165.5), C-3 (δ166.7), and C-4a (δ152.9) in the ¹³C NMR spectrum, which indicated that this aromatic ring was a phloroglucinol ring [12]. Therefore, the α, α, β -trimethyldihydrofuran ring was clearly fused at C-4 through an oxygen at C-3 position. This result was further confirmed by the long-range correlations of H-12 (δ 1.33) and H-13 (δ 1.62) with C-4 (δ 113.2) and H-14 (δ 4.56) with C-3 (δ 166.7) in the HMBC spectrum. The methoxyl group was



Figure 2. Key HMBC and ROESY correlations of compound **1**.

deduced to be located at C-5 position by ROESY experiment revealing the cross-peaks from the methoxyl (δ 3.98) to H-12 (δ 1.33) and H-13 (δ 1.62). The positions of the γ,γ -dimethylallyl moiety and the remaining phenolic hydroxyl group were established as follows. In the HMBC spectrum (Figure 2), one of the olefinic proton H-16 (δ 7.89) was correlated to three aromatic carbons C-8a $(\delta 112.0)$, C-7 $(\delta 112.8)$, and C-8 $(\delta 138.2)$. The resonance of C-8 (δ 138.2) also gave crosspeak with the other olefinic proton H-17 (δ 6.20). These results demonstrated clearly that the γ,γ -dimethylallyl moiety was located at C-8. The cross-peaks from the aromatic proton (\$ 7.02) to C-8a (\$ 112.0), C-16 (\$ 126.8), and C-5 (8 135.2) in the HMBC spectrum indicated that the aromatic proton (δ 7.02) was assigned to be at C-7 position. The downfield shifts of C-6 (δ 156.3) and C-18 (δ 70.8) revealed that these two carbons should be substituted by hydroxyl group. The ROESY experiment showed that H-14 (δ 4.56) correlated with H-13 (δ 1.62), while not with H-12 (δ 1.33). This result indicated that when H-14 and CH₃-13 were assigned β -orientation, CH₃-12 and CH₃-15 were in α -orientation. On the basis of the above results, the structure of compound 1 was thus elucidated as 1, named caloxanthone N.

The structures of compounds 2 and 3 were identified to be gerontoxanthone C (2) [13] and 2-hydroxy-xanthone (3) [14] by comparing their corresponding spectroscopic properties (NMR and MS) with the values reported in the literatures.

Compounds 1-3 were evaluated for their cytotoxic activity against chronic myelogenous leukemia cell line (K562) using the MTT method. Compounds 1 and 2 showed cytotoxic activity with IC₅₀ values of 7.2 and 6.3 µg ml⁻¹, respectively, while compound 3 was inactive (>20 µg ml⁻¹).

3. Experimental

3.1 General experimental procedures

Melting point determinations were obtained on a Kofler hot stage apparatus, and are Q. Xiao et al.

uncorrected. Optical rotation was measured at room temperature using a J-20C polarimeter. The UV spectra were measured on a Shimadzu UV-210A spectrometer. The IR spectra were measured on a Bio-Tad FTS 135 instrument, as KBr pellets. The ¹H, ¹³C, and 2D NMR spectra were run on Bruker AM-400 and DRX-500 spectrometers, using TMS as an internal standard. The FAB-MS spectra were measured with a VG Auto Spectrometer. Column chromatography was performed with silica gel (Marine Chemical Industry Factory, Qingdao, China) and Sephadex LH-20 (Merck, Darmstadt, Germany). TLC was performed with silica gel 60 F254 (Merck) and developed by spraying with 10% H₂SO₄ followed by heating.

3.2 Plant material

The twigs of *Calophyllum inophyllum* L. used in this research were collected in Wenchang county, Hainan province, China, in May 2006, and authenticated by associate researcher Zheng-Fu Dai (Institute of Tropical Bioscience and Biotechnology, Chinese Academy of Tropical Agricultural Sciences, Haikou, China). The voucher specimen (No. 20060508) is deposited in the Institute of Tropical Bioscience and Biotechnology, Chinese Academy of Tropical Agricultural Sciences.

3.3 Extraction and isolation

The dried and crushed twigs of *C. inophyllum* (19.9 kg) were extracted with 95% EtOH thrice at room temperature. After the removal of EtOH by evaporation, the EtOH extract was suspended in water (6.01) and successively partitioned with petroleum ether to give Petro-soluble extract (290.0 g) and an aqueous residue. The aqueous residue was subjected to chromatography on D-101 resin with H₂O, a gradient of 50, 95, and 100% MeOH as the eluents, to afford four fractions. The 50, 95, and 100% MeOH eluents were collected and concentrated *in vacuo* to yield 65.0, 245.0, and 70.0 g residue, respectively.

The Petro-soluble extract (290.0 g) was applied to a silica gel (200-300 mesh) column packed in CHCl₃. The column was then eluted in gradient elution with CHCl3acetone to afford 16 fractions. The active fraction (fraction 16, 44.0g) was then subjected to repeated column chromatography over silica gel using Pet-EtOAc as eluent and further separated by column chromatography over Sephadex LH-20 using 95% EtOH as eluent to afford 1 (24.7 mg). The 95% MeOH fraction (245.0 g) was subjected to vacuum liquid chromatography over silica gel, eluting with gradient elution CHCl₃-MeOH to afford 10 fractions. The active fraction (fraction 3, 5.7 g) was then subjected to repeated column chromatography over silica gel using Petacetone as eluent and further separated by column chromatography over Sephadex LH-20 using 95% EtOH as eluent to afford 2 (15.6 mg) and 3 (10.5 mg).

3.3.1 Caloxanthone N(1)

Yellow needles from EtOH (95%), 24.7 mg, mp 179–181°C, $[\alpha]_D^{17}$ + 59.32 (*c* 1.2, acetone); UV (MeOH) λ_{max} (log ε): 238 (2.06), 249 (2.05), 255 (2.60), 280 (sh) and 331 (2.32) nm; IR (KBr) ν_{max} : 3473, 3244, 2976, 1648, 1605, 1565, 1505, 1479, 1411, 1380, 1272, 1129, 1062, 1002, 960, 868, 823 cm⁻¹; ¹H-NMR and ¹³C-NMR spectral data: see Table 1; HR-FAB-MS: *m/z* 449.1576 [M + Na]⁺ (calcd for C₂₄H₂₆O₇Na, 449.1571).

3.3.2 Gerontoxanthone C(2)

Pale yellow needles from EtOH (95%), 15.6 mg, ¹H-NMR and ¹³C-NMR spectral data were consistent with the literature [13].

3.3.3 2-Hydroxyxanthone (3)

Yellow needles from EtOH (95%), 10.5 mg. ¹H-NMR and ¹³C-NMR spectral data were consistent with the literature [14].

3.4 Bioassay

The 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay was performed according to the previously reported method [15]. The inhibition rates (IR%) were calculated using OD mean values from IR% = (OD_{control} – OD_{sample})/ OD_{control}. The IC₅₀ value, which was defined as the concentration of sample needed to reduce a 50% of absorbance relative to the vehicle-treated control, was determined using the Bliss method. The same experiment was repeated independently thrice to obtain a mean IC₅₀ and its SD.

Acknowledgements

This research was financially supported by National Basic Research Program of China (2007CB116306) and the Science and Technology Foundation of Chinese Academy of Agricultural Sciences (RKY0421).

References

- H.Y. Chen, *Flora of Hainanica*, Vol. 2, (Science Press, Beijing, 1965), p. 56.
- [2] H.F. Dai and W.L. Mei, Modern Research on Medicinal Plants in Hainan, (China Science and Technology Press, Beijing, 2007), p. 31.
- [3] A.D. Patil, A.J. Freyer, D.S. Eggleston, R.C. Haltiwanger, M.F. Bean, P.B. Taylor,

M.J. Caranfa, A.L. Breen, H.R. Bartus, R.K. Johnson, R.P. Hertzberg, and J.W. Westley, *J. Med. Chem.* **36**, 4132 (1993).

- [4] M. Iinuma, H. Tosa, T. Tanaka, and S. Yonemori, *Phytochemistry* 35, 527 (1994).
- [5] M. Iinuma, H. Tosa, T. Tanaka, and S. Yonemori, *Phytochemistry* 38, 725 (1995).
- [6] C. Spino, M. Dodier, and S. Sotheeswaran, *Bioorg. Med. Chem. Lett.* 8, 3475 (1998).
- [7] M. Itoigawa, C. Ito, H.T-W. Tan, M. Kuchide, H. Tokuda, H. Nishino, and H. Furukawa, *Cancer Lett.* 169, 15 (2001).
- [8] Y. Wu, P.C. Zhang, R.Y. Chen, D.Q. Yu, and X.T. Liang, *Acta Chimica. Sinica* 61, 1047 (2003).
- [9] M.C. Yimdjo, A.G. Azebaze, A.E. Nkengfack, A.M. Meyre, B. Bodo, and Z.T. Fomum, *Phytochemistry* 65, 2789 (2004).
- [10] T.R. Govindachari, N. Viswanathan, B.R. Pai, U.R. Rao, and M. Srinivasan, *Tetrahedron* 23, 1901 (1967).
- [11] Y.Z. Li, L.Z. li, H.M. Hua, Z.G. Li, and M.S. Liu, J. Chin. Materia Medica 32, 692 (2007).
- [12] M. Iinuma, T. Ito, R. Miyake, H. Tosa, T. Tanaka, and V. Chelladurai, *Phytochemistry* 47, 1169 (1998).
- [13] C.H. Chang, C.C. Lin, H. Masao, and N. Tsuneo, *Phytochemistry* 28, 595 (1989).
- [14] C. Gnerre, U. Thull, P. Gaillard, P.A. Carrupt, B. Testa, E. Fernandes, F. Silva, M. Pinto, M.M.M. Pinto, J.L. Wolfender, K. Hostettmann, and G. Cruciani, *Helv. Chim. Acta* 84, 552 (2001).
- [15] T. Mosmann, J. Immunol. Methods 65, 55 (1983).