## Neaumycin: A New Macrolide from *Streptomyces* sp. NEAU-x211

Sheng-Xiong Huang,<sup>†</sup> Xiang-Jing Wang,<sup>§</sup> Yijun Yan,<sup>§,†</sup> Ji-Dong Wang,<sup>§</sup> Ji Zhang,<sup>§</sup> Chong-Xi Liu,<sup>§</sup> Wen-Sheng Xiang,<sup>\*,§</sup> and Ben Shen<sup>\*,†,!,‡</sup>

School of Life Science, Northeast Agricultural University, Harbin, Heilongjiang 150030, China, and Department of Chemistry, Department of Molecular Therapeutics, and Natural Products Library Initiative at TSRI, The Scripps Research Institute, Jupiter, Florida, 33458, United States

xiangwensheng@yahoo.com.cn; shenb@scripps.edu

## Received January 12, 2012



Neaumycin, a new 30-membered macrolide featuring an internal diester bridge, a molecular architecture that is unprecedented among known macrolide natural products, was isolated from a soil actinomycete strain *Streptomyces* sp. NEAU-x211. The structure of neaumycin was elucidated on the basis of comprehensive mass and NMR spectroscopic interpretation, including the relative stereochemistry of four independent coupling systems.

Natural products remain the best sources of drugs and drug leads and serve as outstanding small molecule probes for dissecting fundamental biological processes.<sup>1</sup> Despite their high value, however, they are poorly represented in all current small molecule libraries, and large libraries of pure natural products are not available.<sup>1,2</sup> We recently initiated a Natural Products Library Initiative to isolate microbial natural products and assemble them into a pure natural

(1) (a) Jesse, W.-H.; Vederas, J. C. Science **2009**, *325*, 161–165. (b) Newman, D. J.; Cragg, G. M. In *RSC Biomolecular Sciences No. 18: Natural Product Chemistry for Drug Discovery*; Buss, A. D., Butler, M. S., Eds.; RSC: Cambridge, U.K., 2010; pp 3–27. (c) Schmitt, E. K.; Moore, C. M.; Krastel, P.; Petersen, F. *Curr. Opinion Chem. Biol.* **2011**, *15*, 497–504.

(2) Dandapani, S.; Marcaurelle, L. A. Nat. Chem. Biol. 2010, 6, 861–863.
(3) (a) Huang, S.-X.; Zhao, L.-X.; Tang, S.-K.; Jiang, C.-L.; Duan, Y.; Shen, B. Org. Lett. 2009, 11, 1353–1356. (b) Huang, S.-X.; Powell, E.; Rajski, S. R.; Zhao, L.-X.; Jiang, C.-L.; Duan, Y.; Xu, W.; Shen, B. Org. Lett. 2010, 12, 1353–1356. (c) Yu, Z.; Zhao, L.-X.; Jiang, C.-L.; Duan, Y.; Wong, L.; Carver, K. C.; Schuler, L. A.; Shen, B. J. Antibiot. 2011, 64, 159–162. (d) Huang, S.-X.; Yu, Z.; Robert, F.; Zhao, L.-X.; Jiang, Y.; Duan, Y.; Pelletier, J.; Shen, B. J. Antibiot. 2011, 64, 164–166. (e) Yu, Z.; Vodanovic-Jankovic, S.; Ledeboer, N.; Huang, S.-X.; Rajski, S. R.; Kron, M.; Shen, B. Org. Lett. 2011, 13, 2034–2037. (f) Zhao, L.-X.; Shen, B. jet al. J. Nat. Prod. 2011, 74, 1990–1995.

10.1021/ol300074d © 2012 American Chemical Society Published on Web 02/14/2012 product library.<sup>3</sup> We strategically selected bacteria, particularly actinomycetes for their proven track record as prolific natural product producers, as the preferred sources of natural products so that quantities of them can be produced by large scale microbial fermentation.<sup>4</sup>

Here, we report the isolation and structural elucidation of a new macrolide, named neaumycin, from *Streptomyces* sp. NEAU-x211. S. sp. NEAU-x211 is a soil actinomycete isolated from HunLunbeier, Inner Mongolia, China, whose 16S rRNA shows 98.89% sequence identity with *Streptomyces caviscabies* ATCC 51928 [Supporting Information (SI)]. Neaumycin is a 30-membered macrolide featuring an internal diester bridge, a molecular architecture that is unprecedented among known macrolide natural products.<sup>5</sup>

Neaumycin was isolated as a white solid from *S*. sp. NEAU-x211 culture (SI). The structure of neaumycin was elucidated on the basis of comprehensive mass and NMR spectroscopic data interpretation (Figure 1; SI). ESI-MS in

LETTERS 2012 Vol. 14, No. 5 1254–1257

ORGANIC

<sup>&</sup>lt;sup>§</sup>Northeast Agricultural University.

<sup>&</sup>lt;sup>†</sup> Department of Chemistry, The Scripps Research Institute.

<sup>&</sup>lt;sup>II</sup> Department of Molecular Therapeutics, The Scripps Research Institute. \*Natural Products Library Initiative at TSRI, The Scripps Research Institute

<sup>(4) (</sup>a) Van Lanen, S. G.; Shen, B. *Curr. Opinion Microbiol.* **2006**, *9*, 252–260. (b) Bérdy, J. *J. Antibiot.* **2005**, *58*, 1–26. (c) Demain, A. L.; Sanchez, S. *J. Antibiot.* **2009**, *62*, 5–16. (d) Genilloud, O.; Gonzalez, I.; Salazar, O.; Martyin, J.; Tormo, J. R.; Vicente, F. *J. Ind. Microbiol. Biotechnol.* **2011**, *38*, 375–389.

<sup>(5)</sup> Omura, S. *Macrolide Antibiotics*, 2nd ed.; Academic Press: San Diego, CA, 2002.



**Figure 1.** Structure of neaumycin. The relative stereochemistry shown in the four coupling systems of C-4/C-5, C-10 to C-14, C-22 to C-33, and C36/C-37 are independent.

negative mode yielded an  $[M - H]^-$  ion at m/z 985.7 and high resolution ESI-MS in positive mode afforded an  $[M + Na]^+$  ion at m/z 1009.58287, establishing the molecular formula of neaumycin as  $C_{55}H_{86}O_{15}$  and indicating 13 degrees of unsaturation. <sup>13</sup>C NMR (Table 1) revealed resonances for three carbonyl groups ( $\delta_C$  168.7, 166.62, and 166.61) and six double bonds ( $\delta_C$  152.5, 144.2, 138.8, 138.3, 134.9, 132.5, 131.6, 130.3, 128.7, 128.0, 126.2, and 121.5), accounting for 9 of the 13 degrees of unsaturation and suggesting neaumycin features four rings.

Three structural fragments, A, B, C, can be readily deduced on the basis of 2D NMR (COSY, HMBC, ROSEY) correlations (Figure 2, Table 1, and Table S1). Fragment A consists of C-1 ( $\delta_{\rm C}$  166.6) to C-13 ( $\delta_{\rm C}$  74.2). The connectivity from C-2 to C-10, including C-42, C-43, and C-53, and from C-11 and C-13 were established based on COSY correlations and supported by HMBC correlations of H<sub>3</sub>-42 ( $\delta_{\rm H}$  1.11) with C-3 ( $\delta_{\rm C}$  152.5), C-4 ( $\delta_{\rm C}$  44.4), and C-5 ( $\delta_{\rm C}$  75.4); H-5 ( $\delta_{\rm H}$  3.56) and H<sub>3</sub>-53 ( $\delta_{\rm H}$  3.24) with C-7 ( $\delta_{\rm C}$  85.3); and H<sub>3</sub>-43 ( $\delta_{\rm H}$  1.02) with C-9 ( $\delta_{\rm C}$  138.8) and C-10 ( $\delta_{\rm C}$  39.9). The conjugated C-2 ( $\delta_{\rm C}$  121.5)/C-3 double bond was assigned to be trans based on the large coupling constant ( $J_{2,3} = 15.5$  Hz), which was further supported by the ROESY correlation between H-2 ( $\delta_{\rm H}$  5.81) and H-4 ( $\delta_{\rm H}$  2.32). The *trans* configuration for the C-8 ( $\delta_{\rm C}$  128.0)/ C-9 double bond was deduced based on the ROESY correlations between H-7 ( $\delta_{\rm H}$  3.75) and H-9 ( $\delta_{\rm H}$  5.78) and between H-8 ( $\delta_{\rm H}$  5.25) and H-10 ( $\delta_{\rm H}$  2.18). Finally the assignment of C-1 to conjugate with the C-2/C-3 double bond was further supported by its HMBC correlations with H-2 and H-3 ( $\delta_{\rm H}$  6.78). Although COSY correlation between H-10 and H-11 ( $\delta_{\rm H}$  3.34) was not observed, the connectivity between C-10 and C-11 ( $\delta_{\rm C}$ 67.6) was established based on the HMBC correlations of H<sub>3</sub>-43 with C-11, H-11 with C-9, and H-12 ( $\delta_{\rm H}$  1.71) with C-10.

Fragment B extends from C-14 ( $\delta_{\rm C}$  81.0) to C-28 ( $\delta_{\rm C}$  34.1). The connectivity from C-14 to C-28, including C-44 and C-45, was similarly established based on COSY correlations and further supported by HMBC correlations. The geometry of three conjugated double bonds from C-16 to C-21 was established by the H–H coupling constants and ROESY correlations. Thus, both the C-16 ( $\delta_{\rm C}$  126.2)/C-17 ( $\delta_{\rm C}$  134.9) and C-20 ( $\delta_{\rm C}$  131.6)/C-21

 $(\delta_{\rm C} 138.3)$  double bonds were assigned to be *trans* based on ROESY correlations between H-15 ( $\delta_{\rm H} 2.12$ ) and H-17 ( $\delta_{\rm H} 5.74$ )/H-16 ( $\delta_{\rm H} 5.32$ ) and H-18 ( $\delta_{\rm H} 6.02$ ) and between H-19 ( $\delta_{\rm H} 5.89$ ) and H-21 ( $\delta_{\rm H} 5.30$ )/H-20 ( $\delta_{\rm H} 6.06$ ) and H-22 ( $\delta_{\rm H} 2.15$ ), respectively. Although, the overlapping or close <sup>1</sup>H NMR resonances between H-17 and H-19 and between H-18 and H-20 precluded from determining their ROESY correlations definitively, the large coupling constant ( $J_{18,19} = 14.3$  Hz) permitted the *trans* configuration assignment for the C-18/C-19 double bond. Finally, the chemical shifts and coupling constants for the three *trans* double bonds also compare favorably with those of similar structural moieties in the literature.<sup>6</sup>

Fragment C covers C-29 ( $\delta_{\rm C}$  98.8) to C-41 ( $\delta_{\rm C}$  14.5), including C-46 ( $\delta_{\rm C}$  10.1), C-55 ( $\delta_{\rm C}$  58.3), and C-47 ( $\delta_{\rm C}$ 17.1). COSY and HMBC correlations easily defined the sequence from the terminal  $CH_3$  (C-41), via an epoxide ring C-37 ( $\delta_{\rm H}$  2.56,  $\delta_{\rm C}$  60.1) and C-36 ( $\delta_{\rm H}$  3.17,  $\delta_{\rm C}$  63.6), to the OCH<sub>3</sub> (C-55,  $\delta_{\rm H}$  3.57,  $\delta_{\rm C}$  58.3) substituted C-35 ( $\delta_{\rm C}$ 80.2); the *trans* configuration of the epoxide was deduced based on ROESY correlations between H-37 ( $\delta_{\rm H}$  2.56) and H-35 ( $\delta_{\rm H}$  3.32) and between H-36 ( $\delta_{\rm H}$  3.17) and H-38 ( $\delta_{\rm H}$ 1.50). No COSY correlation between H-35 and H-34 ( $\delta_{\rm H}$ 1.58) was observed, but HMBC correlations of H<sub>3</sub>-46 ( $\delta_{\rm H}$ 0.93) with C-35, C-34 ( $\delta_{\rm C}$  44.1), and C-33 ( $\delta_{\rm C}$  72.7) allowed the connectivity between C-35 and C-34/C-33. The structural moiety from C-34 to C-30 ( $\delta_{\rm C}$  36.4) was readily assigned based on COSY correlations. The quaternary carbon C-29 was assigned to be a ketal on the basis of its characteristic <sup>13</sup>C chemical shift ( $\delta_{\rm C}$  98.8). HMBC correlation of H-30 ( $\delta_{\rm H}$  1.40) with C-29 finally completed the assembly of fragment C.

The remaining <sup>1</sup>H and <sup>13</sup>C NMR resonances were suggestive of a mesaconate residue. The latter was confirmed by HMBC correlations of H<sub>3</sub>-52 ( $\delta_{\rm H}$  2.36) with C-48 ( $\delta_{\rm C}$  168.7), C-49 ( $\delta_{\rm C}$  144.2), and C-50 ( $\delta_{\rm C}$  128.7) and by comparison of the NMR data with mesaconic acid or other known derivatives.<sup>7</sup>

The assembly of fragments A, B, C, as well as the mesaconate moiety, into the planar neaumycin structure was achieved by careful analysis of key COSY, HMBC, and ROESY correlations as depicted in Figure 2. First, although COSY correlation between H-13 and H-14 was not observed, the HMBC correlation of H-12 with C-14 and the ROESY correlation between H-13 and H-14 gave rise to the connectivity of fragments A and B through C-13 and C-14. Second, the ROESY correlation between H-28 and H-33 not only defined the relative stereochemistry of C-29 and C-33 but also supported the ring closure to a trans-disubstituted tetrahydropyran and connection between fragments B and C through C-28 and C-29. The latter was further supported by the HMBC correlation of

<sup>(6) (</sup>a) Findlay, J. A.; Radics, L. *Can. J. Chem.* **1980**, *58*, 579–590. (b) Fielder, S.; Rawan, D. D.; Sherburn, M. S. *Tetrahedron* **1998**, *54*, 12907–12922.

<sup>(7) (</sup>a) Braun, S. Org. Magn. Reson. **1978**, *11*, 197–203. (b) Wolff, M.; Seemann, M.; Grosdemange-Billiard, C.; Tritsch, D.; Campos, N.; Rodríguez-Concepción, M.; Boronat, A.; Rohmer, M. Tetrahedron Lett. **2002**, *423*, 2555–2559. (c) Tripp, J. C.; Schiesser, C. H.; Curran, D. P. J. Am. Chem. Soc. **2005**, *127*, 5518–5527.

**Table 1.** <sup>1</sup>H and <sup>13</sup>C NMR Data of Neaumycin (CDCl<sub>3</sub>,  $\delta$  in ppm, *J* in Hz)<sup>*a*</sup>

no.	<sup>1</sup> H NMR	<sup>13</sup> C	no.	<sup>1</sup> H NMR	<sup>13</sup> C
1		166.6	29		98.8
2	5.81 (d 15.5)	121.5	30	1.40 (o)	36.4
3	6.78 (dd 15.5, 10.0)	152.5	30	1.68 (0)	
4	2.32 (o) <sup>b</sup>	44.4	31	1.53 (o)	20.3
5	3.56 (o)	75.4	31	1.75 (o)	
6	1.50 (o)	40.6	32	0.998 (o)	29.2
6	1.59 (o)		32	1.72 (o)	
7	3.75 (t like 8.6)	85.3	33	3.49 (t like 11.5)	72.7
8	5.25 (o)	128.0	34	1.58 (o)	44.1
9	5.78 (o)	138.8	35	3.32 (o)	80.2
10	2.18 (o)	39.9	36	3.17 (brd 7.0)	63.6
11	3.34 (o)	67.6	37	2.56 (o)	60.1
12	1.71 (o)	37.7	38	1.50 (o)	35.1
12	1.91 (t 13.0)		39	1.30 (o)	35.6
13	4.95 (brd 10.5)	74.2	39	1.43 (o)	
14	3.39 (o)	81.0	40	1.38 (o)	20.4
15	2.12 (o)	34.6	41	0.95 (o)	14.5
15	2.80 (m)		42	1.11 (o)	17.2
16	5.32 (o)	126.2	43	1.02 (d 6.5)	10.4
17	5.74 (o)	134.9	44	1.13 (o)	17.4
18	6.02 (dd 11.3, 14.3)	130.3	45	0.81 (d 6.5)	13.6
19	5.89 (dd 11.3, 14.3)	132.5	46	0.93 (o)	10.1
20	6.06 (dd 11.3, 15.0)	131.6	47	1.07 (d 6.5)	17.1
21	5.30 (o)	138.3	48		168.7
22	2.15 (0)	45.0	49		144.2
23	3.31 (0)	77.6	50	6.70 (brs)	128.7
24	1.24 (o)	37.4	51	× /	166.6
24	2.165 (0)		52	2.36 (brs)	14.1
25	3.90 (t 10.5)	77.2	53	3.24 (s)	55.8
26	1.48 (o)	21.8	54	3.43 (s)	57.9
27	4.88 (dt 10.5, 3.6)	71.4	55	3.57 (s)	58.3
28	1.31 (0)	34.1			
28	2.54 (o)				

<sup>*a* <sup>1</sup></sup>H NMR at 500 MHz, <sup>13</sup>C NMR at 125 MHz, and assignments were based on HSQC, COSY, HMBC, and ROESY experiments (Table S1). <sup>*b*</sup> 'o' denotes overlapping signals.

H-28 with C-29. Third, the upfield <sup>13</sup>C chemical shifts of the two carbonyl groups (C-51 at  $\delta_{\rm C}$  166.6 and C-48 at  $\delta_{\rm C}$ 168.7) in the mesaconate moiety suggested that both carbonyls would be esterified.<sup>7</sup> The HMBC correlation of H-13 with C-48 and the downfield chemical shift of H-13  $(\delta_{\rm H} 4.95)$  positioned an ester bond between C-48 and C-13. Siminarly, H-27, another downfield proton ( $\delta_{\rm H}$  4.88), showed HMBC correlation with either carbonyl group C-1 ( $\delta_{\rm C}$  166.62) or C-51 ( $\delta_{\rm C}$  166.61) (due to overlapping signals). The assignment of esterification of C-27 by C-51 was subsequently based on the ROSEY correlation between H<sub>3</sub>-45 and H-50. Finally, there was still one degree of unsaturation unaccounted for, requiring another ring in the final structure. The upfield chemical shift of C-1 indicated it to be an ester carbonyl, which could possibly connect to one of the oxygens left (i.e., O-5, O-11, O-23, O-25, and O-29) to form a macrocyclic ring. The upfield chemical shifts of H-5 ( $\delta_{\rm H}$  3.56), H-11 ( $\delta_{\rm H}$  3.34), H-23 ( $\delta_{\rm H}$ 3.31), and H-25 ( $\delta_{\rm H}$  3.90), however, suggested that all of these positions would unlikely be esterified. Therefore, C-1 had to be connected with C-29 through an ester bond, and this assignment was further supported by the ROESY correlation between H-5 and H-27. This completed the strucral elucidation of neaumycin as a new



Figure 2. Key COSY, HMBC, and ROESY correlations supporting the structures of fragments A, B, C, as well as the mesaconate moiety, and their final assembly into neaumycin.

30-membered macrolide featuring an internal diester bridge (Figures 1 and 2).

The neaumycin structure contained 19 chiral centers. While complete stereochemistry assignment is beyond the scope of the current study, we adopted the *J*-based method,<sup>8</sup> combining with ROESY correlations, to narrow down possible diastereomers. Signal overlap and cross peak broadening hampered the  ${}^{2,3}J_{\rm H,C}$  measurements for H<sub>2</sub>-6 with C-5 or C-7, H-34 with C-35 and C-36, and H-35 with C-34 and C-46. Nevertheless, the relative stereochemistry of four individual coupling systems of C-4/C-5, C-10 to C-14, C-22 to C-33, and C-36/C-37 were established based on the informative  ${}^{2,3}J_{\rm H,C}$  and  ${}^{3}J_{\rm H,H}$  values obtained and key ROESY correlations (Figure 3).

Figure 3A depicted the relative stereochemistry of the C-4/C-5 coupling system. The *anti* orientation of H-4/H-5 was suggested by a large  ${}^{3}J_{\rm H,H}$  (10.5 Hz), while the *gauche* orientations of H-4/HO-C-5 and H-5/C-42 were indicated by a large  ${}^{2}J_{\rm H,C}$  (6.5 Hz) and a small  ${}^{3}J_{\rm H,C}$  (1.9 Hz), respectively. This assignment was supported by ROESY correlations between H-5 and H<sub>3</sub>-42, H-5 and H-3, and H<sup>1</sup>-6 and H-3. The coupling constants between H<sub>2</sub>-6 with C-5 and C-7 were not determined due to the close chemical shifts of  ${}^{1}$ H-6 ( $\delta_{\rm H}$  1.50) and  ${}^{\rm h}$ H-6 ( $\delta_{\rm H}$  1.59). The relative stereochemistry between C-5 and C-7 therefore remains unknown.

Figure 3B depicted the relative stereochemistry of the C-10 to C-14 coupling system. First, the *gauche* orientation of H-10/H-11 was suggested by the small  ${}^{3}J_{H,H}$  (not observed). The *anti* orientation of H-10/HO-C-11 was determined based on the small  ${}^{2}J_{H,C}$  (3.1 Hz). The ROESY correlations between H-9 and H-11 and between H-10 and H-11 supported the *guache* orientation of H-11 to both C-9

<sup>(8) (</sup>a) Matsumori, N.; Kaneno, D.; Murata, M.; Nakamura, H.; Tachibana, K. *J. Org. Chem.* **1999**, *64*, 866–876. (b) Menche, D.; Arikan, F.; Perlova, O.; Horstmann, N.; Ahlbrecht, W.; Wenzel, S. C.; Jansen, R.; Irschik, H.; Muller, R. *J. Am. Chem. Soc.* **2008**, *130*, 14234–14243. (c) Luo, Y.; Shen, B.; et al. *J. Am. Chem. Soc.* **2010**, *132*, 6663–6671. (d) Paterson, I.; Wright, A. E.; et al. *Angew. Chem., Int. Ed.* **2011**, *50*, 3219– 3223.

and H-10. Second, the *anti* orientation of H-11/<sup>1</sup>H-12 was suggested by the large  ${}^{3}J_{H,H}$  (7.2 Hz). The *anti* orientation of <sup>h</sup>H-12/HO-C-11 was consistent with a small  ${}^{3}J_{H,H}$  of H-11 with <sup>h</sup>H-12 (2.3 Hz), a small  ${}^{3}J_{H,C}$  of H-11 with C-13 (2.0 Hz), a small  ${}^{2}J_{H,C}$  of  ${}^{h}H$ -12 with C-11 (1.6 Hz), and a large  ${}^{2}J_{\text{H,C}}$  of  ${}^{1}\text{H}$ -12 with C-11 (6.6 Hz). Similarly, the *anti* orientation of H-13/<sup>h</sup>H-12 was suggested by the large  ${}^{3}J_{\rm H H}$ (10.5 Hz), while the anti orientation of <sup>1</sup>H-12/RO-C-13 was concluded based on a small  ${}^{3}J_{H,H}$  of  ${}^{1}H$ -12 with H-13 (on observed), a small  ${}^{3}J_{H,C}$  of H-13 with C-11 (1.3 Hz), a small  $^{2}J_{\rm H,C}$  of  $^{1}$ H-12 with C-13 (0.5 Hz), and a large  $^{2}J_{\rm H,C}$  of  $^{\rm h}$ H-12 with C-13 (7.8 Hz). The ROSEY correlation between <sup>1</sup>H-12 and H-13 also supported this assignment. Finally, the gauche orientation of H-13/H-14 was suggested by the large  ${}^{2}J_{H,C}$  of H-13 with CH<sub>3</sub>O–C-14 (7.5 Hz) and H-14 with RO-C-13 (6.3 Hz), respectively. This assignment was consistent with the small  ${}^{3}J_{H,H}$  between H-13 and H-14 (not observed) and supported by the ROSEY correlation between H-13 and H-14. Taken together, these observations established the relative configuration from C-10 to C-14.

Figure 3C summarized the assignment of the relative stereochemistry of the C-22 to C-33 coupling system. First, the relative configuration of C-22/C-23 was established by the large  ${}^{3}J_{\text{H,H}}$  and  ${}^{2}J_{\text{H,C}}$  of H-22 with H-23 (8.5 Hz) and C-23 (6.8 Hz), respectively. Second, extension from C-22/ C-23 to C-25 was based on (i) a large  ${}^{3}J_{H,H}$  of H-23 with <sup>h</sup>H-24 (9.7 Hz), a small  ${}^{3}J_{H,C}$  of H-23 with C-25 (2.5 Hz), a small  ${}^{2}J_{H,C}$  of  ${}^{1}$ H-24 with C-23 (0.9 Hz), and a large  ${}^{2}J_{H,C}$  of <sup>h</sup>H-24 with C-23 (5.9 Hz) and (ii) a large  ${}^{3}J_{H,H}$  of  ${}^{1}$ H-24 with C-23 (5.9 Hz) and (ii) a large  ${}^{3}J_{H,H}$  of  ${}^{1}$ H-24 with H-25 (10.1 Hz), a large  ${}^{2}J_{H,C}$  of  ${}^{1}$ H-24 with C-25 (7.4 Hz), and a small  ${}^{2}J_{H,C}$  of  ${}^{h}$ H-24 with C-25 (2.1 Hz), respectively. This assignment was supported by the ROSEY correlations between <sup>1</sup>H-24 and H-22 and between H-23 and H-25. Third, the correlation from C-25 to C-27 was based on the large  ${}^{3}J_{H,H}$  of H-25 with H-26 (9.2 Hz) and H-26 with H-27 (10.5 Hz) and the large  ${}^{2}J_{H,C}$  of H-26 with C-27 (7.5 Hz), which was fully supported by the NOSEY correlations between H-24 and H-45, H-25 and H-45, H-25 and H-27, H-27 and H-45, and H-28 and H-45. Fourth, the relative conformation of C-27/C-28 was based on a large  ${}^{3}J_{H,H}$  of H-27 with  ${}^{1}$ H-28 (12.0 Hz), a small  ${}^{3}J_{H,H}$ of H-27 with <sup>h</sup>H-28 (3.6 Hz), and a large  ${}^{2}J_{\text{H,C}}$  of <sup>1</sup>H-28 with C-27 (6.7 Hz) and supported by the ROESY correlation between <sup>h</sup>H-28 and H-27. Finally, the ROESY correlation between <sup>h</sup>H-28 and H-33 indicated that both C-28 and H-33 are at the axial orientation. As the tetrahydropyran ring is relatively rigid and the C-27 side chain apparently adopted one dominant configuration, the ROESY correlations between <sup>h</sup>H-28 and H-27, <sup>h</sup>H-28 and H<sub>3</sub>-46, and especially H-27 and H-5 led the assignment of a gauche orientation between H-27 and C-29 and an anti orientation between C-27 and O-29/33. Taken together, these observations established the relative stereochemistry from C-22 to C-33.

The relative stereochemistry from C-33 to C-35 was not assigned because the key  ${}^{2,3}J_{\rm H,C}$  constants were not



**Figure 3.** Determination of relative stereochemistry of three individual coupling systems of (A) C-4/C-5, (B) C-10 to C-14, and (C) C-22 to C-33 on the basis of  ${}^{3}J_{\rm H,H}$  and  ${}^{2,3}J_{\rm C,H}$  constants and key ROSEY correlations.

observed. The C-36/C-37 epoxide was established to be *trans* based on the ROESY correlations between H-36 and H-38 and between H-37 and H-35 (Figure 2).

Neaumycin was tested for cytotoxicity against selected cancer cell lines. No significant inhibitory activity was observed at  $1.0 \,\mu\text{M}$  under the conditions tested.

Acknowledgment. We thank the Analytical Instrumentation Center of the School of Pharmacy, UW–Madison, for support in obtaining MS and NMR data and Drs. John Cleveland and Chunying Yang, Scripps Florida for cytotoxicity assays. This work was supported in part by grants from NKPBR (No. 2010CB126102) and NNSF (Nos. 30971937 and 30771427) of China (W.X.) and by NIH grant GM086184 (B.S.).

**Supporting Information Available.** Detailed experimental procedures, HR-ESI-MS spectrum, and assorted 1D and 2D NMR spectra for neaumycin (Table S1 and Figures S1 to S11). This material is available free of charge via the Internet at http://pubs.acs.org.

The authors declare no competing financial interest.