

## Identification of triosephosphate isomerase (TIM) genes from *Microcystis* (Cyanobacteria: Chroococcales) and theoretical analyses of the potential of cyanobacterial TIM as a target for designing specific inhibitors

WANG SHEN<sup>1,2</sup>, JIAN-FAN WEN<sup>1\*</sup>, GUI-LING SUN<sup>1,2</sup> AND YUAN LU<sup>1</sup>

<sup>1</sup>Key Laboratory of Cellular and Molecular Evolution, Kunming Institute of Zoology, Chinese Academy of Sciences, Kunming 650223, People's Republic of China

<sup>2</sup>Graduate School of the Chinese Academy of Sciences, Beijing 100039, People's Republic of China

W. SHEN, J.-F. WEN, G.-L. SUN AND Y. LU. 2005. Identification of triosephosphate isomerase (TIM) genes from *Microcystis* (Cyanobacteria: Chroococcales) and theoretical analyses of the potential of cyanobacterial TIM as a target for designing specific inhibitors. *Phycologia* 44: 447–452.

The genes encoding triosephosphate isomerase (TIM) in three species of *Microcystis* (*M. aeruginosa*, *M. viridis* and *M. wesenbergii*) were investigated. Reverse transcriptase–polymerase chain reaction indicated that they were transcribed in the cells. Analyses showed that their DNA and deduced amino acid sequences were highly conserved between all the three species, only a single nonsynonymous substitution was seen at position 31, from an Asp in *M. aeruginosa* and *M. viridis* to Glu in *M. wesenbergii*. Sequence alignment of these with 12 other known cyanobacterial TIM sequences showed that all the cyanobacterial TIMs had a very high level of amino acid identity (over 50% between each two). Comparison of the cyanobacterial TIMs with other reported TIMs (from diverse lineages of the three Domains) showed that they possessed common active-site residues and sequence motifs. All cyanobacterial TIMs have two common cysteine residues (Cys127 and Cys176), and the Cys176 is almost cyanobacteria-specific with only one exception in *Streptomyces coelicolor*. Both secondary structure alignment and comparative modelling of *Synechocystis* sp. TIM showed that Cys176 was located at the hinge region of the flexible loop-6 and might therefore be critical to the movement of TIM's loop-6, which is important to the function of the enzyme. Thus, the cyanobacterial TIM-specific Cys176 may be a potential site for the discovery of suitable drugs against cyanobacteria, and such drugs may have utility in controlling water blooms due to cyanobacteria.

### INTRODUCTION

Triosephosphate isomerase (TIM) is responsible for the reversible isomerization between glyceraldehyde 3-phosphate and dihydroxyacetone phosphate. This reaction plays an important role in both glycolysis and gluconeogenesis. TIM has the prototypical  $\alpha/\beta$  barrel structural motif. The kinetic properties (Lambeir *et al.* 1987, 1991), three-dimensional structure and catalytic mechanism (Knowles 1991; Lodi *et al.* 1994; Kursula *et al.* 2001) have been described previously in a series of studies.

From a practical application perspective, TIM has been considered as a target for designing inhibitors used as medicines against pathogenic parasites (Garza-Ramos *et al.* 1996, 1998; Velanker *et al.* 1997; Maithal *et al.* 2002). For this purpose, inhibitors of TIM must be highly selective because TIM is a ubiquitous enzyme existing in both hosts and invading organisms. Gomez-Puyou *et al.* (1995) suggested that it was feasible to achieve specific inhibition of homologous enzymes with a high level of selectivity by targeting amino acid residues that were not conserved in evolution but were important for the proper functioning of these enzymes. Using this hypothesis, specific inhibition of TIMs have been studied to restrain some parasitic protozoans such as *Trypanosoma*, *Leishmania* and *Plasmodium* (Garza-Ramos *et al.* 1996, 1998; Velanker *et al.* 1997; Maithal *et al.* 2002).

In recent years, the problem of cyanobacterial blooms,

resulting from water pollution and eutrophication, has become more and more serious (Pitois *et al.* 2001). Some cyanobacteria, such as the toxin-producing strains of *Microcystis*, contribute to cyanobacterial blooms and release toxins, which result in health and environmental hazards in waters (Dawson 1998; Frazier *et al.* 1998; van Buynder *et al.* 2001). Today, the prevention and control of cyanobacterial blooms is still an outstanding unsolved issue. So far, eight cyanobacterial genomes have been sequenced, and only one TIM gene has been identified (GenBank accession number: CAE07338.1, NP\_442075.1, NP\_923986.1, NP\_681756.1, NP\_488425.1, CAE20992.1, NP\_875293.1, CAE19288.1) in each of their genomic databases. In addition, one TIM gene was reported in each of four other cyanobacteria (GenBank accession number: ZP\_00325988.1, ZP\_00164362.2, ZP\_00174972.1, ZP\_00111349.1). Thus, unlike photosynthetic eukaryotes, which have two copies of TIM genes, cyanobacteria have only one copy of the TIM gene in their genomes, and the same TIM functions in both their glycolysis and photosynthesis. This means that if TIM is inhibited, both these metabolic pathways in cyanobacteria will be blocked simultaneously. Therefore, cyanobacterial TIM might be an effective target enzyme for designing cyanobacterial inhibitors to control water blooms.

Prior to this work, no TIM genes were identified in *Microcystis*, which is generally the dominant genus contributing to cyanobacterial blooms (Romanowska-Duda & Tarczynska 2002). It was not known yet whether all cyanobacterial TIMs shares some common features. However, in cyanobacterial

\* Corresponding author (wenjf@mail.kiz.ac.cn).

TIMs, to employ the strategy suggested by Gomez-Puyou *et al.* (1995), it is important to identify an amino acid residue that is not conserved in evolution but plays an important role in the function of the enzyme. In the present work, firstly, TIM genes in three species of *Microcystis* (*M. aeruginosa* Ktzing Lemmermann, *M. viridis* Lemmermann and *M. wesenbergii* Komrek) were amplified, cloned and sequenced. Then, the three *Microcystis* TIMs, together with TIMs of other cyanobacteria and noncyanobacteria, were analyzed to find potential sites for specific inhibitors of cyanobacterial TIMs.

## MATERIAL AND METHODS

### Materials

*Microcystis aeruginosa* (FACHB-315), *M. viridis* (FACHB-930) and *M. wesenbergii* (FACHB-574) were purchased from Algal Culture Collection of the Institute of Hydrobiology, CAS, and were cultured with Endo medium (Lu *et al.* 2001) in sunlight at 20–25°C.

### Amplification, cloning and sequencing of TIM genes

Genomic DNA was extracted from *Microcystis* spp. by a modified phenol–chloroform method (Lu *et al.* 2001). Two pairs of primers were used in polymerase chain reaction (PCR). The first pair was designed by matching two highly conserved regions of all the reported TIM amino acid sequences through alignment, and the nucleotide sequences of the primers were according to TIM gene of *Synechocystis* sp.: Pu1, 16–36: 5'-ATTGCTGGTAATTGGAAAATG-3'; Pd1, 507–488: 5'-AGCCCAAATCGGTTCTAGG-3' (the nucleotide numbers was according to the TIM gene of *Synechocystis* sp.), and was used to amplify an expected segment of ~ 500 bp of TIM genes from *Microcystis* genomic DNA. The PCR was performed as follows: 95°C for 3 min, followed by 40 cycles of 95°C for 1 min, 48°C for 1 min and 72°C for 1 min, followed by 72°C for 10 min. The PCR products of the expected size were purified from 2% agarose gel by using gel extraction kits (Watson Biotechnological Inc., Shanghai, China). The purified products were cloned into pMD 18-T Vectors (TaKaRa Biotechnology (Dalian) Co., Ltd., Dalian, China), and at least five clones of each sample were sequenced by ABI 377 DNA Sequencer (Applied Biosystems, Foster City, CA, USA) with BigDye Terminator Cycle Sequencing Kits (Applied Biosystems, USA). To get much longer sequences, a second pair of primers was designed: Pu2, 236–255: 5'-AAATCTCGGCGGATATGTTA-3' (according to a conserved region of the three *Microcystis* TIM gene segments already obtained through alignment), Pd2, 718–700: 5'-A(AG)TT(AG)AC(AG)AT(TCG)C(TG)(ACG)GC(AG)AA-3' (corresponding to the C-terminal conserved region of the reported cyanobacteria *Synechocystis* sp. and *Nostoc* sp. TIM genes). The PCR, cloning and sequencing were carried out as described above.

### Reverse transcriptase–polymerase chain reaction and PCR with the same primer pair

Total RNA was isolated from axenic *M. viridis* using the UNIQ-10 total RNA minipreps classic kit (Shanghai Sangon Biological Engineering and Technology and Service Co., Ltd.,

Shanghai, China) according to the manufacturer's instructions. The RNA was subsequently incubated with RQ RNase-free DNase (Promega Co., Madison, WI, USA) for 15 min at 37°C to remove contamination of genomic DNA, and terminated by adding Terminator mix. After treatment with chloroform and ethanol, the total RNA was purified with a column and used as templates for reverse transcriptase–polymerase chain reaction (RT-PCR). The reverse transcription reaction (with Pd1 as the RT primer) was carried out at 50°C for 30 min using the TaKaRa RNA PCR Kit (Avian Myeloblastosis Virus (AMV)) V.2.1 (TaKaRa Biotechnology (Dalian) Co., Ltd., Dalian, China) according to the manufacturer's instructions. A control without adding AMV Reverse Transcriptase XL was set simultaneously. Subsequent PCR with the primer pair of Pu1/Pd1 was carried out using the same Kit for 35 cycles at 94°C for 30 s, 48°C for 30 s and 72°C for 45 s. In contrast with the RT-PCR, a PCR with the same primer pair was performed using *M. viridis* genomic DNA as templates. Both the RT-PCR and PCR products were analyzed in the same 1.5% agarose gels.

### Analyses of sequences

DNA sequence analyses were performed by DNASTAR v.5.0 (DNASTAR, Inc., USA). Multiple alignments of nucleotide sequences and amino acid sequences were carried out using CLUSTALW (Thompson *et al.* 1994).

### Modelling of *Synechocystis* sp. TIM

The tertiary structure of mature *Synechocystis* sp. TIM was predicted by comparative modelling using a combination of the Swiss-Pdb Viewer and the SWISS-MODEL server (Peitsch 1995, 1996; Guex & Peitsch 1997). The *Bacillus stearothermophilus* TIM [PDB entry: 1BTM; ExpDB entry: 1BTMA and 1BTMB (Delboni *et al.* 1995)] was chosen as the template to model the *Synechocystis* sp. TIM because it was the best hit and shared 50.2% identity with the *Synechocystis* sp. TIM when using *Synechocystis* sp. TIM as the target sequence to find suitable modelling templates in the ExpDB template database. Optimise Mode method was used to carry out the modelling. First, the sequences alignment between *Synechocystis* sp. TIM and *B. stearothermophilus* TIM was modified manually with Swiss-Pdb Viewer. Then the modelling request was submitted to Swiss-Model server to get the initial model. Fixing  $\alpha$  carbon atoms of the initial model, energy minimization was performed by 500 steps steepest descent and 500 steps conjugate gradient. Root-mean-squared (RMS) values for the topologically equivalent  $\alpha$  carbon atoms between the model and the template were calculated. Finally, protein structure superposition of the modelled *Synechocystis* sp. TIM over *Homo sapiens* TIM (PDB entry: 1HTI), *Saccharomyces cerevisiae* TIM (1YPI, 3YPI) and *Escherichia coli* TIM (1TRE) was performed with the DiCE structural alignment program (Sali & Blundell 1990).

## RESULTS AND DISCUSSION

### Amplification, sequencing and identification of TIM genes in *Microcystis* spp.

With the two pairs of primers (Pu1 and Pd1 for the 5' end of the gene, and Pu2 and Pd2 for the 3' end), two expected over-

lapping PCR products (both about 500 bp), were obtained from all the three species (*M. aeruginosa*, *M. wesenbergii* and *M. viridis*). After sequencing and assembling the two overlapping sequences, pairwise sequence comparisons among the nucleotide sequences of the three species showed that they shared over 98% identity. The deduced amino acid sequences were more highly conserved: only one residue difference was observed, the Asp31 in *M. aeruginosa* and *M. viridis* TIMs was substituted by Glu31 in *M. wesenbergii*. When the three deduced polypeptides were aligned with 153 reported TIM sequences (from diverse lineages of the three Domains, including 47 eukaryotes, 74 eubacteria and 22 archaebacteria) obtained from the NCBI Protein database, a high degree of homology was observed. The three deduced amino acid sequences of *Microcystis* genes had an identity of greater than 50% to each of those sequences from 12 previously reported cyanobacterial sequences. These alignments clearly demonstrated that we had obtained the TIM gene sequences for the three species of *Microcystis* (GenBank accession number AY238889, AY238890 and AY238891). A sequence file with the accession numbers and alignments reported here is available from the authors upon request.


In this work, the first pair of primers was designed to correspond to two highly conserved regions of all known TIM gene sequences, and the PCR products, which showed a single ~ 500 bp band in agarose gels, were very pure; the sequences obtained from five clones were completely identical. All these suggested that, as for the eight cyanobacteria whose genomes had already been sequenced, TIM genes probably occur as a single copy in the three *Microcystis*.

To verify the transcription of the three genes, RT-PCR and PCR with the same primer pair were carried out simultaneously on *M. viridis*. Both of the products showed the same expected size amplicons of about 500 bp (the calculated length is 500 bp) (Fig. 1, Lanes a and c). The negative result of the control of RT-PCR verified that the products of RT-PCR were derived from RNA and not from DNA contamination (Fig. 1, Lane b). This implied that the gene was transcribed in the cells and might not contain introns.

#### Analyses of potential sites of cyanobacterial TIM as targets for designing specific inhibitors

Including the present work, there are now 15 cyanobacterial TIM sequences known. When they were aligned with the other 153 known TIMs mentioned above, all the 15 cyanobacterial TIM sequences were shown to share common features with the other TIMs. For example, they also possessed the conserved active-site residues (Lys14, His96 and Glu165) and some common sequence motifs (part of the alignment results are showed in Fig. 2). In contrast to TIMs of other eubacteria and eukaryotes, the cyanobacterial TIMs had one deletion before the seventh  $\beta$ -strand (see Fig. 2). All the cyanobacterial TIMs had two common cysteine residues (Cys127 and Cys176). Cys127 was also found in other noncyanobacterial TIMs and was a strictly conserved residue of TIMs, but Cys176 was found almost exclusively in cyanobacterial TIMs (with the only exception of TIM of a noncyanobacterium *Streptomyces coelicolor*), and was almost a cyanobacterial TIM-specific cysteine.

TIM is a dimeric glycolytic enzyme, consisting of two iden-



**Fig. 1.** Demonstration of transcription of TIM gene in *Microcystis viridis* by RT-PCR and PCR with the same primer pair flanking the TIM gene. Both the RT-PCR and PCR products showed the same expected size amplicons of about 500 bp. Lane a, RT-PCR with *M. viridis* total RNA as templates; Lane b, RT-PCR with *M. viridis* total RNA as templates but without adding AMV Reverse Transcriptase XL (as negative control); Lane c, PCR with *M. viridis* genomic DNA as templates.

tical monomers. The monomer has a typical TIM-barrel fold with an inner core cylinder of eight parallel  $\beta$ -strands surrounded by a concentric outer cylinder formed by eight  $\alpha$ -helices. The eight central  $\beta$ -strands of the scaffold are labelled from the N-terminus to the C-terminus as  $\beta$ 1–8, the corresponding  $\alpha$ -helices are labelled  $\alpha$ 1–8. The loops connecting the  $\beta$ -strands to the subsequent helices are called loop 1–8. In loop-6 the conserved residues form a rigid lid and the non-conserved residues on both sides of the lid form two hinges (Joseph *et al.* 1990). According to the secondary structural definitions for TIMs of several species (including *Homo sapiens*, *S. cerevisiae*, *E. coli* and *B. stearothermophilus*; PDB entries 1HTI, 1YPI, 1TRE, 1BTM, respectively), the cyanobacterial Cys176 was located in the nonconserved region between the conserved lid region (WAIGTG) and the  $\alpha$ 6 helix, that is, in the hinge region of loop-6.

In our modelling of *Synechocystis* sp. TIM, the final total energy of the *Synechocystis* sp. TIM structure model was  $-13,184.065$  kJ/mol after energy minimization, and the RMS of 240 topologically equivalent  $\alpha$  carbon atoms between the model and the template was  $0.96$  Å when using the DiCE structural alignment program to superpose the modelled *Synechocystis* sp. TIM structure over the A-chain structures of 1HTI, 1YPI, 3YPI, 1TRE mentioned above, the backbone atoms aligned very closely (Fig. 3). All these strongly suggest that the tertiary structural model of *Synechocystis* sp. TIM had the same structural features of all reported crystallographic TIM molecules.

In addition, the model also showed that the cyanobacterial TIM-specific Cys176 was clearly located at the hinge of loop-6 (Fig. 3). The importance of the flexible loop-6 had been


	$\beta$ 1	$\alpha$ -helix 1	$\beta$ 2	$\alpha$ -helix 2	$\beta$ 3	$\alpha$ -helix 3
S. yn	---MRKIIAGNWKMHKTQAEQAFLQGFKPLIEDAAESREVLVCPVFTDLGMSQQLHGGRVRLGAQNVHWEASGAYTGEISAAMLTEI					
N. os	---MRKIVIAGNWKMFKTQAESQEFLEKFLPALEETPQEREVLVCPVFTDLAILSLSQSLHGSLSVQLGAQNVHWAENGAYTGEISGPMLEI					
T. er	-----MYKTQLEAKEFLEGLMSQLTKTPEEREVLCTPFTALDFMSKILHGSLIRLGAQNVHWEDEGAYTGEISGLMLKDV					
M. ae	-----HKTQREALEFLQDFKSHLDETPDDREVVLCPFTALAVLSKTLHGGRIRLGAQNVHWEKSGAYTGEISADMLEI					
P. ma	---MRKPVIAGNWKMMHTCAQAREYTA MF LPIEATPNDRHVVIAPPFTA ISSMAESVGGTRLELSSQN VHWEDD GAYTAEV SPTMLLEH					
B. st	---MRKPIIAGNWKMHKT LAEAVQFVEDVKGHVVP-ADEVISVVCAPFLFLDRLVQAADGTDLKIQAQTMHFADQGAYTGEVSPVMLKDL					
H. sa	MAPSRKFFVGGNWKMNGRKQSLGELIGTLN-AAKVPAD-TEVVCAPPTAYIDFARQKLD-PKIAVAQAQNCYKVTNGAFTGEISPGMIKDC					
S. ce	MA--RTFFVGGNFKLNKSGSKSIKEIVERLN-TASIPEN-VEVVICPPATYLDYSVSLVKKPQVTVGAQNAYLKASGAFTGENSVQDIKDV					
E. co	---MRHPLVMGNWKLNGSRHMVHELVSNLK- <u>KELAGVAGCAVAIAPPEMYIDMAKREAEGSHIMLGAQNV</u> DLNLSGAFGTGETSAAMLKDI					
	$\beta$ 4	$\alpha$ -helix 4	$\beta$ 5	$\alpha$ -helix 5	$\beta$ 6	loop-6
S. yn	GIHYVVIGHSERRQYFGETDETANLRVLAQAQKAGLIPILCVGESKAQRDAGETEQVIVDQVKKGLVNVDS---NLVIAYEPIWAIGTGD					
N. os	GVRVYVIGHSERRQFFGETDETVNLRQAQKYGLTPIILCVGETKQQRDSGETESLIVSGLDKDLINVDQT---NLVIAYEPIWAIGTGD					
T. er	GVSIVVIGHSERRQYFGETDETVNMRKAAQKHGLTPIILCVGETKQQRDSGETESHIFSQLANDLVVDVQDQ---NLVIAYEPIWAIGTGD					
M. ae	GVHYVVIGHSERRQYFGETDETVNLRVISAQKQGLIPIICVGESKAQRDAGETEKVVIKQIQGGLVNVQDQ---NLVIAYEPIWAIGTGE					
P. ma	QVRYAIVGHSEPRKYFSEDEQINRRARSAQAHLIPIVCGESDEQERGEREAERVIRRQVEQGLEDDTDPD---KLVVAYEPIWAIGTGK					
B. st	GVTVVILGHSERRQMF AETDETVNKKVLA AFT RGLIPIICCGESLEEREAGQTN AVVASQVEKALAGLTPEQVQAVIAYEPIWAIGTGK					
H. sa	GATWVVLGHSERRRVFGESDELIGQKVAHALAEGLVIAICIGEKLDEREAGITEKVVFEQTKVIADN--VKDWSKVVLAYEPVWAIGTGK					
S. ce	GAKWVILGHSERRSYFHEDDKFIADKTKFALGQGVVILCIGETLEEKKAGKTLDVVERQLNAVLEE--VKDWTNVVVAIYEPVWAIGTGL					
E. co	GAQYIIIGHSERRTYHKESDELI <del>AKKFAVLKEQGLTPVLCIGETE</del> AENAGKTEEV <del>CARQIDAVLKTQ</del> GA <del>AAFE</del> GAVIAYEPIWAIGTGK					
	* $\alpha$ -helix 6	$\beta$ 7	$\alpha$ -helix 7	$\beta$ 8	$\alpha$ -helix 8	
S. yn	TCAATEANRVIGLIREQLT-----NSQVTIQYGGSVNANNVDEIMAQPEIDGALVGGASLEPQSFARIVNFQP-----					
N. os	TCETTEANRVIGLIRSQK-----NSDVPIQYGGSVKPNNIDEIMAQPEIDGVLVGGASLEAASFARIVNYL-----					
T. er	TCEAKEANRVIGLIRGKLN-----NSNVTIQYGGSVKPNNVDDIMAQEEIDGALVGGASLNPESFSRLVNYQ-----					
M. ae	TCESEANRVIGLIRQQLD-----NPEVTIQYGGSVKPDNIDEIMAQSQINGALVGGASLDPAV-----					
P. ma	TCEASEANRICGLIRRVWG-----ASELIIQYGGSVKPGNIDELMAMSDIDGVLVGGASLDPEFARIVNYQTS-----					
B. st	SSTPEDANSVCGHIRSVSRLFGPEAAEAIRIQYGGSVKPDNIRDFLAQQQIDGPLVGGASLEPASFLQLVEAGRHE---					
H. sa	TATPQQAQVEHKLGRWLKSNVSDAVAQSTRIIYGGSVTGATCKELASQPDVDGFLVGGASLKP-EFVDIINAKQ-----					
S. ce	AATPEDAQDIHASIRKFLASKLGDKAASELRILYGGSSANGSNAVTFKDKADVDFLGGASLKP-EFVDIINSRN-----					
E. co	SATPAQAQAVHKFIRDHIA-KVDANIAEQVIIQYGGSVNANNAELFAQPDIDGALVGGASLKADAFVIVKAAEAAKQA					

**Fig. 2.** Partial results of TIM sequence alignment. Five cyanobacteria: *Synechocystis* sp. PCC 6803 (S.yn), *Nostoc* sp. PCC 7120 (N.os), *Trichodesmium erythraeum* IMS101 (T.er), *Microcystis aeruginosa* (M.ae) and *Prochlorococcus marinus* str. MIT 9313 (P.ma); two nonphotosynthetic bacteria: *Escherichia coli* (E.co) and *Bacillus stearothermophilus* (B.st); two eukaryotes: *Homo sapiens* (H.sa) and *Saccharomyces cerevisiae* (S.ce). Cys176 of *Synechocystis* sp. TIM is indicated with an asterisk (\*) above the sequence. The secondary structural elements of the three-dimensional structure of TIMs of *H. sapiens* (1HTI), *S. cerevisiae* (1YPI), *E. coli* (1TRE), *B. stearothermophilus* (1BTM) and *Synechocystis* sp. (modelled) are labelled with underlines (thin line,  $\alpha$ -helix; bold line  $\beta$ -strand) and shadows (loop-6).

demonstrated very clearly in previous work. The proposed role of this loop is to ensure efficient throughput of substrate to product and to stabilize the reaction intermediate (*cis*-enediolate) (Knowles 1991). During the catalysis, the binding of substrates or substrate analogues was accompanied by a change in the conformations of loops-5, 6 and 7. The most remarkable change occurred in loop-6 – the tip of the loop undergoes a C $\alpha$  movement of about 7 Å in response to ligand binding to provide new hydrogen bonds to the substrate phosphate group, and the motion of the loop-6 lid was caused by the change of dihedral angles of hinge residues (Joseph *et al.* 1990). These imply that the hinge residues of loop-6 in TIM play a pivotal role in catalysis.

In our work, both the secondary structural comparisons and

the modelled *Synechocystis* sp. TIM structure indicated that the Cys176 was just located at an important region, the hinge of loop-6. The thiol group of cysteine can be modified with many chemical agents, for example thiol nucleophiles. Such modifications can affect the structure and function of proteins. Thus, cysteine has often been considered to be a target residue for designing enzyme inhibitors. As mentioned in the introduction, there have been some studies showing that some molecules could successfully selectively inactivate TIMs from some parasites by acting on a certain Cys (Garza-Ramos *et al.* 1998; Maithal *et al.* 2002; Tellez-Valencia *et al.* 2002). Cyanobacterial TIM Cys176 may be such a target residue. Because, on the basis of the analysis above, Cys176 meets the criteria suggested by Gomez-Puyou *et al.* (1995), that is,



**Fig. 3.** Result of superposition of TIM structures. The protein structure of the modelled *Synechocystis* sp. TIM was superposed with the TIM A-chain structures of *Homo sapiens* (1HTI), *Saccharomyces cerevisiae* (1YPI, 3YPI) and *Escherichia coli* (1TRE). The loop-6 of 1HTI, 1YPI and 1TRE are shown in an opened state; the loop-6 of 3YPI and the modelled *Synechocystis* sp. TIM are shown in a closed state. Cys176 of *Synechocystis* sp. TIM (displayed with balls) is located at the hinge of loop-6. (viewed by WebLab ViewerLite, Molecular Simulations Inc., San Diego, CA, USA)

it is not a conserved amino acid residue, rather it is a cyanobacteria TIM-specific one, and it plays an important role in the function of TIM. Therefore, we suggest that Cys176 is suitable choice as a target residue for designing cyanobacteria TIM-specific inhibitors. Although Cys176 also exists in TIM of *S. coelicolor*, this will not affect the use of such inhibitors in the control of water blooms. Certainly, further work is needed to validate this potential target site, which includes designing, synthesis and testing of inhibitors that specifically interact with cyanobacterial TIM Cys176.

## ACKNOWLEDGEMENTS

We are grateful to Prof. Xiufan Shi and Mr Jun Lin (Kunming Institute of Zoology, CAS) for their helpful discussion. This work was supported by Grants (XiBu ZhiGuang Project 2004 and KSCX2-SW-101C) to J.-F.W. from CAS, a Grant (2000YP19) to J.-F.W. from Yunnan Province and a Grant (TF051795) from GEF-MSP.

## REFERENCES

DAWSON R.M. 1998. The toxicology of microcystins. *Toxicon* 36: 953–962.

DELBONI L.F., MANDE S.C., RENTIER-DEL RUE F., MAINFROID V., TURLEY S., VELLIEUX F.M., MARTIAL J.A. & HOL W.G. 1995. Crystal structure of recombinant triosephosphate isomerase from *Bacillus stearothermophilus*. An analysis of potential thermostability factors in six isomerases with known three-dimensional structures points to the importance of hydrophobic interactions. *Protein Science* 4: 2594–2604.

FRAZIER K., COLVIN E., POURIA S., HULLINGER G. & GARCIA R. 1998. Microcystin toxicosis in cattle due to overgrowth of blue-green algae. *Veterinary and Human Toxicology* 40: 23–24.

GARZA-RAMOS G., PEREZ-MONTFORT R., ROJO-DOMINGUEZ A., DE GOMEZ-PUYOU M.T. & GOMEZ-PUYOU A. 1996. Species-specific inhi-

bition of homologous enzymes by modification of nonconserved amino acids residues. The cysteine residues of triosephosphate isomerase. *European Journal of Biochemistry* 241: 114–120.

GARZA-RAMOS G., CABRERA N., SAAVEDRA-LIRA E., DE GOMEZ-PUYOU M.T., OSTOA-SALOMA P., PEREZ-MONTFORT R. & GOMEZ-PUYOU A. 1998. Sulfhydryl reagent susceptibility in proteins with high sequence similarity—triosephosphate isomerase from *Trypanosoma brucei*, *Trypanosoma cruzi* and *Leishmania mexicana*. *European Journal of Biochemistry* 253: 684–691.

GOMEZ-PUYOU A., SAAVEDRA-LIRA E., BECKER I., ZUBILLAGA R.A., ROJO-DOMINGUEZ A. & PEREZ-MONTFORT R. 1995. Using evolutionary changes to achieve species-specific inhibition of enzyme action—studies with triosephosphate isomerase. *Chemistry and Biology* 62: 847–855.

GUEX N. & PEITSCH M.C. 1997. SWISS-MODEL and the Swiss-PdbViewer: an environment for comparative protein modeling. *Electrophoresis* 18: 2714–2723.

JOSEPH D., PETSCH G.A. & KARPLUS M. 1990. Anatomy of a conformational change: hinged “lid” motion of the triosephosphate isomerase loop. *Science* 249: 1425–1428.

KNOWLES J.R. 1991. Enzyme catalysis: not different, just better. *Nature* 350: 121–124.

KURSULA I., PARTANEN S., LAMBEIR A.M., ANTONOV D.M., AUGUSTYNS K. & WIERENGA R.K. 2001. Structural determinants for ligand binding and catalysis of triosephosphate isomerase. *European Journal of Biochemistry* 268: 5189–5196.

LAMBEIR A.M., OPPERDOES F.R. & WIERENGA R.K. 1987. Kinetic properties of triose-phosphate isomerase from *Trypanosoma brucei*. A comparison with the rabbit muscle and yeast enzymes. *European Journal of Biochemistry* 168: 69–74.

LAMBEIR A.M., LOISEAU A.M., KUNTZ D.A., VELLIEUX F.M., MICHELS P.A. & OPPERDOES F.R. 1991. The cytosolic and glycosomal glyceraldehyde-3-phosphate dehydrogenase from *Trypanosoma brucei*. Kinetic properties and comparison with homologous enzymes. *European Journal of Biochemistry* 198: 429–435.

LODI P.J., CHANG C.L., KNOWLES J.R. & KOMIVES E.A. 1994. Triosephosphate isomerase requires a positively charged active site: the role of lysine-12. *Biochemistry* 33: 2809–2814.

LU Y., WEN J.F. & LÜ T.W. 2001. Isolation, pure cultivation and total DNA extraction of *Microcystis aeruginosa* kutz in Dianchi Lake. *Journal of Lake Science* 13: 285–288. In Chinese.

MAITHAL K., RAVINDRA G., BALARAM H. & BALARAM P. 2002. Inhibition of *Plasmodium falciparum* triose-phosphate isomerase by chemical modification of an interface cysteine. Electrospray ionization mass spectrometric analysis of differential cysteine reactivities. *Journal of Biological Chemistry* 277: 25106–25114.

PEITSCH M.C. 1995. Protein modeling by E-mail. *Bio-Technology* 13: 658–660.

PEITSCH M.C. 1996. ProMod and Swiss-Model: internet-based tools for automated, comparative protein modelling. *Biochemical Society Transactions* 24: 274–279.

PITOIS S., JACKSON M.H. & WOOD B.J. 2001. Sources of the eutrophication problems associated with toxic algae: an overview. *Journal of Environmental Health* 64: 25–32.

ROMANOWSKA-DUDA Z. & TARCZYNSKA M. 2002. The influence of microcystin-LR and hepatotoxic cyanobacterial extract on the water plant *Spirodela oligorrhiza*. *Environmental Toxicology* 17: 434–440.

SALI A. & BLUNDELL T.L. 1990. Definition of general topological equivalence in protein structures. A procedure involving comparison of properties and relationships through simulated annealing and dynamic programming. *Journal of Molecular Biology* 212: 403–428.

TELLEZ-VALENCIA A., AVILA-RIOS S., PEREZ-MONTFORT R., RODRIGUEZ-ROMERO A., TUENA DE GOMEZ-PUYOU M., LOPEZ-CALAHORRA F. & GOMEZ-PUYOU A. 2002. Highly specific inactivation of triosephosphate isomerase from *Trypanosoma cruzi*. *Biochemical and Biophysical Research Communications* 295: 958–963.

THOMPSON J.D., HIGGINS D.G. & GIBSON T.J. 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence align-

ment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Research* 22: 4673–4680.

VAN BUYNDER P.G., OUGHTRED T., KIRKBY B., PHILLIPS S., EAGLESHAM G., THOMAS K. & BURCH M. 2001. Nodularin uptake by seafood during a cyanobacterial bloom. *Environmental Toxicology* 16: 468–471.

VELANKER S.S., RAY S.S., GOKHALE R.S., SUMA S., BALARAM H., BAL-

ARAM P. & MURTHY M.R. 1997. Triosephosphate isomerase from *Plasmodium falciparum*: the crystal structure provides insights into antimalarial drug design. *Structure* 5: 751–761.

*Received 12 February 2004; accepted 3 April 2005*  
*Communicating editor: J. Beardall*