

Three new koninginins from *Trichoderma neokongii* 8722



Xiao-Xue Zhou^{a,b}, Jing Li^{a,b}, Yin-He Yang^a, Ying Zeng^a, Pei-Ji Zhao^{a,*}

^a State Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650204, China

^b University of Chinese Academy of Sciences, Beijing 100049, China

ARTICLE INFO

Article history:

Received 9 January 2014

Received in revised form 25 February 2014

Accepted 4 March 2014

Available online 2 April 2014

Keywords:

Trichoderma neokongii

Chemical constituents

Structure identification

Koninginin

ABSTRACT

Three new fungal metabolites, named koninginins I (**1**), J (**2**) and K (**3**) together with four known koninginins A (**4**), B (**5**), D (**6**) and E (**7**), were isolated from solid fermentation products of *Trichoderma neokongii* 8722. Three new structures were elucidated by extensive spectroscopic methods, including 1D NMR and 2D NMR, and HR-ESI-MS experiments.

© 2014 Phytochemical Society of Europe. Published by Elsevier B.V. All rights reserved.

1. Introduction

Trichoderma can produce a lot of secondary metabolites using for biological control (Reino et al., 2008; El-Hasan et al., 2009; Stoppacher et al., 2010; Mukherjee et al., 2012). A new polyketide type compound, koninginin, was obtained from *Trichoderma* sp. (Reino et al., 2008). Koninginins A, B, C, E and G showed a growth inhibition of etiolating wheat coleoptiles (Cutler et al., 1989, 1991, 1999; Parker et al., 1995a, 1995b). Koninginins A, B, D and G also could affect the growth of some plant pathogen fungi (Dunlop et al., 1989; Ghisalberti and Rowland, 1993; Cutler et al., 1999). In order to search for new structure koninginin type compounds against plant diseases, the extract of *Trichoderma neokongii* 8722 was investigated and three new koninginin type compounds, together with four known koninginins were obtained. This report describes three new compounds structures.

2. Results and discussion

From the extracts of solid fermentation products of *T. neokongii* 8722, seven compounds including three new koninginins I–K (**1–3**) (Fig. 1) were identified. The structures of the known compounds were determined to be koninginins A (**4**) (Cutler et al., 1989), B (**5**) (Cutler et al., 1991), D (**6**) (Dunlop et al., 1989; Song et al., 2010), and E (**7**) (Parker et al., 1995a).

Compound **1** was obtained as colorless amorphism. The HR-ESI-MS data indicated a molecular formula of C₁₆H₂₄O₅ based on the [M+H]⁺ ion signal at *m/z* 297.1699 (calc. 297.1697). The NMR data (Table 1) revealed one quaternary carbon at δ_C 199.5, 175.1 and 112.9, five methines at δ_C 67.1 (δ_H 4.39), 66.9 (δ_H 4.27), 78.1 (δ_H 4.11), 73.4 (δ_H 3.70) and 68.7 (δ_H 3.73), and seven methylenes and one methyl, which suggested compound **1** was koninginin type compound (Dunlop et al., 1989; Reino et al., 2008). According to the NMR and MS spectra, compound **1** had one more hydroxyl than koninginin D (Cutler et al., 1991; Song et al., 2010). A preliminary linear skeleton bearing two branches was deduced to be C-2–C-3–C-4 (–branch)–C-7–C-8–C-9–C-10–C-11 (–branch) from complete interpretation of key cross-peaks in the COSY spectrum (H-2/H-3/H-4; H-7/H-8/H-9/H-10/H-11) and key correlations in the HMBC spectrum: H-2 (δ_H 2.32 and 2.62) correlated with the carbons at δ_C 199.5 (C-1), 30.6 (C-3), 67.1 (C-4), H-3 (δ_H 1.97 and 2.18) with the carbons at δ_C 199.5 (C-1), 34.3 (C-2), 67.1 (C-4) and 175.1 (C-5), H-4 (δ_H 4.39) with the carbons at δ_C 175.1 (C-5), 112.9 (C-2), 34.3 (C-2) and 30.6 (C-3), and H-7 (δ_H 4.27) with the carbons at δ_C 199.5 (C-1), 175.1 (C-5), 112.9 (C-6), 78.1 (C-9), and 68.7 (C-15) (w), H-9 (δ_H 4.11) with the carbons at δ_C 66.9 (C-7) and 28.6 (C-8), H-16 (δ_H 1.16) with the carbons at δ_C 40.3 (C-14), 68.7 (C-15), and other correlations (Fig. 2). In the ¹H NMR spectrum, two intermediate coupling constants (*J* = 4.8, 6.6 Hz) were observed for H-4, which confirmed a pseudoequatorial position for H-4. Except for the signals of H-15 and C-15 for **1**, the NMR data were very similar to those of koninginin D (Dunlop et al., 1989; Song et al., 2010). According to a biogenetic perspective, and comparing the specific rotation, chemical shift and coupling constant of **1** with that of the

* Corresponding author. Tel.: +86 871 65223111.

E-mail address: zhaopeiji@mail.kib.ac.cn (P.-J. Zhao).




Fig. 1. The structures of compounds 1–3.




Fig. 2. HMBC and key correlations of compounds 1–2.

literature data (Song et al., 2010), the relative configuration of **1** was same with koniginin D (Dunlop et al., 1989; Song et al., 2010). Based on above data, compound **1** was elucidated to be koniginin I.

Compound **2** was obtained as colorless powder. The HR-ESI-MS data indicated a molecular formula of C₁₆H₂₄O₅ based on the [M+Na]⁺ ion signal at *m/z* 319.1519 (calc. 319.1516). The MS and NMR spectroscopic data of compound **2** were very similar to those of koniginin B except that one methylene (C-15) in koniginin B was oxidized to the ketone in compound **2** (Cutler et al., 1991; Liu and Wang, 2001). The 2D-NMR data (Table 2) showed the detail: H-2 (δ_H 2.14) correlated with the carbons at δ_C 209.0 (C-15) and 43.5 (C-14), H-14 (δ_H 2.46) with the carbons at δ_C 209.0 (C-15), 30.0 (C-16) and 23.4 (C-13), and other correlations (Fig. 2). In the ¹H NMR spectrum, two intermediate coupling constants (*J* = 6.8, 16.4 Hz) were observed for H-2, which confirmed a axial position for H-2. Except that one methylene (C-15) in koniginin B was oxidized to the ketone in compound **2**, the NMR data were very similar to those of koniginin B (Cutler et al., 1991; Liu and Wang, 2001). From a biogenetic perspective, the configuration of **2** should be identical to that of the co-occurring koniginin B. So, compound **2** was identified to be koniginin J.

Compound **3** was obtained as colorless oil. The HR-ESI-MS data indicated a molecular formula of C₁₆H₂₄O₆ based on the [M+Na]⁺ ion signal at *m/z* 335.1466 (calc. 335.1465). The MS and NMR spectroscopic data of compound **3** were very similar to those of koniginin B except that terminal methyl (CH₃-16) in koniginin B was oxidized to the carboxyl group in compound **3** (Cutler et al., 1991; Liu and Wang, 2001). The 2D-NMR data (Table 2) showed the correlations between H-14 (δ_H 1.65) and carbons at δ_C 178.3 (C-16), 33.8 (C-15) and 29.1 (C-13), H-15 (δ_H 2.35) and carbons at δ_C 178.3 (C-16), 33.8 (C-15) and 24.7 (C-14). In the ¹H NMR spectrum, two intermediate coupling constants (*J* = 6.0, 18.0 Hz) were observed for H-2, which confirmed a axial position for H-2. Except that

terminal methyl (CH₃-16) in koniginin B was oxidized to the carboxyl group in compound **3**, the NMR data were very similar to those of koniginin B (Cutler et al., 1991; Liu and Wang, 2001). From a biogenetic perspective, the configuration of **3** should be identical to that of the co-occurring koniginin B. So, compound **3** was identified to be koniginin K.

Compounds **1**–**7** were assayed for antifungal activity (*Gaeumannomyces graminis*, *Fusarium moniliforme*, *Verticillium cinnabarium* and *Phyricularia oryzae*), but all compounds did not show the inhibition activity to the tested phytopathogenic fungi at 100 μ g/disk. Nematicidal activity result indicated that only koniginin A (**4**) had weak activity against *Panagrellus redivivus* and *Caenorhabditis elegans*.

3. Experimental

3.1. General

UV spectra were measured on a Shimadzu UV-2401PC spectrophotometer, λ_{max} (log ϵ) in nm. NMR experiments were carried out on Bruker AM-400 and Bruker DRX-500 NMR spectrometers with TMS as internal standard. ESI-MS and HR-ESI-MS were recorded on a Finnigan LCQ-Advantage mass spectrometer and a VG Auto-Spec-3000 mass spectrometer. Optical rotations were measured on a Jasco DIP-370 digital polarimeter. Column chromatography was carried out on silica gel (G, 200–300 mesh and H, Qingdao Marine Chemical Factory, Qingdao, China), and Sephadex LH-20 (Pharmacia). Precoated silica gel GF254 plates (Qingdao Marine Chemical Factory, Qingdao, China) were used for thin layer chromatography (TLC).

3.2. Fungal material

The strain of *T. neokongii* 8722 was deposited in Southwest Forestry University. Four pathogenic fungi (*Gaeumannomyces graminis*, *Fusarium moniliforme*, *Verticillium cinnabarium* and *Phyricularia oryzae*) were provided by Dr. Fan L. M. at Yunnan Agricultural University. The culture medium consisted of potato (peeled, 200 g), agar (15 g) and glucose (20 g), per L of deionized H₂O. *T. neokongii* 8872 (20 L) was cultured on potato-dextrose agar dish at temperature of 26 °C for 21 days.

3.3. Extraction and isolation

Solid fermentation products of *T. neokongii* 8722 (20 L) was cut into small pieces and extracted with mixture solution (EtOAc:MeOH:HAc = 80:15:5, v/v/v) by three times to afford of rude extracts. The extracts were dissolved in water, and extracted with EtOAc and then *n*-butanol three times, respectively.

The EtOAc (31.0 g) residue was subjected to a column of silica gel G (200–300 mesh) using petroleum ether–EtOAc and CHCl₃–MeOH gradient solvent system to produce 11 fractions (Fr.1–Fr.11). The fraction Fr.5 (320 mg) was subjected to Sephadex LH-20 CHCl₃–MeOH (1:1) and subsequent purified by preparative TLC to give compound **5** (11 mg). The fraction Fr.8 (1.70 g) was subjected

Table 1
NMR data of compounds **1** (in CD₃OD, *J* in Hz).

Position	¹ H	¹³ C	HMBC
1	–	199.5	–
2	2.32 (1H, ddd, 4.8, 8.4, 16.8)	34.3	1, 3, 4
	2.62 (1H, ddd, 4.8, 7.8, 16.8)		1, 3, 4
3	1.97 (1H, m)	30.6	1, 2, 4
	2.18 (1H, m)		1, 2, 4
4	4.39 (1H, dd, 4.8, 6.6)	67.1	2, 3, 5, 6
5	–	175.1	–
6	–	112.9	–
7	4.27 (brs)	66.9	1, 5, 6, 9, 15 (w)
8	2.06 (1H, m)	28.6	6, 7
	1.57 (1H, m)		9
9	4.11 (1H, m)	78.1	7, 8, 5 (w)
10	3.70 (1H, m)	73.4	11, 12
11	1.66 (2H, m)	33.8	9, 10, 12
12	1.54 (2H, m)	26.9	9
13	1.44 (2H, m)	27.0	12, 14, 15
14	1.44 (2H, m)	40.3	12, 15
15	3.73 (1H, m)	68.7	13
16	1.16 (3H, d, 6.18)	23.7	14, 15

Table 2NMR data of compounds **2** and **3** (in CDCl₃, *J* in Hz).

Position	2			3		
	¹ H	¹³ C	HMBC	¹ H	¹³ C	HMBC
1	–	198.1	–	–	198.4	–
2	4.04 (1H, dd, 6.8, 16.4)	71.0	1, 3	4.04 (1H, dd, 6.0, 18.0)	71.2	1, 2
3	1.79 (1H, m)	28.9	1, 2, 4	1.78 (1H, ddd, 5.4, 12.6, 25.4)	29.2	1, 2, 4
	2.36 (1H, m)		1, 2, 4	2.34 (1H, m)		1, 2, 4, 5
4	2.48 (1H, m)	27.1	5, 6	2.46 (1H, m)	27.3	2, 3, 5, 6
	2.58 (1H, m)		Overlap	2.55 (1H, m)		Overlap
5	–	171.2	–	–	171.5	–
6	–	109.1	–	–	109.4	–
7	2.09 (1H, m)	17.6	8	2.07 (1H, m)	17.8	6, 8
	2.58 (1H, m)		Overlap	2.55 (1H, m)		Overlap
8	1.65 (1H, m)	22.6	7, 9	1.63 (1H, m)	22.9	7, 9
	1.95 (1H, dd, 8.2, 17.2)		6, 7	1.94 (1H, dd, 6.6, 13.8)		6, 7
9	3.77 (1H, m)	80.7	7, 10, 11	3.75 (1H, ddd, 1.9, 5.3, 11.3)	81.0	7, 10
10	3.65 (1H, m)	72.8	9, 11, 12	3.62 (1H, m)	73.3	8, 11, 12
11	1.57 (1H, m)	32.5	9, 10, 12	1.55 (2H, m)	32.7	Overlap
	2.57 (1H, m)		9, 10, 12			
12	1.41 (1H, m)	25.0	–	1.40 (1H, m)	25.2	11, 13
	1.52 (1H, m)		–	1.55 (1H, m)		Overlap
13	1.64 (2H, m)	23.4	–	1.38 (2H, m)	29.1	12
14	2.46 (2H, t, 8.7)	43.5	13, 15	1.65 (2H, m)	24.7	12, 13, 15, 16
15	–	209.0	–	2.35 (2H, t, 7.2)	33.8	13, 14, 16
16	2.14 (3H, s)	30.0	14, 15	–	178.3	–

on silica gel G eluted with a gradient solvent system of petroleum ether–EtOAc to provide compound **4** (60 mg). The fraction Fr.10 (2.10 g) was eluted with a gradient solvent system of petroleum ether–EtOAc to give fractions of Fr.10.1–Fr.10.7. Fr.10.5 was repeatedly purified by silica gel G and eluted with petroleum ether–EtOAc to obtain compound **7** (6 mg). Fr.10.7 was further purified by preparative TLC to give compound **6** (65 mg). The fraction of Fr.10.6 (912 mg) was separated by semipreparative HPLC (LC3000 Semi-preparation Gradient HPLC System, China) to yield compounds **2** (4.0 mg) (Sample was performed in a RP-C₁₈ column (250 mm × 10 mm) at ambient temperature with a detection wavelength at 254 nm and a mobile phase of methanol/water (the water reduce from 60% to 0%) at a flow rate of 3 ml/min). The application of Sephadex LH-20 CHCl₃–MeOH (1:1) helps us to obtain the new compound **3** (2 mg). The *n*-butanol residue (20.1 g) was subjected to a column of silica gel (200–300 mesh) using CHCl₃–MeOH gradient solvent system to produce 3 fractions (Fbr.1 to Fbr.3). The Fbr.2 (120 mg) was subjected to silica gel H using CHCl₃–MeOH gradient solvent system and then purified by Sephadex LH-20 (Acetone) to produce compound **1** (5 mg).

3.3.1. Koninginin I (**1**)

Colorless amorphism: $[\alpha]_D^{18} = +96.36$ ($c = 0.63$, MeOH); UV (MeOH) λ_{\max} (log ϵ): 254 (4.04); NMR data see Table 1; ESI–MS: 297 [M+H]⁺; HR–ESI–MS: 297.1699 ([M+H]⁺, calc. 297.1697).

3.3.2. Koninginin J (**2**)

Colorless powder: $[\alpha]_D^{18} = +103.67$ ($c = 0.80$, MeOH); UV (MeOH) λ_{\max} (log ϵ): 199.6 (3.45), 261 (3.89); NMR data see Table 2; ESI–MS: 319 [M+Na]⁺; HR–ESI–MS: 319.1519 ([M+Na]⁺, calc. 319.1516).

3.3.3. Koninginin K (**3**)

Colorless amorphism: $[\alpha]_D^{18} = +169.42$ ($c = 0.15$, MeOH); UV (MeOH) λ_{\max} (log ϵ): 199 (3.59), 261 (4.08); NMR data see Table 2; ESI–MS: 335 [M+Na]⁺; HR–ESI–MS: 335.1466 ([M+Na]⁺, calc. 335.1465).

3.4. Assay activities

The antifungal activity was assessed against phytopathogenic fungi (*G. graminis*, *F. moniliforme*, *V. cinnabarium* and *P. oryzae*)

using disk diffusion method (Espinell-Ingroff et al., 1999). The method of nematocidal activity against *Panagrellus redivivus* and *Caenorhabditis elegans* was based on the literature (Li et al., 2005).

Acknowledgments

This work was supported by grants from the “973” Program of China (2013CB127505), the NSFC (31170061), and Applied Basic Research Foundation of Yunnan Province (2013FA018), the Young and Technical Leader Raising Foundation of Yunnan Province (2009CI071) and Program of Tobacco Corporation (HYHH2012CL04). We acknowledge the Department of Instrumental Analysis of Kunming Institute of Botany for measuring the optical rotations, UV, NMR and mass spectra.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.phytol.2014.03.004>.

References

- Cutler, H.G., Cutler, S.J., Ross, S.A., Sayed, K.E., Dugan, F.M., Bartlett, M.G., Parker, S.R., 1999. Koninginin G, a new metabolite from *Trichoderma aureoviride*. *J. Nat. Prod.* 62 (1) 137–139.
- Cutler, H.G., Himmelsbach, D.S., Arrendale, R.F., Cole, P.D., Cox, R.H., 1989. Koninginin A: a novel plant growth regulator from *Trichoderma koningii*. *Agric. Biol. Chem.* 53 (10) 2605–2611.
- Cutler, H.G., Himmelsbach, D.S., Yagen, B., Arrendale, R.F., Jacyno, J.M., Cole, P.D., Cox, R.H., 1991. Koninginin B: a biologically active congener of koninginin A from *Trichoderma koningii*. *J. Agric. Food Chem.* 39 (5) 977–980.
- Dunlop, R.W., Simon, A., Sivasithamparan, K., Ghisalberti, E.L., 1989. An antibiotic from *Trichoderma koningii* active against soilborne plant pathogens. *J. Nat. Prod.* 52 (1) 67–74.
- El-Hasan, A., Walker, F., Schöne, J., Buchenauer, H., 2009. Detection of viridifungin A and other antifungal metabolites excreted by *Trichoderma harzianum* active against different plant pathogens. *Eur. J. Plant Pathol.* 124 (3) 457–470.
- Espinell-Ingroff, A., White, T., Pfaller, M.A., 1999. Antifungal agents and susceptibility tests. In: Murray, P.R. (Ed.), *Manual of Clinical Microbiology*. 7th edn. American Society for Microbiology, Washington, DC, pp. 1640–1652.
- Ghisalberti, E.L., Rowland, C.Y., 1993. Antifungal metabolites from *Trichoderma harzianum*. *J. Nat. Prod.* 56 (10) 1799–1804.
- Li, G.H., Shen, Y.M., Zhang, K.Q., 2005. Nematicidal activity and chemical component of *Poria cocos*. *J. Microbiol.* 43, 17–20.
- Liu, G., Wang, Z., 2001. Total synthesis of koninginin D, B and E. *Synthesis* 2001 (01) 0119–0127.

- Mukherjee, M., Mukherjee, P.K., Horwitz, B.A., Zachow, C., Berg, G., Zeilinger, S., 2012. *Trichoderma*-plant-pathogen interactions: advances in genetics of biological control. *Indian J. Microbiol.* 52 (4) 522–529.
- Parker, S.R., Cutler, H.G., Schreiner, P.R., 1995a. Koninginin E: isolation of a biologically active natural product from *Trichoderma koningii*. *Biosci. Biotechnol. Biochem.* 59 (9) 1747–1749.
- Parker, S.R., Cutler, H.G., Schreiner, P.R., 1995b. Koninginin C: a biologically active natural product from *Trichoderma koningii*. *Biosci. Biotechnol. Biochem.* 59 (6) 1126–1127.
- Reino, J.L., Guerrero, R.F., Hernandez-Galan, R., Collado, I.G., 2008. Secondary metabolites from species of the biocontrol agent *Trichoderma*. *Phytochem. Rev.* 7 (1) 89–123.
- Song, F., Dai, H., Tong, Y., Ren, B., Chen, C., Sun, N., Zhang, L., 2010. *Trichoderma* ketones A-D and 7-O-methylkoninginin D from the marine fungus *Trichoderma koningii*. *J. Nat. Prod.* 73 (5) 806–810.
- Stoppacher, N., Kluger, B., Zeilinger, S., Krska, R., Schuhmacher, R., 2010. Identification and profiling of volatile metabolites of the biocontrol fungus *Trichoderma atroviride* by HS-SPME-GC-MS. *J. Microbiol. Methods* 81 (2) 187–193.