

Dammarane-Type Glycosides from Steamed
NotoginsengPENG-YING LIAO,^{†,‡} DONG WANG,[†] YING-JUN ZHANG,^{*,†} AND
CHONG-REN YANG[†]

State Key Laboratory of Phytochemistry and Plant Resources of West China, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650204, People's Republic of China, and Graduate School of the Chinese Academy of Sciences, Beijing 100039, People's Republic of China

Notoginseng, the root of *Panax notoginseng* (Burk.) F. H. Chen (Araliaceae), is a Chinese traditional medicine, which is used in both raw and processed forms due to their different pharmacological activities. Detailed chemical investigation on steamed notoginseng led to the isolation of 27 dammarane-type triterpenoids (1–27), including 4 new glycosides, namely, notoginsenosides ST-1–ST-3 and ST-5 (1–4), in addition to 3 other known compounds. Of the new compounds, 1–3 possess new aglycones. Their structures were elucidated on the basis of detailed analyses of their 1D and 2D NMR spectra and chemical reactions. The known compounds, koryoginsenoside-R₁ (19), yesanichinoside D (20), 6''-O-acetylginsenoside Rg₃ (24), and 3β,6α,12β-trihydroxydammar-20(21),24-diene (25), were isolated from notoginseng for the first time.

KEYWORDS: Steamed notoginseng; *Panax notoginseng*; dammarane; notoginsenosides ST-1–ST-3 and ST-5

INTRODUCTION

Notoginseng, the root of *Panax notoginseng* (Burk.) F. H. Chen (Araliaceae), a traditional Chinese medicinal herb, is mainly cultivated in Yunnan province of China. Traditionally, notoginseng has been used in two forms, raw and processed. The former is mainly used for injuries from falls and removing blood stasis, whereas the latter is used as a tonic to promote blood circulation (1). It is well-known that dammarane-type triterpenoidal saponins are the main bioactive constituents of notoginseng, from which more than 30 saponins have been isolated (2). In the 1980s, we compared the saponin compositions in processed notoginseng to those in the raw one and found that the contents of bisdesmosides (such as ginsenosides Rg₁, Rb₁, Rd, Re, and notoginsenoside R1) in the processed form were lower, whereas the monodesmosides (such as ginsenosides Rg₂, Rg₃, and notoginsenoside R2) increased (3). In recent years, Lau et al. (4, 5) reported that the concentrations of less polar saponins, for example, ginsenosides 20(S)-Rh₁, 20(R)-Rh₁, Rk₃, Rh₄, 20(S)-Rg₃, 20(R)-Rg₃, Rk₁, and Rg₅, increased in steamed notoginseng, by HPLC analyses. However, no systematic phytochemical studies on the steamed notoginseng were reported. As a part of our continuing studies on the chemical constituents of *Panax* medicinal plants, the present study was undertaken to isolate and structurally elucidate new dammarane

glycosides, together with known dammarane-type triterpenoids and other types of known compounds from the steamed notoginseng.

MATERIALS AND METHODS

Material. Air-dried notoginseng was obtained from Wenshan County, Yunnan Province, China, in May 2006. The raw materials were crushed into small grains and then steamed at 120 °C for 12 h directly, without mixing with water, to give the steamed notoginseng, which was used for extraction and isolation in this study.

Apparatus. Optical rotations were performed on a P-1020 polarimeter (JASCO, Tokyo, Japan). IR spectra were measured on a Bruker Tensor 27 spectrometer with KBr pellets. ¹H and ¹³C NMR, ¹H–¹H COSY, HMQC, HMBC, ROSEY, and TOCSY spectra were recorded in C₅D₅N with Bruker AM-400 and DRX-500 spectrometers. Coupling constants are expressed in hertz, and chemical shifts are given on a δ (parts per million) scale with TMS as an internal standard. FAB-MS were recorded on an AutoSpec 3000 spectrometer (VG, Manchester, U.K.) with glycerol as the matrix, in *m/z*. HRESI-MS were recorded on an API Qstar Pulsa LC/TOF spectrometer. Melting points (mp) were determined on an XRC-1 melting point apparatus, produced by Sichuan University, China (uncorrected).

Column chromatography (CC) was performed with Diaion 101 resin (Shandong Lukang Pharmaceutical Co., Ltd., China), silica gel (200–300 mesh) (Qingdao Marine Chemical and Industrial Factory, China), MCI-gel CHP20P (75–100 μm) (Mitsubishi Chemical Co., Ltd., Japan), and Rp-8 or Rp-18 gel (40–60 μm) (Merck, Darmstadt, Germany). TLC was performed with silica gel H pre-coated plates (Qingdao Marine Chemical and Industrial Factory, China) with CHCl₃/MeOH/H₂O (75:25:3; 80:20:2; 85:15:1; or 90:10:1, v/v) and CHCl₃/MeOH (90:10 to

* Author to whom correspondence should be addressed (telephone +86-871-5223235; fax +86-871-5150124; e-mail zhangyj@mail.kib.ac.cn).

[†] Kunming Institute of Botany.

[‡] Graduate School of the Chinese Academy of Sciences.




Figure 1. New compounds 1–4 isolated from the steamed notoginseng.

25:1, v/v) and Rp-8 or Rp-18 precoated plates (Merck) with MeOH/H₂O (60:40 to 100:0, v/v). Spots were detected by spraying with 10% H₂SO₄ in EtOH followed by heating.

Extraction and Isolation. The air-dried steamed notoginseng (4.7 kg) was powdered and extracted with ethanol four times. The ethanolic extract was concentrated under reduced pressure to afford a residue (778 g). The residue was dissolved in H₂O and passed through a Diaion 101 column (90 × 8.5 cm) two times, eluting with H₂O and MeOH, successively. The methanol eluates were combined and concentrated under reduced pressure to give a residue (440 g), which was further subjected to silica gel column chromatography eluted with (CHCl₃: MeOH:H₂O, 85:15:1) to afford nine fractions (1–9). Fraction 1 (13 g) was repeatedly chromatographed over silica gel (CHCl₃/MeOH/H₂O), Rp-8, Rp-18, and MCI-gel CHP20P columns (MeOH/H₂O) to afford **5** (252 mg) and **6** (89 mg). In a similar way, compounds **25** (30 mg), **26** (23 mg), **27** (32 mg), **28** (666 mg), **29** (157 mg), and **30** (191 mg) from fraction 3 (21 g), compounds **1** (14 mg), **2** (51 mg), **3** (41 mg), **7** (25 mg), **8** (6 mg), **17** (33 mg), **18** (83 mg), **19** (10 mg), **20** (15 mg), **21** (20 mg), **22** (48 mg) and **24** (16 mg) from fraction 5 (60 g), compounds **9** (151 mg), **10** (23 mg), **11** (71 mg), **16** (20 mg), and **23** (238 mg) from fraction 7 (38 g), and compounds **4** (21 mg), **12** (114 mg), **13** (720 mg), **14** (1076 mg), and **15** (36 mg) from fraction 9 (40 g) were obtained, respectively, through repeated column chromatography (Figures 1 and 2).

Notoginsenoside ST-1 [$3\beta,6\alpha,12\beta,24,25$ -pentahydroxydammar-20(22)(*E*)-ene-6-*O*- β -D-glucopyranoside] (**1**) was obtained as a white amorphous powder: mp 168–171 °C; $[\alpha]_D^{26} +29.1^\circ$ (c 0.249, MeOH); IR (KBr), ν_{\max} (cm⁻¹) 3424, 2961, 2931, 2876, 1637, 1462, 1384, 1156, 1075, 1031, 928, 890; FAB-MS (negative ion mode), m/z 653 [M - H]⁻, 491 [M - 162(glucosyl) - H]⁻; HRESIMS (negative ion mode), m/z 653.4270 [M - H]⁻, (calcd for 653.4264, C₃₆H₆₁O₁₀); ¹H and ¹³C NMR data, see Tables 1–3.

Notoginsenoside ST-2 [$3\beta,12\beta,24,25$ -tetrahydroxy-23-methoxydammar-20(22)(*E*)-ene-3-*O*- β -D-glucopyranosyl-(1 \rightarrow 2)- β -D-glucopyranoside] (**2**) was obtained as a white amorphous powder: mp 172–177 °C; $[\alpha]_D^{26} -9.4^\circ$ (c 0.264, MeOH); IR (KBr), ν_{\max} (cm⁻¹) 3417, 2944, 2877, 1640, 1549, 1465, 1452, 1388, 1370, 1077, 1033, 978; FAB-MS (negative ion mode), m/z 831 [M + 1]⁻; HRESIMS (negative ion mode), m/z 829.4942 [M - H]⁻ (calcd for 829.4949, C₄₃H₇₃O₁₅); ¹H and ¹³C NMR data, see Tables 1–3.

Notoginsenoside ST-3 [$3\beta,12\beta,24,25$ -tetrahydroxy-23-methoxydammar-20(22)(*Z*)-ene-3-*O*- β -D-glucopyranosyl-(1 \rightarrow 2)- β -D-glucopyranoside] (**3**) was obtained as a white amorphous powder: mp 171–175 °C; $[\alpha]_D^{26} +20.3^\circ$ (c 0.254, MeOH); IR (KBr), ν_{\max} (cm⁻¹) 3418, 2944, 2877, 1640, 1549, 1465, 1453, 1389, 1372, 1161, 1078, 1038, 978; FAB-MS (negative ion mode), m/z 830 [M]⁻, 668 [M - 162(glucosyl)]⁻; HRESIMS (negative ion mode), m/z 829.4955 [M - H]⁻ (calcd for 829.4949, C₄₃H₇₃O₁₅); ¹H and ¹³C NMR data, see Tables 1–3.

Notoginsenoside ST-5 [$3\beta,12\beta,20(S),25$ -tetrahydroxydammar-23-ene-3-*O*- β -D-xylopyranosyl-(1 \rightarrow 2)- β -D-glucopyranosyl-(1 \rightarrow 2)- β -D-glucopyranoside] (**4**) was obtained as a white amorphous powder: mp 193–196 °C; $[\alpha]_D^{26} +2.3^\circ$ (c 0.257, MeOH); IR (KBr), ν_{\max} (cm⁻¹) 3422, 2940, 2878, 1639, 1461, 1375, 1077, 1043, 896; FAB-MS (negative ion mode), m/z 932 [M]⁻, 800 [M - 132(xylosyl)]⁻, 476 [M - 132(xylosyl) - 162(glucosyl) - 162(glucosyl)]⁻; HRESIMS (negative ion mode), m/z 931.5241 [M - H]⁻ (calcd for 931.5266, C₄₇H₇₉O₁₈); ¹H and ¹³C NMR data, see Tables 1–3.

Acid Hydrolysis of Compounds 1–4. Compounds 1–3 (each 5 mg) were hydrolyzed with 2 M HCl/dioxane (1:1, 4 mL) under reflux for 8 h, respectively. The reaction mixture was partitioned between H₂O and CHCl₃ (2 mL × 3). The aqueous layer was neutralized with 2 M NaOH and then dried to give a monosaccharide mixture. A solution of the sugar mixture in pyridine (2 mL) was added to L-cysteine methyl ester hydrochloride (about 1.5 mg) and kept at 60 °C for 1 h. Then trimethylsilylimidazole (about 1.5 mL) was added to the reaction mixture and kept at 60 °C for 30 min. The mixture was subjected to GC analysis, run on a Shimadzu GC-14C gas chromatograph equipped with an H₂ flame ionization detector. The column was a 30 m × 0.32 mm i.d. 30QC2/AC-5 quartz capillary column with the following conditions: column temperature, 180–280 °C; programmed increase, 3 °C/min; carrier gas, N₂ (1 mL/min); injector and detector temperature, 250 °C; injection volume, 4 μ L; and split ratio, 1/50. The configuration of the sugar moiety was determined by comparing the retention time with the derivatives of the authentic samples. The retention times of D-/L-glucose were 21.115/21.565 min. The configuration of the sugar moiety from compounds 1–3 was D-glucose.

Compound **4** (8 mg) was treated as described for compounds 1–3 to give the monosaccharide mixture, which was detected by GC analysis running on an Agilent Technologies HP5890 gas chromatograph. The retention times of D-/L-glucose and D-/L-xylose were 19.715/20.159 and 14.606/15.256 min, respectively. The configuration of the sugar moiety from compound **4** was D-glucose and D-xylose.

RESULTS AND DISCUSSION

The ethanolic extracts of the steamed notoginseng were chromatographed repeatedly over Diaion 101 resin, silica gel, Rp-8, Rp-18, and MCI-gel CHP20P column to yield 4 new dammarane-type glycosides, notoginsenosides ST-1–ST-3 (**1–3**) and ST-5 (**4**), and 26 known compounds (**5–30**).

The known compounds were identified as ginsenosides Rh₄ (**5**) (**6**), Rk₃ (**6**) (**6**), Rg₅ (**7**) (**6**), Rk₁ (**8**) (**6**), Rg₁ (**9**) (**7**), Rg₂ (**10**) (**7**), Rf (**11**) (**8**), Re (**12**) (**7**), Rb₁ (**13**) (**9**), and Rd (**14**) (**9**), notoginsenosides R1 (**15**) and R2 (**16**) (**8**), sanchinoside-B1 (**17**) (**10**), 25-hydroxy-20(*R*)-ginsenoside Rh₁ (**18**) (**7**), koryoginsenoside-R₁ (**19**) (**11**), yesanchinoside D (**20**) (**12**), 20(*S*)-ginsenoside Rh₁ (**21**) (**7**), 20(*R*)-ginsenoside Rh₁ (**22**) (**7**), 20(*R/S*)-ginsenoside Rg₃ (**23**) (**13**), 6''-*O*-acetyl-ginsenoside Rg₃ (**24**) (**14**, **15**), 3 $\beta,6\alpha,12\beta$ -trihydroxydammar-20(21),24-diene (**25**) (**16**, **17**), 20(*S*)-protopanaxatriol (**26**) (**18**), 20(*R*)-protopanaxatriol (**27**) (**18**), panaxytriol (**28**) (**19**, **20**), (*Z,Z*)-9,12-octadecadienoic acid 2-hydroxy-1,3-propanediny ester (**29**) (**21**), and 5-hydroxymethyl-2-furancarboxaldehyde (**30**) (**22**), respectively, on the basis of FAB-MS, NMR spectra data, and comparison with those reported in the literature. Among them, glycosides **19**, **20**, and **24**, as well as the aglycone **25**, were isolated from *P. notoginseng* for the first time.

Compound **1** was obtained as a white amorphous powder. Its molecular formula C₃₆H₆₂O₁₀ was determined by negative high-resolution electrospray ionization (HRESI) mass spectrometry and confirmed by DEPT spectra. The IR spectrum showing strong absorbances at 3424, 1637, 1384, 1075, 1031, 928, and 890 cm⁻¹ suggested the presence of hydroxyl groups and a double bond in **1**. The fragment ion peaks at 653 [M - H]⁻ and 491 [M - H - 162(hexosyl)]⁻ in fast atom bombardment (FAB)-MS, and the anomeric proton and




Figure 2. Known compounds 5–30 from the steamed notoginseng.

carbon signals at δ_H 5.03 (d, $J = 8.0$ Hz) and δ_C 106.1, respectively, in 1H and ^{13}C NMR suggested the presence of a β -hexosyl unit in **1**. Acid hydrolysis of **1** gave D-glucose as the sugar component, indicating **1** was a β -D-glycoside. The NMR data of **1** were similar to those of sanchinoside-B₁ (**17**), except that **1** had one more hydroxyl group located on the side chain of the aglycone. The structure of **1** was further elucidated by 2D NMR experiments. In the proton–proton homonuclear shift correlated spectroscopy (1H – 1H COSY) spectrum, the methylene signal at δ_H 2.58 (m, H-23) was found to be coupled with both an olefinic proton at δ_H 5.74 (t, $J = 7.0$ Hz, H-22) and an oxygenated methine at δ_H 3.80 (m, H-24). The heteronuclear multiple-bond correlation (HMBC) spectrum exhibited the following long-range correlation signals: δ_H 2.82 (H-17) with δ_C 141.8 (C-20), 13.1 (C-21) and 123.9 (C-22); δ_H 1.83 (H-21) with δ_C 51.3 (C-17), C-20 and C-22; δ_H 5.74 (H-22) with δ_C 31.2 (C-23), C-17, and C-21; δ_H 2.58 (H-23) with δ_C 79.0 (C-24), C-20, and C-22; δ_H 3.80 (H-24) with δ_C 72.3 (C-25) and C-22; and δ_H 1.51 (H-26) and 1.49 (H-27) with C-24 and C-25, respectively. These data confirmed that the additional hy-

droxyl group was located at C-24. In addition, the HMBC correlation between δ_H 5.03 (H-1') and δ_C 80.1 (C-6) revealed that the β -D-glucosyl unit was located at C-6 of the aglycone.

In the rotating frame Overhauser effect spectroscopy (ROSEY) spectrum of **1**, the methyl proton at δ_H 1.83 (s, H-21) was correlated with δ_H 2.58 (H-23), but not with the olefinic proton at δ_H 5.74 (H-22). This observation indicated that the C-20(22) double bond was of the *E* form. The chemical shift of the C-21 methyl group (δ_C 13.1) was similar to those of ginsenoside Rh₄ (**23**), sanchinoside-B₁ (**10**), and notoginsenoside T₁ (**24**). Thus, the structure of notoginsenoside ST-1 (**1**) was determined as 3 β ,6 α ,12 β ,24,25-pentahydroxydammar-20(22)*E*-ene-6-*O*- β -D-glucopyranoside.

The molecular formula of compound **2** was established as C₄₃H₇₄O₁₅ by the quasimolecular ion peak in the negative HRESIMS. Its NMR features were similar to that of ginsenoside Rg₅ (**7**) (**6**), except for the side chain of the aglycone. Compared with compound **7**, compound **2** had one fewer double bond, but one more methoxyl and two more hydroxyl groups (δ_H 3.32, δ_C 55.1, 72.3, and 80.7) on the side chain. In the 1H – 1H COSY spectrum of **2**, the oxygen-bearing methine proton at δ_H 4.31

Table 1. ^{13}C NMR Spectroscopic Data of Compounds **1–4** (δ , in $\text{C}_5\text{D}_5\text{N}$)

no.	1	2	3	4
1	39.5	39.3	39.3	39.2
2	28.0	26.8	26.8	26.8
3	78.6	88.9	89.0	89.0
4	40.4	39.8	39.8	39.8
5	61.5	56.4	56.4	56.5
6	80.1	18.5	18.5	18.5
7	45.4	35.4	35.4	35.2
8	41.4	40.3	40.3	40.2
9	50.5	50.7	51.0	50.5
10	39.8	37.1	37.0	37.0
11	32.7	32.3	33.2	32.2
12	72.5	71.9	72.2	71.2
13	50.3	51.0	52.0	49.0
14	50.7	50.9	51.1	51.8
15	32.5	32.7	32.7	31.3
16	27.7	28.6	30.0	26.8
17	51.3	51.4	50.1	54.2
18	16.9	16.5	16.5	16.5
19	17.4	15.9	15.9	15.9
20	141.8	143.5	145.3	73.4
21	13.1	13.3	15.5	27.7
22	123.9	123.5	123.2	40.1
23	31.2	79.5	78.2	123.9
24	79.0	80.7	80.7	142.0
25	72.3	72.3	72.5	69.9
26	26.6	29.4	28.5	30.8
27	26.0	24.8	26.1	30.8
28	31.8	28.2	28.2	28.2
29	16.4	16.7	16.6	16.7
30	17.7	17.0	17.1	17.1
OCH ₃		55.1	55.4	
GlcI-1'	106.1	105.2	105.2	104.8
GlcI-2'	75.5	83.4	83.4	83.0
GlcI-3'	79.7	78.0	78.0	78.7
GlcI-4'	71.9	71.6	71.6	71.1
GlcI-5'	78.2	78.4	78.4	78.3
GlcI-6'	63.1	62.7	62.9	63.0
GlcII-1''		106.1	106.1	103.2
GlcII-2''		77.2	77.2	84.6
GlcII-3''		78.2	78.2	78.0
GlcII-4''		71.6	71.6	71.9
GlcII-5''		78.4	78.4	77.8
GlcII-6''		62.8	62.7	63.0
Xyl-1'''				106.5
Xyl-2'''				76.0
Xyl-3'''				77.8
Xyl-4'''				70.8
Xyl-5'''				67.5

(H-23) was coupled with both olefinic proton at δ_{H} 5.68 (H-22) and another oxygen-bearing methine proton at δ_{H} 3.85 (H-24). Moreover, the HMBC correlations of H-17 (δ_{H} 2.83) with C-20 (δ_{C} 143.5), C-21 (δ_{C} 13.3), and C-22 (δ_{C} 123.5); H-21 (δ_{H} 1.91) with C-17 (δ_{C} 51.4), C-20 and C-22; H-22 with C-17, C-21, and C-24 (δ_{C} 80.7); H-23 with C-20 and C-24; H-24 (δ_{H} 3.85) with C-22, C-23 (δ_{C} 79.5), C-25 (δ_{C} 72.3), C-26 (δ_{C} 29.4), and C-27 (δ_{C} 24.8); H-26 (δ_{H} 1.60) and H-27 (δ_{H} 1.45) with C-24 and C-25, and the methoxyl protons (δ_{H} 3.32) with C-23 revealed the structure of the side chain in **2**.

The *E* form of the C-20(22) double bond was determined by the ROSEY spectrum, in which correlation of the methyl proton at δ_{H} 1.91 (H-21) with the methoxyl proton at δ_{H} 3.32 was observed; however, there was no correlation between H-21 and the olefinic proton at δ_{H} 5.68 (H-22). The sugar sequence and location were confirmed by the HMBC correlations of one anomeric proton at δ_{H} 4.90 (d, $J = 7.5$ Hz, H-1') with C-3 (δ_{C} 88.9) of the aglycone and another anomeric proton at δ_{H} 5.35 (d, $J = 7.7$ Hz, H-1'') with C-2' (δ_{C} 83.4) of the inner glucosyl unit. On the basis of the above evidence, compound **2** was

Table 2. ^1H NMR Spectroscopic Data for the Aglycone Moieties of Compounds **1–4** [δ (J, Hertz), in $\text{C}_5\text{D}_5\text{N}$]

no.	1	2	3	4
1	1.70 m	1.50 m	1.45 m	1.48 m
	1.03 m	0.74 m	0.72 m	0.78 m
2	1.93 m	2.18 m	2.17 m	2.19 m
	1.85 m	1.81 m	1.80 m	1.86 m
3	3.53 dd	3.26 m	3.28 dd	3.30 m
	(4.5, 11.5)		(4, 11.5)	
5	1.44 m	0.66 m	0.66 m	0.69 m
6	4.43 m	1.49 m	1.47 m	1.49 m
		1.36 m	1.34 m	1.38 m
7	2.53 m	1.46 m	1.45 m	1.48 m
	1.97 m	1.22 m	1.21 m	1.25 m
9	1.57 m	1.36 m	1.35 m	1.40 m
11	2.04 m	1.91 m	1.91 m	2.03 m
	1.46 m	1.41 m	1.36 m	1.49 m
12	3.94 m	3.83 m	3.87 m	3.90 m
13	1.99 m	1.97 m	2.09 m	2.06 m
15	1.72 m	1.67 m	1.69 m	1.60 m
	1.16 m	1.10 m	1.08 m	1.04 m
16	1.86 m	1.94 m	2.01 m	1.91 m
	1.42 m	1.57 m	1.52 m	1.47 m
17	2.82 m	2.83 m	2.77 m	2.36 m
18	0.82 s	0.79 s	0.75 s	0.80 s
19	1.22 s	1.01 s	0.97 s	1.02 s
21	1.83 s	1.91 s	2.08 s	1.43 s
22	5.74 t (7.0)	5.68 d (9.8)	5.80 d (9.7)	2.78 m
				2.49 m
23	2.58 m	4.31 m	4.48 m	6.31 m
24	3.80 m	3.85 d (6.5)	3.77 d (4.9)	6.02 d (15.5)
26	1.51 s	1.60 s	1.56 s	1.55 s
27	1.49 s	1.45s	1.52 s	1.54 s
28	2.07 s	1.26 s	1.27 s	1.28 s
29	1.61 s	1.08 s	1.08 s	1.11 s
30	1.03 s	0.91 s	0.93 s	0.94 s
OCH ₃		3.32 s	3.41 s	

Table 3. ^1H NMR Spectroscopic Data for the Sugar Moieties of Compounds **1–4** [δ (J, Hertz), in $\text{C}_5\text{D}_5\text{N}$]

no.	1	2	3	4
GlcI-1'	5.03d (8.0)	4.90 d (7.5)	4.92 d (7.5)	4.93 d (7.5)
GlcI-2'	4.10 t (8.5)	4.20 m	4.23 m	4.14 m
GlcI-3'	4.27 m	4.21 m	4.24 m	4.35 m
GlcI-4'	4.22 m	4.11 m	4.13 m	4.09 m
GlcI-5'	3.96 m	3.90 m	3.92 m	3.95 m
GlcI-6'	4.53 dd (2.5, 11.5)	4.53 m	4.56 m	4.57 br.d (10)
	4.37 dd (5.5, 12.0)	4.32 m	4.35 m	4.34 m
GlcII-1''		5.35 d (7.7)	5.37 d (7.6)	5.53 d (7.5)
GlcII-2''		4.08 m	4.11 m	4.18 m
GlcII-3''		4.28 m	4.31 m	4.28 m
GlcII-4''		4.30 m	4.31 m	4.18 m
GlcII-5''		3.90 m	3.92 m	3.87 m
GlcII-6''		4.46 m,	4.48 m,	4.50 m br.d
		4.44 m	4.45 m	
Xyl-1'''				5.39 d (6.5)
Xyl-2'''				4.09 m
Xyl-3'''				4.32 m
Xyl-4'''				4.14 m
Xyl-5'''				4.34 m
				3.68 m

elucidated to be $3\beta,12\beta,24,25$ -tetrahydroxy-23-methoxydammar-20(22)(*E*)-ene-3-*O*- β -D-glucopyranosyl-(1 \rightarrow 2)- β -D-glucopyranoside and named notoginsenoside ST-2.

Compound **3** showed the same molecular formula, $\text{C}_{43}\text{H}_{74}\text{O}_{15}$, as **2**. The NMR data of **3** were identical to those of **2**, except for the upfield shifts of C-17 and C-23, both by -1.3 ppm, and the downfield shifts of C-20, C-21, and C-27 by $+1.8$, $+2.2$, and $+1.3$ ppm, respectively. This suggested that **3** was a geometric isomer of **2** at the C-20(22) double

bond. The configuration of the C-20(22) double bond in **3** was further confirmed to be of the *Z* form due to the NOESY correlation between the methyl proton at δ_{H} 2.08 (H-21) and the olefinic proton at δ_{H} 5.80 (H-22), as seen in the ROESY spectrum. Thus, compound **3** was determined to be 3 β ,12 β ,24,25-tetrahydroxy-23-methoxydammar-20(22)(*Z*)-ene-3-*O*- β -D-glucopyranosyl-(1 \rightarrow 2)- β -D-glucopyranoside and named notoginsenoside ST-3.

Compound **4** had a molecular formula $\text{C}_{47}\text{H}_{80}\text{O}_{18}$ on the basis of the quasimolecular ion peak in the negative HRESIMS. Comparisons of the ^1H and ^{13}C NMR data with those of notoginsenoside Ft₁ (**25**) indicated that the only difference between them was the side chain. Both compounds presented the same degrees of unsaturation as eight; however, **4** had one more oxygen atom. Besides two quaternary oxygen-bearing carbon signals at δ_{C} 73.4 (C-20) and 69.9 (C-25), there were two olefinic methines [δ_{C} 123.9 (C-23) and 142.0 (C-24); δ_{H} 6.31 (H-23) and 6.02 (H-24)] in the side chain of **4**. The structure of the side chain was further established by an HMBC experiment, in which the long-range cross peaks from H-21 (δ_{H} 1.43) to C-17 (δ_{C} 54.2) and C-20 (δ_{C} 73.4); from H-22 (δ_{H} 2.78 and 2.49) to C-20, C-21 (δ_{C} 27.7), C-23, and C-24; from H-23 to C-22 (δ_{C} 40.1) and C-25 (69.9); and from H-24 to C-22, C-25, C-26 (δ_{C} 30.8), and C-27 (δ_{C} 30.8) were observed. It indicated that the double bond was located between C-23 and C-24, and two hydroxyl groups were at C-20 and C-25, respectively. The fragment ion peaks at m/z 932 [M^-] and 800 [$\text{M} - 132$] in the FAB/MS, along with the HMBC correlations of δ_{H} 4.93 (d, $J = 7.5$ Hz, H-1') with δ_{C} 89.0 (C-3); δ_{H} 5.53 (d, $J = 7.5$ Hz, H-1'') with δ_{C} 83.0 (C-2'); and δ_{H} 5.39 (d, $J = 6.5$ Hz, H-1''') with δ_{C} 84.6 (C-2''), confirmed the sugar moiety of **4** to be the same as that of notoginsenoside Ft₁. Hence, compound **4** was determined to be 3 β ,12 β ,20(*S*),25-tetrahydroxydammar-23-ene-3-*O*- β -D-xylopyranosyl-(1 \rightarrow 2)- β -D-glucopyranosyl-(1 \rightarrow 2)- β -D-glucopyranoside and named notoginsenoside ST-5.

On the basis of traditional Chinese medical theories, the herbs need processing after harvest for different purposes. The chemical compositions of the steamed notoginseng were significantly different from those in the raw materials. Because the process of steaming notoginseng was carried out under high temperature, the new monodesmosides should be formed by the degradation of the C-20 glycosyl moiety of the dammarane skeleton during the processing. It is apparent that the rich dammarane-type monodesmosides presented are not only chemically characteristic of steamed notoginseng but also give special biological activities to this processed herb. The pharmaceutical evaluation of the steamed notoginseng is now in progress.

ACKNOWLEDGMENT

We are grateful to the members of the Analytical Group of State Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, for measurements of all spectra.

LITERATURE CITED

- (1) Ye, D. J.; Zhang, S. C. Herbs from roots and rhizomes. *Chinese Medicinal Herbs Preparation*; People's Medical Publishing House: Peking, China, 1999; p 187.
- (2) Wang, C. Z.; McEntee, E.; Wicks, S.; Wu, J. A.; Yuan, C. S. Phytochemical and analytical studies of *Panax notoginseng* (Burk.) F. H. Chen. *J. Nat. Med.* **2006**, *60*, 97–106.
- (3) Yang, C. R.; Cui, Z. H.; Wu, M. Z.; Zhou, J. Isolation and structural elucidation of the glycoside saponins from the raw and steamed notoginseng. *Zhong Yao Tong Bao* **1985**, *10*, 407–418.
- (4) Lau, A. J.; Seo, B. H.; Woo, S. O.; Koh, H. L. High-performance liquid chromatographic method with quantitative comparisons of whole chromatograms of raw and steamed *Panax notoginseng*. *J. Chromatogr., A* **2004**, *1057*, 141–149.
- (5) Lau, A. J.; Woo, S. O.; Koh, H. L. Analysis of saponins in raw and steamed *Panax notoginseng* using high-performance liquid chromatography with diode array detection. *J. Chromatogr., A* **2003**, *1011*, 77–87.
- (6) Park, I. H.; Kim, N. Y.; Han, S. B.; Kim, J. M.; Kwon, S. W.; Kim, H. J.; Park, M. K.; Park, J. H. Three new dammarane glycosides from heat processed ginseng. *Arch. Pharm. Res.* **2002**, *25*, 428–432.
- (7) Teng, R. W.; Li, H. Z.; Chen, J. T.; Wang, D. Z.; He, Y. N.; Yang, C. R. Complete assignment of ^1H and ^{13}C -NMR data for nine protopanaxtriol glycosides. *Magn. Reson. Chem.* **2002**, *40*, 483–488.
- (8) Zhou, J.; Wu, M. Z.; Taniyasu, S.; Besso, H.; Tanaka, O.; Saruwatari, Y.; Fuwa, T. Dammarane-saponins of Sanchi-Ginseng, roots of *Panax notoginseng* (Burk.) F. H. Chen (Araliaceae): structures of new saponins, notoginsenosides-R1 and R2, and identification of ginsenosides-Rg₂ and Rh₁. *Chem. Pharm. Bull.* **1983**, *29*, 2844–2850.
- (9) Matsuura, H.; Kasai, R.; Tanaka, O.; Saruwatari, Y. I.; Fuwa, T.; Zhou, J. Further studies on dammarane-saponins of Sanchi-ginseng. *Chem. Pharm. Bull.* **1983**, *31*, 2281–2287.
- (10) Wei, J. X.; Wang, L. A.; Du, H.; Li, R. Isolation and identification of sanchinoside B1 and B2 from rootlets of *Panax notoginseng* (Burk.) F. H. Chen. *Acta Pharm. Sinica* **1985**, *20*, 288–293.
- (11) Kim, D. S.; Chang, Y. J.; Zedk, U.; Zhao, P.; Liu, Y. Q.; Yang, C. R. Dammarane saponins from *Panax ginseng*. *Phytochemistry* **1995**, *40*, 1493–1497.
- (12) Zou, K.; Zhu, S.; Tohda, C.; Cai, S. Q.; Komatsu, K. Dammarane-type triterpene saponins from *Panax japonicus*. *J. Nat. Prod.* **2002**, *65*, 346–351.
- (13) Yang, T. R.; Kasai, R.; Zhou, J.; Tanaka, O. Dammarane saponins of leaves and seeds of *Panax notoginseng*. *Phytochemistry* **1983**, *22*, 1473–1478.
- (14) Teng, R. W.; Ang, C. S.; McManus, D.; Armstrong, D.; Mau, S.; Bacic, A. Regioselective acylation of ginsenosides by Novozyme 435 to generate molecular diversity. *Helv. Chim. Acta* **2004**, *87*, 1860–1872.
- (15) Gebhardt, S.; Bihler, S.; Manfred, S. Z.; Riva, S.; Monti, D.; Falcone, L.; Danieli, B. Biocatalytic generation of molecular diversity: modification of ginsenoside Rb₁ by β -1,4-galactosyl-transferase and *Candida antarctica* lipase part 4. *Helv. Chim. Acta* **2002**, *85*, 1943–1959.
- (16) Zhao, Y. Q.; Wu, L. J.; Yuan, C. L.; Li, X.; Chen, Y. J. A new triterpene saponin from the stems and leaves of *Panax ginseng*. *Chin. Chem. Lett.* **1992**, *3*, 887–888.
- (17) Ryu, J. H.; Park, J. H.; Eun, J. H.; Jung, J. H.; Sohn, D. H. A dammarane glycoside from Korean red ginseng. *Phytochemistry* **1997**, *44*, 931–933.
- (18) Yu, M.; Zhao, Y. Q. Identification and structure elucidation of a pair of configurational isomers from the fruits of *Panax ginseng*. *Chin. Trad. Herbal Drugs* **2002**, *33*, 404–405.
- (19) Zhao, P.; Liu, Y. Q.; Yang, C. R. Minor constituents from the roots of *Panax notoginseng*. *Acta Bot. Yunnanica* **1993**, *15*, 409–412.
- (20) Hirakura, K.; Morita, M.; Nakajima, K.; Ikeya, Y.; Mitsushashi, H. Polyacetylenes from the roots of *Panax ginseng*. *Phytochemistry* **1991**, *30*, 3327–3333.
- (21) Yang, H.; Xie, J. L.; Sun, H. D. Study on chemical constituents of *Saussurea lappa* II. *Acta Bot. Yunnanica* **1997**, *1*, 92–96.
- (22) Hearn, M. T. W. Carbon-13 chemical shifts in some substituted furans and thiophenes. *Aust. J. Chem.* **1976**, *29*, 107–113.

- (23) Beak, N.; Kim, D.; Lee, Y.; Park, J.; Lee, C.; Kim, S. Ginsenoside Rh4, a genuine dammarane glycoside from Korean red ginseng. *Planta Med.* **1996**, 62, 86–87.
- (24) Teng, R. W.; Li, H. Z.; Wang, D. Z.; Yang, C. R. Hydrolytic reaction of plant extracts to generate molecular diversity: new dammarane glycoside from the mild acid hydrolysate of root saponins of *Panax notoginseng*. *Helv. Chim. Acta* **2004**, 87, 1270–1278.
- (25) Chen, J. T.; Li, H. Z.; Wang, D.; Zhang, Y. J.; Yang, C. R. New dammarane monodesmosides from the acidic deglycosylation of notoginseng-leaf saponins. *Helv. Chim. Acta* **2006**, 89, 1442–1448.

Received for review October 11, 2007. Revised manuscript received December 26, 2007. Accepted December 28, 2007. This work was partly supported by the NSFC (30472156 and U0632010).

JF073000S