

Communications to the Editor

[Chem. Pharm. Bull.]
34(9)3974—3977(1986)

**TUBEIMOSIDE I, A NEW CYCLIC BISDES MOSIDE FROM CHINESE
CUCURBITACEOUS FOLK MEDICINE "TU BEI MU",
A TUBER OF *Bolbostemma paniculatum***

Ryoji Kasai,^a Masazumi Miyakoshi,^a Kazuhiro Matsumoto,^a Rui-Lin Nie,^b
Jun Zhou,^b Toshinobu Morita,^a and Osamu Tanaka^{*,a}
Institute of Pharmaceutical Sciences, Hiroshima University School
of Medicine,^a Kasumi, Minami-ku, Hiroshima 734, Japan
and Kunming Institute of Botany, Academia
Sinica,^b Kunming Yunnan, China

An oleanane type triterpenoid saponin, named tubeimoside I was isolated from tubers of *Bolbostemma paniculatum* (Maxim.) Franquet (Cucurbitaceae), a Chinese folk medicine, "Tu Bei Mu". Its novel cyclic structure, including the chirality of the 3-hydroxy-3-methylglutarate moiety, is reported. The name cyclic bisdesmoside is proposed for this type of saponin. The potent solubilizing effect of this compound is also described.

KEYWORDS—— tubeimoside I; cyclic bisdesmoside; bayogenin; saponin; *Bolbostemma paniculatum*; Cucurbitaceae; Chinese folk medicine; 3-hydroxy-3-methylglutarate; mevalonolactone chirality; solubilizing effect

A Chinese crude drug "Tu Bei Mu", steamed and dried tubers of *Bolbostemma paniculatum* (Maxim.) Franquet (Cucurbitaceae), has been used in China as an anti-inflammatory agent for mastitis and an antidote for snake poison.

A suspension of a methanolic extract of the tubers in water was passed through a column of highly porous polymer resin to give a mixture of glycosides which was chromatographed on silica gel and then on reversed phase silica gel. This afforded a new glycoside named tubeimoside I (1), a white powder, $[\alpha]_D^{17} +14.6^\circ$ ($c=1.09$, MeOH), $C_{63}H_{98}O_{29} \cdot 2H_2O$, yield 1.93%.

The 1H - and ^{13}C -NMR of 1 showed the presence of five monosaccharide units. On acid hydrolysis, 1 afforded an aglycone (2), D-glucose, L-arabinose, L-rhamnose and D-xylose. The identification of these monosaccharides, including their absolute configuration, was established according to the method recently reported by Oshima *et al.*¹⁾ The aglycone 2 was obtained as colorless needles (from MeOH-H₂O), mp 333-335°C, $[\alpha]_D^{17} +98.8^\circ$ ($c=0.43$, pyridine). On inspection of the ^{13}C -NMR spectral data, 2 was proved to be identical with bayogenin, the oleanane type triterpene already isolated from *Castanospermum australe* Cunn. et Fras. as a minor sapogenin.²⁾

The IR spectrum (in Nujol) of 1 revealed the presence of ester groups ($1730cm^{-1}$) and the ^{13}C -NMR spectrum (in pyridine-*d*₅) of 1 showed three ester carbonyl signals at δ 171.2, 171.4 and 176.0, the last of which can be assigned as the 28-O-glycosyl ester.³⁾ Mild alkaline treatment of 1 with 0.5% KOH-H₂O yielded 3 and a dicarboxylic


acid (4). The latter was identified as 3-hydroxy-3-methylglutaric acid by comparison of spectral data and GC-MS with an authentic sample.

In comparing the ^{13}C -NMR spectra (in pyridine- d_5) of 3 and 2, glycosylation shifts⁴⁾ were observed for the signals due to C-2 (-1.1ppm), -3 (+9.5ppm), -23 (-3.0ppm)⁴⁾ and -28 (-4.0ppm)³⁾. This indicates that 3 is a bisdesmoside of 2 with glycosyl linkage at both 3-OH and 28-COOH. The EI-MS spectrum of a trimethylsilyl (TMSi) ether of 3 exhibited fragment ions due to sugar moieties at m/z 915 (pentose-methylpentose-pentose), 727 (pentose-hexose), 639 (pentose-methylpentose) and 349 (terminal pentose). But there were no ions associated with the terminal hexosyl and the terminal methylpentosyl units. On selective cleavage of an ester glycoside linkage with LiI in a mixture of abs. MeOH and 2,6-lutidine,⁵⁾ 3 gave a monodesmoside 5 and a methyl trisaccharide 6. On acid hydrolysis, 5 gave arabinose and glucose. The EI-MS spectrum of a peracetate of 5 exhibited the fragment ions characteristic of terminal arabinose (m/z 259) and arabinosyl-glucose (m/z 547). The sequencing analysis of a permethyl ether of 5 indicated the presence of a terminal arabinopyranosyl unit and a 2-linked glucopyranosyl unit. These results coupled with the inspection of carbon signals due to the sugar moiety (Table I) led to the formulation of 5 as shown in Chart 1. The sequencing analysis of a permethyl ether of 6 showed the presence of a 2-linked arabinopyranosyl unit, a 3-linked rhamnopyranosyl unit and a terminal xylopyranosyl unit. Coupled with the fragmentation of the EI-MS of TMSi ether of 3 (*vide supra*), 6 was formulated as methyl D-xylopyranosyl-(1 \rightarrow 3)-L-rhamnopyranosyl-(1 \rightarrow 2)-L-arabinopyranoside. By reference to the study on the unusual glycosylation shifts observed for the 2-linked arabinopyranosyl esters of oleanane type triterpenes,⁶⁾ the sugar carbon signals of 3 were assigned as shown in Table I. Based on these results, the structure of 3 was established as shown in Chart 1.

On very mild alkaline hydrolysis with 0.25% KOH-H₂O at 0°C, 1 gave 7. The ^{13}C -NMR spectrum (in pyridine- d_5) of 7 showed signals due to two ester carbonyl carbons at δ 176.3 (28-glycosyl ester) and 171.7 as well as a signal due to a free carboxyl carbon at δ 174.9, indicating that one of two ester groups of the hydroxymethylglutarate moiety of 1 was saponified by this reaction. Hydrolysis of 1 with 0.5N BaO in MeOH at room temperature yielded a methyl ester (8) of 7. The EI-MS spectrum of a TMSi ether of 8 exhibited the fragment ions associated with terminal pentose (m/z 349), arabinosyl-glucose (m/z 727) and (xylosyl-rhamnose)-OCO-CH₂-CCH₃(OH)-CH₂-COOCH₃ (m/z 797). This indicated that the hydroxymethylglutarate moiety of 8 (and 7) must be located on either the rhamnosyl unit or the terminal xylosyl unit. Comparison of the ^{13}C -NMR spectrum of 7 with that of 3 (Table I) in view of the acylation shift⁷⁾ revealed the allocation of the acyl group at the 4-hydroxyl group of the rhamnosyl unit. Recently, R.-S. Xu and his co-workers also isolated 1 from the same crude drug as ours and found by 2D-NMR analysis that another ester linkage of the hydroxymethylglutarate moiety of 1 must be located at the 4-hydroxyl group of the terminal arabinosyl unit.^{8,9)}

Finally, the chirality of the asymmetric carbon of the acyl moiety of 1 was established as follows. In order to reduce free COOH of 7 to CH₂OH, a peracetate of 7 was treated with diborane in tetrahydrofuran. Alkaline saponification of the product gave mevalonolactone (9) and 3. Hirai *et al.*¹⁰⁾ reported the micro-scale identification of the chirality of optically active mevalonolactone by conversion into 3(*R* or *S*)-5-*O*-acetyl-1-[(*R*)-phenylethyl]-mevalonamide (10) and subsequent HPLC analysis. We found that TMSi ethers (11) of the isomers of 10 can be distinguished from each

Chart 1

Table I. ^{13}C -Chemical Shifts of Sugar Moieties in Pyridine- d_5

	3	5	7	3	7
3-O-				28-O-	
Glc-1	103.4	103.4	103.4	Ara-1	93.5
2	83.5	83.6	83.4	2	75.6
3	77.9 ^a	78.0 ^a	78.0	3	70.1
4	71.8 ^b	71.8	71.6 ^b	4	66.2
5	78.0 ^a	78.1 ^a	78.0	5	63.1
6	62.4	62.5	62.3		
				Rha-1	101.4
Ara-1	106.5	106.6	106.4	2	71.2 ^b
2	73.8	73.9	73.7	3	83.2
3	74.3	74.4	74.2	4	72.8
4	69.2	69.3	69.2	5	70.1
5	67.2	67.3	67.1	6	18.4
				Xyl-1	107.2
				2	75.0
				3	78.3 ^a
				4	71.0 ^b
				5	67.2

a.b: These assignments may be reversed.

12: R=-H

13: R=- β -Glc- β -Glc

Table II. Solubilizing Effect of 1, 3 and 13 (1mg/ml) on Saponin A (12) in H₂O

Compd.	A(mg/ml)	B(mg/ml)	C
None	1.6	0.017	-
13	24	16*	22
3	24	21*	28
1	30	30	44

A: amount of 12, B: solubility of 12, C: mol ratio of solubilized 12 to bisdesmosides, at 37°C, *: saturated soln.

Table III. Solubilizing Effect of 1, 3 and 13 (1mM) on Yellow OB in Buffer

Compd.	solubility of Yellow OB
None	Insoluble
13	13.0μM
3	12.3μM
1	78.4μM

M/80 phosphate buffer (pH 6.5 ionic strength 0.02). at 37°C.

other by GC-MS (on a column packed with 5% SE-52 on Chromosorb-W). By this modified procedure, the chirality of the C-3 of 9 was found to be the S-configuration. Consequently, the structures of 1, 7 and 8 including the stereochemistry of the acyl moiety were established as shown in Chart 1. The name, cyclic bisdesmoside is proposed for glycosides of this type.

Recently, it has been found that the solubility in water of the monodesmosides of pericarps of *Sapindus mukurossi* Gaertn., such as saponin A (12) is markedly increased by the bisdesmosides such as mukurozi-saponin Y₂ (13)¹¹⁾ and the sesquiterpene-oligoglycosides¹²⁾ which co-occur in this plant. It was also found that the water solubility of a synthetic dye, Yellow OB was also increased by 13.¹³⁾ As shown in Table II and Table III, 1 is a more potent solubilizer than non-cyclic bisdesmosides 3 and 13. This indicates the significance of the cyclic structure of this type in the surface activities of saponins.

ACKNOWLEDGEMENT We are grateful to Prof. S. Nozoe, Pharmaceutical Institute, Tohoku University, for the gift of an authentic sample of 3(R)-mevalonolactone. One of the authors, O. Tanaka, wishes to thank to Dr. R.-S. Xu, Shanghai Institute of Materia Medica, Science Academy of China for his communication about the 2D-NMR analysis prior to publication. Thanks are also due to Dr. H. Matsuura, Wakunaga Pharm. Ind. Co. Ltd., Hiroshima for measurement of the NMR spectra.

REFERENCES AND NOTES

- 1) R. Oshima, J. Kumanotani and C. Watanabe, *J. Chromatogr.*, **259**, 159 (1983).
- 2) R. A. Eade, J. J. H. Simes and B. Stevenson, *Aust. J. Chem.*, **16**, 900 (1963).
- 3) H. Ishii, I. Kitagawa, M. Matsushita, K. Shirakawa, K. Tori, T. Tozyo, M. Yoshikawa and Y. Yoshimura, *Tetrahedron Lett.*, **22**, 1529 (1981).
- 4) R. Kasai, M. Okihara, J. Asakawa, K. Mizutani and O. Tanaka, *Tetrahedron*, **35**, 1427 (1979).
- 5) K. Ohtani, K. Mizutani, R. Kasai and O. Tanaka, *Tetrahedron Lett.*, **25**, 4537 (1984).
- 6) K. Mizutani, K. Ohtani, R. Kasai, O. Tanaka and H. Matsuura, *Chem. Pharm. Bull.*, **33**, 2266 (1985).
- 7) K. Yamasaki, R. Kasai, Y. Masaki, M. Okihara and O. Tanaka, *Tetrahedron Lett.*, **1977**, 1231 (1977).
- 8) R.-S. Xu *et al.*, *Tetrahedron Lett.*, to be submitted.
- 9) The allocation of the ester linkage on the 2-hydroxyl group of the terminal arabinoside moiety, proposed by our group at 27th Symposium on The Chemistry of Natural Products (Oct., 1985 in Hiroshima), should now be revised.
- 10) N. Hirai and K. Koshimizu, *Phytochem.*, **20**, 1867 (1981).
- 11) H. Kimata, T. Nakashima, S. Kokubun, K. Nakayama, Y. Mitoma, N. Yata and O. Tanaka, *Chem. Pharm. Bull.*, **31**, 1998 (1983).
- 12) R. Kasai, H. Fujino, T. Kuzuki, W.-H. Wong, C. Goto, N. Yata, O. Tanaka, F. Yasuhara and S. Yamaguchi, *Phytochem.*, **25**, 871 (1986).
- 13) K. Nakayama, H. Fujino (*née* Kimata), R. Kasai, Y. Mitoma, N. Yata and O. Tanaka, *Chem. Pharm. Bull.*, **34**, 3279 (1986).

(Received July 23, 1986)