



Agapetes subsessilifolia (Ericaceae), a New Species from the Eastern Himalayas

Author(s): Huang Su-hua, Sun Hang, Zhou Zhe-kun

Reviewed work(s):

Source: *Novon*, Vol. 8, No. 4 (Winter, 1998), pp. 378-379

Published by: Missouri Botanical Garden Press

Stable URL: <http://www.jstor.org/stable/3391859>

Accessed: 19/12/2011 03:32

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at

<http://www.jstor.org/page/info/about/policies/terms.jsp>

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.



Missouri Botanical Garden Press is collaborating with JSTOR to digitize, preserve and extend access to *Novon*.

<http://www.jstor.org>

Agapetes subsessilifolia (Ericaceae), a New Species from the Eastern Himalayas

Huang Su-hua

Department of Biology, Yunnan University, Kunming, Yunnan 650091,
People's Republic of China

Sun Hang and Zhou Zhe-kun

Herbarium, Kunming Institute of Botany, The Chinese Academy of Sciences, Kunming,
Yunnan 650204, People's Republic of China

ABSTRACT. *Agapetes subsessilifolia* S. H. Huang, H. Sun & Z. K. Zhou, a new species of Ericaceae from the Gorge of Yalu Tsangpo, Medog, Tibet, in the eastern Himalayas, is described and illustrated.

During a botanical expedition to the big bend gorge of Yalu Tsangpo of southeastern Tibet, the following distinct new species of Ericaceae was discovered.

Agapetes subsessilifolia S. H. Huang, H. Sun & Z. K. Zhou, sp. nov. TYPE: China. Tibet: Medog, Hammi, 1600 m, Oct. 1992, *Sun Hang et al.* 0997 (holotype, KUN). Figure 1.

Species *Agapetes pseudo-griffithii* Airy-Shaw affinis, sed petiolis valde brevis, pedicellis apice leviter in clavatos expansis, calycis segmentis anguste triangulatis, corollis cylindricis, 2.5 cm longis differt.

Frutex epiphyticus, rami graciles, brunnei, glabri, leviter angulati. Folia subopposita vel in ramulis congestis, coriacea, ovato-oblonga vel late elliptica, 7.5–14 cm longa, 3–3.5 cm lata, apice longe acuminata vel caudata, basi rotundata, margine subintegra, utrique glabra, costa utrinque prominula. Nervis lateralibus obscuris; petiolus valde brevis, 2–3 mm longus, diam. 3 mm, glaber. Inflorescentiae corymbiformes, 3–5 florae, axillares vel in ramis veteribus, pedunculo 2.5–3.5 cm longo; pedicelli graciles, 3 cm longi, apice leviter clavati expansi; receptaculum 4 mm longum, 3 mm latum; calyxis segmenta 5, anguste triangulares, 5 mm longa, 4 mm lata, acuta. Corolla cylindrica, rubra, transversaliter V-formiter purpureo-striata, 2.5 cm longa, supra leviter ampliata, 1 cm lata, lobis 5, stamina 10, deltoidea; filamenta valde brevia, ca. 1 mm longa, leviter puberula; thecae 5–6 mm longae, dense papillose, rostris 16 mm longis, dorso ecalcaratis apice poris

dehiscentibus; discus annulatus, ca. 0.5 mm altus; ovarium 5-loculare. Stylus cylindricus, 2.5 cm longus; Stigma truncatum, glabrum. Frutus ignotus.

Epiphytic shrub, glabrous. Leaves subopposite, clustered at the top of branches, coriaceous, ovate-oblong to widely elliptic, 7.5–14 × 3–5.5 cm, apex narrowly acuminate, base rounded, glabrous, margins entire; petioles very short, 2–3 mm long, or sessile, glabrous. Corymbs 3–5-flowered, axillary on old branches, peduncle 2.5–3.5 cm, pedicels 3 cm, glabrous, clavate-shaped at apex. Sepals narrowly triangular, 5 × 4 mm, acute. Corolla tubular, red with V-shaped purplish striated, 2.5 × 1 cm, triangular lobes. Filaments very short, ca. 1 mm, slightly pubescent; thecas 5–6 mm; rostrum 16 mm long, glabrous, not spurred. Disk annular, 0.5 mm tall; ovary 5-locular, style 2.5 cm, stigma truncate, glabrous. Fruit unknown.

Agapetes subsessilifolia is similar to *A. pseudo-griffithii* Airy-Shaw but differs in having very short or sessile petioles; pedicels that are clavate-shaped at the apex; calyx lobes narrowly triangular; and cylindrical corollas to 2.5 cm long.

Paratypes. CHINA. Tibet: Medog, Gelin, 1750 m, Dec. 1992, *Sun Hang et al.* 2537 (KUN), 2661 (KUN); Buqiong Lake, 1600 m, Nov. 1992, *Sun Hang et al.* 1140 (KUN).

Acknowledgments. We are very grateful to the National Natural Science Foundation of China for supporting this expedition to Tibet; to Wu Xi-lin for the excellent illustration, and to Zhu Guanghua for reviewing the manuscript.




Figure 1. *Agapetes subsessilifolia* S. H. Huang, H. Sun & Z. K. Zhou. —A. Flowering branch ($\times 1$). —B. Opened corolla, adaxial view showing androecium ($\times 3$). —C. Stamen, abaxial view ($\times 3$). —D. Stamen, lateral view ($\times 3$). —E. Calyx and persistent style ($\times 3$).