

野牡丹科异药花属一新种

陈 介

(中国科学院昆明植物研究所)

A NEW SPECIES OF GENUS FORDIOPHYTON STAPF (MELASTOMATACEAE)

CHEN CHEIH

(Kunming Institute of Botany, Academia Sinica)

短茎异药花(新种)

本新种与本属其他种的主要区别是，植株矮，具匍匐茎；茎长2—3厘米，节上具刺毛，其余无毛；花序总梗(或花萼)长4.5—11厘米。

Fordiophyton brevicaule C. Chen, sp. nov.

A congenericis caulibus erectis 2—3 cm longis, praeter ad nodos setosis glabris; pedunculis (vel scapigera) 4.5—11 cm longis recedens.

Herba ca. 15 cm alta (cum inflorescentia), stolonifera; caule erecto obtuse quadrangulari, 2—3 cm longo, glabro, ad nodum setoso. Folia papryacea vel submembranacea, elliptica ad ovato-elliptica, apice acuta, basi cordata, 3.5—8 cm longa, 2—5 cm lata, margine paulo vadoseque serrulata, 7-nervia, supra dense parveque verruculata, glabra, nervis primariis planis lateribus inconspicuis subtus prominentibus conspicuis laxe setosis, ceterum glabra; petiolis sulcatis, 1—3 cm longis, utrinque latere setosis, subtus pilis glandulosis laxe ornatis. Panicula terminalis, e cymis constans, tantum

semel ramosa, 9—15 cm longa, pedunculata (vel scapigera); pedunculis 4.5—11 cm longis, glabris; bracteis 2, lanceolatis, ca. 6 mm longis; pedicellis ca. 5 mm longis, fructificatione ad 1.3 cm longis, glabris; calyce anguste infundibulari, tetragono, tubo 4—6 mm longo glabro diaphano, lobis lanceolatis apice acuminatis, ca. 7 mm longis 2 mm latis; petalis ovatis, rubris, ca. 9 mm longis, 5 mm latis, apice glanduloso-pilosus; staminibus longioribus ca. 14 mm longis, anthera linearis ca. 8 mm longa basi obtuse cornuta, connectivo apice leviter producto, eis brevioribus ca. 6.2 mm longis, anthera ovata ca. 1.2 mm longa basi non tuberculata, connectivo non producto; ovario obovato, apice truncato, membranaceo-coronato. Capsulae late obconicae vel subpoculiformes, ca. 6 mm diam.

Hongkong: in humidis juxta fluvium, Jun. 1. 1933, C. Wang 32407 (Typus! Herb. Inst. Bot. Austro-Sin., Acad. Sin.).