

北美外来植物中原产东亚类群的学名考证^{*}

马金双

(中国科学院昆明植物研究所, 云南 昆明 650204)

摘要: 作者在整理北美外来入侵植物中发现一些起源(或主要分布)于东亚植物的学名, 在北美乃至欧洲使用非常混乱。本文特将有关重要类群整理出来, 包括异名、原产地、北美的分布以及必要的讨论等。

关键词: 外来植物; 北美; 原产地; 东亚; 分类群; 修订

中图分类号: Q 949

文献标识码: A

文章编号: 0253-2700 (2010) 01-014-11

Nomenclatural Notes on Alien Vascular Plants in North America Originated from East Asia

MA Jin-Shuang

(Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, 650204, China)

Abstract: The scientific names of twenty-three alien vascular plants in North America, originating from East Asia, are reviewed here, since their uses in North America are different from those in East Asia. Most are taken from the invasive plants literatures in North America as well as in Europe, but a few are also from native floras in East Asia. The synonyms for each accepted species, along with their distributions both in North America and in East Asia, with discussion if necessary, are also provided.

Key words: Invasive Plants; North America; Native to East Asia; Taxonomic Notes

1. *Alhagi maurorum* Medik. in Vorles. Churpfalz. Phys. -Ocon. Ges. 2 : 397, 1781. Syn. *A. pseudalhagi* (M. Bieb.) Desv. in J. Bot. Agric. 1 : 120, 1813, *A. graecorum* Boiss., Diagn. Pl. Or. Nov. 2 (9) : 114, 1849 (Fabaceae).

Camel-thorn is a native species found in northwest China (north Gansu, west Nei Menggu, west Qinghai, Xinjiang), northern India, and central and west Asia as well as North Africa, naturalized in many parts of world, including southern Africa, Australia and USA: AZ, CA, CO, ID, NM, NV, TX, UT, WA.

Yakovlev (1979) suggested that the five species of *Alhagi* in 'Flora of USSR' should be treated as one species, and Awmack & Lock (2002) treated

all species of *Alhagi* as *A. maurorum*, with synonyms including *A. sparsifolia* of central Asia and China (Xu, 1998), *A. pseudalhagi* (Ball, 1968; Bossard et al., 2000) and *A. graecorum* of Europe (Ball, 1968).

2. *Ampelopsis heterophylla* (Thunb.) Sieb. & Zucc. in Abh. Math. -Phys. Cl. Konigl. Bayer. Akad. Wiss. 4 (2) : 197, 1845. Syn. *Vitis heterophylla* Thunb. in Murray, Syst. Veg. ed. 14 : 244, 1784, & Fl. Jap. 103, 1784 (Vitaceae).

Porcelain Berry, a native species from far east Russia, northern Japan, Korea and eastern China, to Himalayas, Indochina and the Philippines, and has been cultivated since the 1870s in

* Received date: 2009-04-20, Accepted date: 2009-11-26

作者简介: 马金双 (1955-) 男, 研究员, 主要从事植物分类学研究。

USA : CT, DC, DE, GA, IL, KY, MA, MD, MI, NC, NH, NJ, NY, OH, PA, RI, VA, WI, WV.

This is an extremely variable species, and many varieties and forms have been described both from the native area and from under cultivation (see below for detail). Moreover, there is still no consensus on nomenclature even in the native area: the Japanese treated it as *A. glandulosa* (Momiyama, 1977; Ohba, 1999), the Chinese treated it as *A. heterophylla* (Li, 1996; 1999), but the same taxon was recorded as *A. brevipedunculata* in North America (Antenen, 1996) and Europe (Richardson, 1997). Here, a broad concept is adopted, without further division, and the most common synonyms of this species includes at least the following:

A. glandulosa (Wall.) Momiy. in Fl. E. Himalaya 2 : 78, 1971, & var. *glabrifolia* (Honda) Momiy. in J. Jap. Bot. 52 : 30, 1977; *A. brevipedunculata* (Maxim.) Trautv. in Trudy Imp. St. Peterb. Bot. Sada 8 : 178, 1883, & var. *hancei* (Planch.) Rehder in J. Arn. Arb. 2 : 177, 1921, & var. *glabrifolia* Honda in Bot. Mag. Tokyo 45 : 423, 1931; *A. heterophylla* (Thunb.) Sieb. & Zucc. var. *brevipedunculata* (Regel) C. L. Li in Chin. J. Appl. Envirn. Biol. 2 (1) : 47. 1996, & var. *hancei* Planch. in DC. Monogr. Phan. 5 : 457, 1887, & var. *glabrifolia* (Honda) Nemoto, Fl. Jap. Suppl. : 463, 1936, & var. *kulingensis* (Rehd.), C. L. Li in Chin. J. Appl. Envirn. Biol. 2 (1) : 47, 1996, & f. *elegans* (K. Koch) Voss, Vilm. Blumeng. ed. 3, 1 : 183, 1894, & var. *citrulloides* (Lebas) Rehder in L. H. Bailey, Stand. Cycl. Hort. 1 : 278, 1914; *A. humulifolia* Bunge var. *heterophylla* (Thunb.) Koch, Hort. Dendr. 48, 1853; *A. brevipedunculata* (Maxim.) Trautv. in Trudy Imp. St. Peterb. Bot. Sada 8 : 178. 1883, & var. *maximowiczii* (Regel) Rehder in L. H. Bailey, Gentes Herb. 1 : 36,

1920, & var. *citrulloides* (Lebas) Rehder in L. H. Bailey, Gentes Herb. 4 : 135, 1923, & var. *kulingensis* Rehder in L. H. Bailey, Gentes Herb. 1 : 36, 1920, & f. *citrulloides* (Lebas) Rehder in J. Arn. Arb. 2 : 176, 1921, & f. *elegans* (K. Koch) Rehder in J. Arn. Arb. 2 : 176, 1921, & var. *citrulloides* (Lebas) Rehder in J. Arn. Arb. 4 : 135, 1923, & var. *elegans* (K. Koch) L. H. Bailey in Gentes Herb. 4 : 135, 1923, & var. *heterophylla* (Thunb.) Hara, Enum. Sperm. Jap. 3 : 133, 1954; *A. regelianiana* Carr. in Rev. Hort. 440, 1866; *Cissus brevipedunculata* Maxim., Prim. Fl. Amur. 68 - 69, 1859. *Vitis glandulosa* Wall., Fl. Ind. 2 : 479, 1824; *V. heterophylla* Thunb. var. *maximowiczii* Regel in Trudy Imp. St. - Peterb. Bot. Sada 2 : 392, 1873, & var. *elegans* Regel in Gartenfl. 22 : 197, 1873, & var. *vestita* Rehd. in Mitt. Deutsch. Dendr. Ges. 21 : 189, 1912; *V. humulifolia* Bunge f. *glabra* Debeaux in Acta Linn. Soc. Bordeaux 31 : 132, 1876.

3. *Aralia elata* (Miq.) Seem in J. Bot. 6 : 134, 1868. Syn. *Dimorphanthus elatus* Miq., Comm. Phytoogr. 95, t. 12, 1840; *A. spinosa* var. *elata* (Miq.) Sarg., Silva 5 : 60, 1903.

Japanese Angelica-Tree, native to northeast, south and southwest China, Japan, Korea and far east Russia (Wen, 2000), has been widely cultivated in North America since the 1830s, but naturalized and escaped from the Middle Atlantic region to New England in the 1900s. However, it was misidentified as *A. spinosa* (Lamont and Fitzgerald, 2001; Zebryk, 2003; Sorrie, 2005), a native species of North America from the Middle Atlantic to the southern and southeastern USA, with a long main axis of inflorescence at the top of the stem. This species, however, has no main axis of inflorescence at all, with fruit and seed smaller than that of *A. spinosa* (Moore *et al.* in print). Naturalized in USA: MA, NH, NJ, NY, PA; and Canada: ON.

4 . *Chenopodium album* L. , Sp. Pl. 1 : 219.
1753 (Chenopodiaceae)

White Goosefoot , native to Eurasia and/or introduced into North America is still uncertain in origin (Zhu , 1979 ; Clemants and Mosyakin , 2003) , but is certainly known as a worldwide weed. This is one of the worst weeds and most widespread plants in the world. Hundreds of segregate microspecies and infraspecific entities of the *C. album* aggregate have been described and/or recognized by various authors. Some authors have recognized numerous segregate intergrading species , while others have developed elaborate infraspecific hierarchies with numerous subspecies , varieties , forms , and even numerous subforms , or have combined both approaches. Unfortunately , neither approach has brought satisfactory and uncontroversial results (Clemants and Mosyakin , 2003). Widespread in USA : AK , AL , AR , AZ , CA , CO , CT , DC , DE , FL , GA , HI , IA , ID , IL , IN , KS , KY , LA , MA , MD , ME , MI , MN , MO , MS , MT , NC , ND , NE , NH , NJ , NM , NV , NY , OH , OK , OR , PA , RI , SC , SD , TN , TX , UT , VA , VT , WA , WI , WV , WY; and Canada : AB , BC , LB , MB , NB , NF , NS , NT , ON , PE , QC , SK , YT.

5 . *Dioscorea polystachya* Turz. in Bull. Soc. Imp. Nat. Mosc. 10 (7) : 158 , 1837 ; Raz in Fl. N. Amer. 26 : 483 , 2002. Syn. *D. batatas* Decne. Rev. Hort. 3 : 243 , 1854 (Dioscoreaceae).

The Chinese yam originates from eastern China (northeast to southwest) , cultivated as edible tubers , and was introduced into USA as an ornamental vine. In 1970s , it had not yet been documented as escaping from cultivation. By 1986 , however , it was reported to have become naturalized and was observed in areas outside of cultivation in USA : AL , AR , CT , DC , GA , IL , IN , KS , KY , LA , MD , MO , MS , NC , NJ , NY , OH , OK , PA , SC , TN , VA ,

VT , WV.

This species has been frequently misidentified as *D. batatas* in various references in Europe (King , 1986) and North America (Zheng *et al.* , 2004) and *D. oppositifolia* L. (Mueller *et al.* , 2003). The name of *D. opposita* Thunb. (Fl. Jap. 151 , 1784) has also been used in the sense of *D. polystachya* , which is an earlier name for *D. batatas*. It was based on a misidentification of Japanese material as the strictly Indian species *D. oppositifolia* L. , which was cited as a synonym in the protologue , thus rendering *D. opposita* nomenclaturally superfluous as well as an illegitimate name (Ting and Gilbert , 2000) .

6 . *Elymus repens* (L.) Gould in Madroño 9 (4) : 127. 1947. Syn. *Elytrigia repens* (Linn.) Nevsk. in Act. Inst. Bot. Acad. Sci. URSS 1 (1) : 14 , 1933 ; *Agropyron repens* (L.) P. Beauv. Ess. Agrostogr. 102 , 146 , 180 , t. 20 , f. 2. 1812 ; *Triticum repens* L. , Sp. Pl. 1 : 86 , 1753 (Poaceae).

Creeping Wild Rye is native to Eurasia (temperate Europe and central Asia , northwest China) , but can also be found in many parts of South America , Australia , New Zealand and Indonesia ; it is also widely distributed across North America , especially invasive in grass prairie zones and/or wetlands of western North America. USA : AK , AR , AZ , CA , CO , CT , DC , DE , IA , ID , IL , IN , KS , MA , MD , ME , MI , MN , MO , MT , NC , ND , NE , NH , NJ , NM , NV , NY , OH , OK , OR , PA , RI , SD , TN , TX , UT , VA , VT , WA , WI , WV , WY; Canada : AB , BC , LB , MB , NB , NF , NS , NT , ON , PE , QC , SK , YT.

This species is widely accepted as *Elymus repens* in Europe (Melderis , 1980) and much of the world (Mabberley , 1992; Plant Book , p. 203) , but the former Russian name is still followed by the Chinese as *Elytrigia* in China (Kuo and Tsui , 1987) .

7. *Fallopia aubertii* (Henry) Holub in *Folia Geobot. Phytotax.* 6 : 176. 1971; A. J. Li in *Fl. Reipubl. Popularis Sin.* 25 (1) : 102, 1998 & A. R. Li, A. E. Grabovskaya-Borodina, S. P. Hong, J. McNeill, H. Ohba, & C. W. Park in *Fl. Chin.* 5 : 317, 2003. Syn. *Polygonum aubertii* Henry in *Rev. Hort.* 1907: 82, f. 23 & 24, 1907 (Polygonaceae).

China Fleece Vine is a native species from north to southwest China (Gansu, Guizhou, Henan, Hubei, Nei Menggu, Ningxia, Qinghai, Shaanxi, Shanxi, Sichuan, Yunnan, and Xizang). Some authors put this species under *Polygonum* (Akeroyd, 1989; Sorrie, 2005), and some even consider it to be the same as *Fallopia baldschuanica* (Regel) Holub (Bailey and Stace, 1992), based on *P. baldschuanicum* Regel (Trudy Imp. St. Peterb. Bot. Sada 8 : 684. 1883) in USA: CO, MA, MI, NY, PA, UT, VA, WA

8. *Firmiana simplex* (L.) W. Wight in *USDA Bur. Pl. Industr. Bull.* 142 : 67, 1909. Syn. *F. platanifolia* (Linn. f.) Marsili in *Sagg. Sci. Lett. Acc. Padova* 1 : 106 - 116, 1786; *Sterculia platanifolia* L. f., *Suppl. Pl.* 423. 1781, *Culhamia simplex* (L.) Nakai in *Bull. Nat. Sci. Mus. Tokyo* 31 : 76, 1952; *Hibiscus simplex* L., *Sp. Pl. ed. 2* : 977, 1763 (Sterculiaceae).

Chinese Parasol Tree, native to China (from north to south, also cultivated widely) and Japan, is treated as *F. platanifolia* in China (Hsue, 1984). USA: AL, AR, FL, GA, LA, MD, MS, NC, SC, TN, TX, VA.

9. *Frangula alnus* Miller, *Gard. Dict.* (ed. 8), No. 1, 1768; Syn. *Rhamnus frangula* L., *Sp. Pl.* 1 : 193. 1753. (Rhamnaceae).

Glossy buckthorn, native to Europe, Asia and North Africa, is commonly accepted as *Frangula alnus* Miller in Europe (McKean, 1997; Tutin, 1968). However, it was treated

as *Rhamnus frangula* L. in China (distributed in Xinjiang only, Chen, 1982) and in North America (McClain, 1996; Krock and Williams, 2002, Frappier and Eckert, 2003; Sorrie, 2005). It was introduced to North America prior to the 1900s for horticultural purposes, and is widespread in USA: CO, CT, IA, IL, IN, KY, MA, MD, ME, MI, MN, NH, NJ, NY, OH, PA, RI, TN, VA, VT, WI, WV, WY; and Canada: MB, NB, NS, ON, PE, QC.

10. *Humulus scandens* (Lour.) Merr. in *Trans. Amer. Philip. Soc.* n. ser. 24, 2 : 138, 1935; S. S. Chang in *Fl. Reip. Pop. Sin.* 23 (1) : 220 - 221, 1998; Z. Y. Wu, Z. K. Zhou & B. Bartholomew in *Fl. Chin.* 5 : 75, 2003. Syn. *Antidesma scandens* Lour. *Fl. Cochinch.* 2 : 157, 1790 (Cannabaceae).

Japanese Hop has probably arrived in North America due to intentional cultivation at the end of the nineteenth century. Native to China (national wide except Qinghai and Xinjiang), Japan, Korea and Vietnam, it is also naturalized in Europe, but frequently used as *Humulus japonicus* Sieb. & Zucc. both in Europe (Nelson, 1989) and in North America (Small, 1978; Sorrie, 2005). Widespread in USA: CT, DC, DE, GA, IA, IL, IN, KS, KY, MA, MD, ME, MI, MN, MO, NC, NE, NH, NJ, NY, OH, PA, RI, SC, TN, VA, VT, WI, WV; and Canada: ON, QC.

11. *Kochia scoparia* (L.) Schrader in *Neues J. Bot.* 3 : 85. 1809; G. L. Zhu in *Fl. Reip. Pop. Sin.* 25 (2) : 102, 1979, G. L. Zhu, S. L. Mosyakin & S. E. Clemants in *Fl. Chin.* 5 : 385, 2003. Syn. *Bassia scoparia* (L.) A. J. Scott. in *Fed. Rep.* 81 (2 - 3) : 108, 1978; *Chenopodium scoparium* L., *Sp. Pl.* 1 : 221. 1753 (Chenopodiaceae).

Mexican Summer Cypress from Eurasia and Africa, cultivated throughout China, widely

naturalized in Australia , and North and South America , is an extremely variable species , and several forms , varieties , and subspecies have been described (Mosyakin , 2003) . The nomenclature is also extremely confusing since it has been classified either as *Bassia* (Ball and Akeroyd , 1993) or *Chenopodium* in history. Widely spread in USA: AZ, CA , CO , CT , DE , IA , ID , IL , IN , KS , KY , MA , ME , MI , MN , MO , MS , MT , NC , ND , NE , NH , NJ , NM , NV , NY , OH , OK , OR , PA , RI , SD , TN , TX , UT , VA , VT , WA , WI , WV , WY; and Canada : AB , BC , MB , NB , NS , ON , QC , SK.

12 . *Landoltia punctata* (G. Mey.) Les & D. J. Crawford , Novon 9 : 532. 1999. Syn. *Spirodelta punctata* (G. Mey.) C. H. Thompson in Ann. Rep. Miss. Bot. Gard. 9 : 28 , 1898 ; *Lemna punctata* G. Meyer in Prim. Fl. Esseq. 262 , 1818 ; *Spirodelta oligorrhiza* (Kurz) Hegelm. (Lemnaceae).

Dotted Water-flaxseed , native to warm-temperate to subtropical regions with mild winters in China (Fujian , Henan , Hubei , Sichuan , Taiwan , Xizang , Yunnan , and Zhejiang) , India , Indonesia , Japan , Malaysia , the Philippines , Thailand , Vietnam , and also found in Africa , Australia , North America , the Pacific Islands , and South America. It has been more often treated in *Lemna* (Crow and Hellquist , 2000) in North America or *Spirodelta* (Lawatree , 1980) in Europe. Wide spread in USA: AL , AR , AZ , CA , DC , DE , FL , GA , IL , KY , LA , MA , MO , MS , NC , OK , OR , PA , SC , TN , TX , VA.

13 . *Lespedeza thunbergii* (DC.) Nakai in Bull. For. Exp. Sta. Chosen. 6 : 1 - 101 , 1927. Syn. *L. formosa* (Vog.) Koehne , Deuts. Dendrol. 343 , 1893 ; *Desmodium formosum* Vog. In Nov. Act. Acad. Leop. -Carol. 19 , suppl. 29 , 1842 ; *D. thunbergii* DC. Prodr. 2 : 337 , 1825

(Fabaceae) .

Thunberg's Bushclover , from eastern Asia (from Japan to India) , has been treated as *L. formosa* in China (Li and Chen , 1995) , and is widespread in USA: AR , CT , FL , GA , IL , IN , KY , LA , MA , MD , MI , MO , NC , NJ , NY , PA , SC , VA ; and Canada : ON.

14 . *Lithospermum arvense* L. Sp. Pl. 132 , 1753 ; Syn: *Buglossoides arvensis* (L.) Johnston in J. Arn. Arb. 35 : 42 , 1954 (Boraginaceae) .

Corn Gromwell , a native of Eurasia , is treated as *Lithospermum* in China (Wang , 1990 ; Zhu et al. , 1995) , but as *Buglossoides* in Europe (Fernandes , 1972) and in North America (Sorrie , 2005) . Wide spread in USA: AL , AR , CA , CO , CT , DC , DE , FL , GA , IA , ID , IL , IN , KS , KY , LA , MA , MD , ME , MI , MN , MO , MS , MT , NC , ND , NE , NH , NJ , NY , OH , OK , OR , PA , RI , SC , SD , TN , TX , UT , VA , VT , WA , WI , WV , WY; and Canada : BC , MB , NS , ON , QC.

15 . *Lonicera standishii* Jacques in J. Soc. Nat. Hort. Paris 5 : 514 , 1859. Syn. *Lonicera standishii* Carr. in Fl. Ser. Jard. 1 Eur. 13 : 63. 1860 (Caprifoliaceae) .

Standish's Honeysuckle , from Eurasia , was introduced in around the 1800s , as ornamental and wildlife plants , and is often recorded as *L. standishii* 'Carr.' , later a synonym in USA : IL , KY , MD , NY , PA , VA.

16 . *Lygodium scandens* (L.) Sw. in Schrad. Journ. 106 , 1801 ; R. C. Ching in Fl. Reip. Pop. Sin. 2 : 109 , 1959. Syn. *L. microphyllum* (Cav.) R. Br. Prod. Fl. N. Holl. 162 , 1810 ; *Ugena microphylla* Cav. Icon. Descrip. Pl. 6 : 76 , pl. 595 , f. 2 , 1801 ; *Ophioglossum scandens* L. Sp. Pl. 2 : 1063 , 1753 (Lygodiaceae) .

The Old World climbing fern , native to Old

World tropics and subtropics, from south and southwest China to tropical Asia, is treated as *L. microphyllum* in North America (Zheng *et al.*, 2005). It was recognized as a potential problem not long after naturalization in USA: FL.

17. *Nephrolepis hirsutula* (Forst.) Presl, Tent. Pterid. 79, 1836. Syn. *Davallia multi-flora* Roxb. in Calcutta J. Nat. Hist. & Misc. Arts & Sci. India 4: 515, pl. 31, 1844; *Poly-podium hirsutulum* Forst., Prodr. 81, 1786; *N. multi-*

flora (Roxb.) Jarret & Morton in Contr. US Nat. Herb. 38: 309, 1974 (Dryopteridaceae or Nephrolepidaceae).

The Asian Sword fern has been treated in Flora Reipublicae Populus Sinicae twice in different families, Dryopteridaceae (Ching, 1959) and Nephrolepidaceae (Wu, 1999), and was also frequently recorded as a member of *Polypodium* in North America. Native to tropical Africa and tropical Asia (China: Taiwan, Fujian, Guangdong, Guangxi, Hainan, and Yunnan and south Japan), it is cultivated in Europe (Page and Bennell, 1986), naturalized in Florida and Hawaii of the United States, the Caribbean, tropical South America, and the Mascarenes.

18. *Polygonum posumbu* Buch.-Ham. ex D. Don, Drodri. Fl. Nep. 71, Feb. 1825; A. J. Li in Fl. Reip. Pop. Sin. 25 (1): 29 - 30, 1998. Syn. *Persicaria posumbu* (Buch.-Ham. ex D. Don) H. Gross in Bot. Jahrb. Syst. 49 (2): 313, 1913; *P. caespitosum* Blume Bijdr. 532, Dec. 1825 (Polygonaceae).

The Asiatic Water Pepper has been treated either as *Persicaria posumbu* (Buch.-Ham. ex D. Don) H. Gross in North America (Costea *et al.*, 2005), or as *P. caespitosum* Blume by many others. It is native to China, India, Indonesia, Japan, Korea, Myanmar, Nepal, the Philippines, Thailand, and was introduced around

1910 from Philadelphia into North America, USA: AL, CT, DC, DE, FL, GA, IA, IL, IN, KS, KY, LA, MA, MD, ME, MI, MN, MO, MS, NC, NE, NJ, NM, NY, OH, PA, SC, TN, TX, VA, VT, WI, WV; and Canada: BC, NB, ON, QC, SK.

19. *Pueraria montana* (Lour.) Merr. in Trans. Amer. Philos. Soc., n. s. 24 (2): 210, 1935. Syn. *Pueraria montana* (Lour.) Merr. var. *lobata* (Willd.) Maesen & S. Almeida in J. Bombay Nat. Hist. Soc. 85 (1): 234, 1988 (Fabaceae).

The Kudzu has a very complex nomenclature: it was first widely used as *P. thunbergiana* (Piper, 1920; Alexander, 1930; Hutchinson and Dalziel, 1958). Van der Maesen (1985) revised the genus, but wrongly recognized as *P. lobata* var. *montana*, which was followed by many works both locally and worldwide, even in recent years (Sun *et al.*, 2005). A few years later, the right name for Kudzu was corrected as *P. montana* (Lour.) Merr. var. *lobata* (van der Maesen and Almeida, 1988), but this correction was published in a local journal given little attention by the world until another report in the USA ten years later (Ward, 1998). Today, it is still confused; for example, a recent industrial monograph of *Pueraria* (van der Maesen, 2002) treated Kudzu as *P. montana* var. *lobata* in the botanical section, but some other chapters in the same work used only *P. lobata* var. *montana*. More confusing is that the three varieties (var. *thomsonii*, var. *lobata* and *montana*) named by van der Maessen are very difficult to identify in the field (Li Zhenyu, PE, personal comm. of March 13, 2006). In fact, it was not accepted by recent work (Cullen, 1995; Cox, 1999; Cronk and Fuller, 2001). There are more names for Kudzu if one checks modern literatures. For example, it is represented by 182 *P. lobata*, 42 *P. thunbergiana*, 13 *P. montana*, 6 *P. thomsonii*, and 3 *P. hirsuta* under Kudzu in BIOSIS (1980 - 2005).

Kudzu is an important traditional medicine in East Asia (Gu *et al.*, 1996; Zhou *et al.*, 1995; Zhu *et al.*, 2002) as well as a kind of food in east, south and southeast Asia (Alexander, 1930). It was introduced in 1876 at the Centennial Exposition in Philadelphia as an ornamental plant from Japan (Bell and Wilson, 1989), developed for use as forage in the 1920s in Florida (Piper, 1920), and promoted in the 1930s by the United States Soil Conservation Service for erosion control (Alexander, 1930; van der Maesen, 1985; Bell and Wilson, 1989) as well as for various uses in other fields (Piper, 1920; Porterfield, 1938). Kudzu coverage was 3 millions hm² in the 1950s in USA, but due to its rapid growth, it smothered trees, houses and telephone poles, and further use as cover was discouraged. It was declared a weed in 1972 (van der Maesen, 1985), and in 1981 it was about to be declared a pest because it affected c. 22000 hm² of commercial forest in the state of Georgia (van der Maesen, 2002; Shores, 1997). As large impenetrable masses, growing over woody vegetation and able to completely engulf unwooded areas (Godfrey, 1988), Kudzu can completely envelop a tree, killing it by shutting out all light (Bell and Wilson, 1989); it is a serious and widespread invader of semi-natural or natural habitat (Cronk and Fuller, 2001) across the USA (McClain *et al.*, 2006). The lesson of Kudzu, deliberately introduced as a useful plant with encouragement in the beginning and resulting in being forbidden as a notorious invasive in the end is an excellent case in invasive biology (Lowney, 2002).

Kudzu is native to east, south to southeast Asia, including Bhutan (Grierson and Long, 1987), China (Wu, 1994; 1995) (except Heilongjiang, Nei Menggu and Qinghai), northeast India (Naithani, 1990; Kumar and Sane, 2003), northern Thailand (Lock and Heald, 1994), Malaysia (Lock and Ford, 2004), and the Pacific

ic Islands (Cronk and Fuller, 2001). However, due to longtime cultivation, the origin is uncertain, but thought to be around warm tropical Asia, including south China, northeast India, Burma and Thailand. From there, it was introduced into Africa (Hutchinson and Dalziel, 1958; Burrows, 1989), Asia, Europe (Kartmazova *et al.*, 1981; Yakovlev *et al.*, 1996), Oceania and America (D Arcy, 1980; Bodner and Hymowitz, 2002). USA: AL, AR, CT, DC, DE, FL, GA, HI, IL, IN, KS, KY, LA, MA, MD, MO, MS, NC, NJ, NY, OH, OK, PA, SC, TN, TX, VA, WV.

20. *Reynoutria japonica* Houtt. Nat. Hist. 2 (8): 640, T. 51, f. 1, 1777; A. J. Li in Fl. Reip. Pop. Sin. 25 (1): 105 - 106, 1998, A. R. Li & C. W. Park in Fl. Chin. 5 : 319, 2003. Syn. *Polygonum cuspidatum* Sieb. & Zucc. in Abh. Math. -Phys. Cl. Konigl. Bayer. Akad. Wiss. 4 (3) : 208. 1846; *Fallopia japonica* (Houtt.) Ronse Decraene in Bot. J. Linn. Soc. 98 (4) : 369, 1988 (Polygonaceae).

Japanese Knotweed is native to China, Japan, Korea, far east Russia, widely cultivated and becoming a weed in other parts of the world. It was probably introduced to North America in the late 1800s as an ornamental plant and has also been used for erosion control and for landscape screening. It has spread through north and central Europe, Australia, New Zealand and most parts of USA: AK, AR, CA, CO, CT, DC, DE, GA, IA, ID, IL, IN, KS, KY, LA, MA, MD, ME, MI, MN, MO, MS, MT, NC, NE, NH, NJ, NY, OH, OR, OK, PA, RI, SC, TN, UT, VA, VT, WA, WI, WV; and Canada: BC, MB, NB, NF, NS, ON, PE, QC.

There are three genera and two epithets for Japanese Knotweed in the nomenclatural field: the plant material brought to Britain from Japan was named *Reynoutria japonica* Houtt (1777),

and additional plant material was named *Polygonum cuspidatum* Sieb. & Zucc. (1846). Later it was discovered the two plants were identical (Makino, 1901; Beerling *et al.*, 1994). However, modern research work suggest this species should be put in the genus *Fallopia* (Ronse Decraene and Akeroyd, 1988; Bailey and Stace, 1992; Beerling *et al.*, 1994; 1995; Seiger, 1996; Cox, 1999), but has never been accepted by local floras in east Asia; it is also known as *Polygonum cuspidatum* in Europe and North America (Akeroyd, 1989; Sorrie, 2005).

21. *Sapium sebiferum* (L.) Roxb., Hort. Bengal. 69, 1814, Annot. in Fl. Ind. ed. 2, 3 : 693, 1832; Syn. *Triadica sebifera* (L.) Small, Florida Trees 59, 1913; *Croton sebiferum* L., Sp. Pl. 2: 1004, 1753 (Euphorbiaceae).

Chinese tallow tree, as a species of *Sapium* (s. l.), a genus with worldwide tropical distribution, centered mainly in the New World (Tseng, 1997), has been widely accepted in the world (Jubinsky, 1996; Cronk and Fuller, 2001; Rogers and Evan, 2002; Rogers, 2003; Wall and Darwin, 1999). Recently, however, it has been assigned to *Triadica* (Esser, 2002) again, a separate genus from *Sapium* but only distributed in Asia. It was introduced from China to South Carolina in the 1700s and then in significant numbers to the Gulf Coast in the early 1900s for seed oil under recommendation from the USDA from the 1920s to 1940s. It is also ornamental (still sold and planted), with waxy seeds traditionally used to make candles. It is native to the south of Yellow River in China, Japan, Vietnam and India, and is also cultivated in Europe and Africa and in USA: AL, AR, FL, GA, LA, MS, NC, SC, TX.

22. *Setaria glauca* (L.) Beauv. Ess. Agrost. 51 : 178, 1812; G. Y. Sheng in Fl. Reip. Pop. Sin. 10 (1) : 357, 1990. Syn. *S. pumila* (Poir.)

Roemer & J. A. Schultes, Syst. Veget. 2 : 891, 1817; *Panicum glaucum* L., Sp. Pl. 3d. 1 : 56, 1753; *P. pumilum* Poir. in Encycl. Méth. Bot. 4 : 273, 1816 (Poaceae).

The Yellow Bristle Grass, from the Old World tropics and Eurasian temperates, was introduced into Australia and North America and is frequently used as *S. pumila* in Europe (Grass Team, Eur. Gard. Fl. 2 : 42, 1984) and North America (Rominger, 1962). It is distributed in USA: DC, DE, KY, LA, OR, WV; and Canada: ON, SK.

23. *Sparganium erectum* L., Pl. Sp. 971, 1753 (Sparganiaceae).

The Exotic Bur Reed, from Eurasia and North America (Cook and Nicolls, 1987), is a widespread aquatic plant, divided into *S. minimum* Wall. and *S. angustifolium* Michx. (Chen, 1992; Cook, 1980; Synott, 1984) in Eurasia, and *S. eurycarpum* Engelm (Kaul, 2000; Crow and Hellquist, 2000) in North America, also distributed in USA: CA, CT, IA, ID, IL, IN, MA, MD, ME, MI, MN, NC, ND, NE, NH, NJ, NY, OH, OR, PA, SD, VA, VT, WA, WI, WV, WY; and Canada: AB, NB, NF, NS, ON, PE, QC, SK.

Reference :

- Akeroyd JR, 1989. *Polygonum* [A]. In European Garden Flora [M]. London: Cambridge University Press, 3 : 128
- Alexander EJ, 1930. *Pueraria thunbergiana* [J]. *Addisonia*, 15 : 57—58, pl. 509
- Antenen, S. 1996. *Ampelopsis* [A]. In: Randall JM, Marinelli J eds., Invasive Plants: Weeds of the Global Garden, 21st Century Gardening Series, Handbook # 149 [M], Brooklyn: Brooklyn Botanic Garden
- Awmack CS, Lock JM, 2002. The genus *Alhagi* (Leguminosae: Papilionoideae) in the Middle East [J]. *Kew Bulletin*, 57 : 435—443
- Bailey JP, Stace CA, 1992. Chromosome number, morphology, pairing, and DNA values of species and hybrids in the genus *Fallopia* (Polygonaceae) [J]. *Plant Systematics and Evolution*, 180 : 29—52

- Ball PW, 1968. *Alhagi* [A]. In *Flora Europaea* [M]. London: Cambridge University Press, 2 : 191
- Ball PW, Akeroyd JR, 1993. *Bassia* [A]. In *Flora Europaea* [M]. London: Cambridge University Press, 2, 1 : 119
- Beerling DJ, Bailey JP, Conolly AP, 1994. *Fallopia japonica* (Houtt.) Ronse Decraene [J]. *Journal of Ecology*, 82 : 959—979
- Beerling DJ, Huntley B, Bailey JP, 1995, Climate and the distribution of *Fallopia japonica*: Use of an introduced species to test the predictive capacity of response surfaces [J]. *Journal of Vegetation Science*, 6 (2) : 269—282
- Bell CR, Wilson CR, 1989, "Kudzu," The Encyclopedia of Southern Culture [M]. Chapel Hill: University of North Carolina Press, 383—384
- Bodner CC, Hymowitz T, 2002. Ethnobotany of *Pueraria* species [A]. In: Keung WM ed., *Pueraria, the Genus Pueraria* [M]. London and New York: Taylor & Francis, 29—58
- Bossard CC, Randall JM, Hosbovsky MC, 2000, Invasive Plants of Californian Wildlands [M], Berkeley: University of California Press
- Burrows JE, 1989. Kudzu vine—a new plant invader of South Africa [J]. *Vald & Flora*, 75 (4) : 116—117
- Chen YD, 1992, Sparganiaceae [A]. In *Flora Reipublicae Popularis Sinicae* [M]. Beijing: Sciences Press, 8 : 25—33 (Chinese)
- Chen YL, 1982. *Rhamnus* [A]. In *Flora Reipublicae Popularis Sinicae* [M]. Beijing: Science Press, 48 (1) : 23 (Chinese)
- Ching RC, 1959. *Pteridophyta* [A]. In *Flora Reipublicae Popularis Sinicae* [M]. Beijing: Sciences Press, 2 : 317 (Chinese)
- Clemants SE, Mosyakin SL, 2003, Chenopodiaceae [A]. In *Flora of North America: north of Mexico* [M]. New York and Oxford: Oxford University Press, 4 : 296
- Costea M, Tardifm FJ, Hinds HR, 2005, Polygonaceae [A]. In *Flora of North America: north of Mexico* [M]. New York and Oxford: Oxford University Press, 5 : 592
- Cook CD K, 1980. *Sparganium* [A]. In *Flora Europaea* [M]. London: Cambridge University Press, 5 : 274—275
- Cook CD K, Nicolls MS, 1987. A monographic study of the genus *Sparganium* (Sparganiaceae) Part. 2. Subgenus *Sparganium* [J]. *Botanica Helvetica*, 91 (1) : 1—44
- Cox GW, 1999. Alien Species in North America and Hawaii [M], Washington DC and Covelo : Island Press, California
- Cronk QCB, Fuller JL, 2001. Plant Invaders [R]. Earthscan, London and Sterling, VA
- Crow GE, Hellquist CB, 2000. Aquatic and Wetland Plants of Northeastern North America [M]. Madison: The University of Wisconsin Press, 2 : 82—84 & 296
- Cullen J, 1995. *Pueraria* [A]. In *European Garden Flora* [M]. London: Cambridge University Press, 4 : 498
- D'Arcy WG, 1980. Flora of Panama, Family 83, Leguminosae [J]. *Annals of the Missouri Botanical Garden*, 67 (3) : 751—753
- Esser, HJ, 2002. A revision of *Triadica* Lour. (Euphorbiaceae) [J]. *Harvard Papers in Botany*, 7 (1) : 17—21
- Fernandes R, 1972. *Buglossoides* [A]. In *Flora Europaea* [M]. London: Cambridge University Press, 3 : 87—88
- Frappier B, Eckert RT, 2003. Potential impacts of the invasive exotic shrub *Rhamnus frangula* L. (glossy buckthorn) on forests of southern New Hampshire [J]. *Northeastern Naturalist*, 10 (3) : 277—296
- Godfrey RK, 1988. Trees, Shrubs, and Woody Vines of Northern Florida and Adjacent Georgia and Alabama [M], Athens: University of Georgia Press
- Grierson AJC, Long DG, 1987. Leguminosae [A]. In *Flora of Bhutan* [M]. Edinburgh: Royal Botanical Garden, 1 (3) : 692
- Gu ZP, Chen BZ, Feng RZ et al., 1996. The source utilization and evaluation of medical Kudzu and roots from the Genus plants *Pueraria* DC in China [J]. *Acta Pharmacologica Sinica*, 31 (5) : 387—393 (Chinese)
- Hsue HH, 1984. *Firmiana* [A]. In *Flora Reipublicae Popularis Sinicae* [M]. Beijing: Science Press, 49 (2) : 134 (Chinese)
- Hutchinson J, Dalziel JM, 1958. Leguminosae [A]. In *Flora of West Tropical Africa* [M]. London: Crown Agents for Oversea Governments and Administrations, 1 (2) : 573
- Jubinsky, G. 1996, *Sampium* [A]. In: Randall JM, Marinelli J eds., *Invasive Plants: Weeds of the Global Garden*, 21st Century Gardening Series, Handbook # 149 [M]. Brooklyn: Brooklyn Botanic Garden
- Kartmazova LS, Tkachenko NM, Borysov MI et al., 1981. Morphology anatomical diagnosis of the racemes of *Pueraria lobata* [J]. *Farmats Zhurnal* (Kiev), 1981 (1) : 56—59 (Ukrainian)
- Kaul RB, 2000, Sparganiaceae [A]. In *Flora of North America: north of Mexico* [M]. New York and Oxford: Oxford University Press, 22 : 270—277
- King CJ, 1986, *Dioscorea* [A]. In *European Garden Flora* [M]. London: Cambridge University Press, 1 : 330
- Krock SL, Williams CE, 2002. Allelopathic potential of the alien shrub glossy buckthorn, *Rhamnus frangula* L.: a laboratory bioassay [J]. *Journal of Pennsylvania Academy of Science*, 76 (1) : 17—21
- Kumar S, Sane PV, 2003. Legumes of South Asia—a check list [M]. Kew: Royal Botanic Gardens, 363
- Kuo PC, Tsui NR, 1987. *Elytrigia* [A]. In *Flora Reipublicae Popularis Sinicae* [M]. Beijing: Science Press, 9 (3) : 105 (Chinese)
- Lamont EE, Fitzgerald JM, 2001. Noteworthy plants reported

- from the Torrey Range-2000 [J]. *Journal of the Torrey Botanical Society*, 128 (4) : 409—414
- Lawatree A, 1980. *Spiroleda* [A]. In *Flora Europaea* [M]. London: Cambridge University Press, 5 : 273
- Li CL, 1996. New Taxa in Vitaceae from China [J]. *Chinese Journal of Applied and Environmental Biology*, 2 (1) : 43—53 (Chinese)
- Li CL, 1999. Vitaceae [A]. In *Flora Reipublicae Popularis Sinicae* [M]. Beijing : Sciences Press, 48 (2) : 32—53 (Chinese)
- Li JY, Chen YA, 1995. *Lespedeza* [A]. In *Flora Reipublicae Popularis Sinicae* [M]. Beijing : Science Press, 41 : 138 (Chinese)
- Lock JM, Ford C, 2004. Legumes of Malesia—a checklist [M]. Kew: Royal Botanic Gardens, 212
- Lock JM, Heald J, 1994. Legumes of Indo-China—a checklist [M]. Kew: Royal Botanic Gardens, 117
- Lowney KS, 2002. Friend or foe? Changing cultural definitions of Kudzu [A]. In : Keung WM ed., *Pueraria*, The genus *Pueraria* [M]. London and New York. : Taylor & Francis, 273—286
- van der Maesen LJG, 1985. Revision of the genus *Pueraria* DC with some notes on Teyleria Backer (Leguminosae) [J]. *Agricultural University Wageningen Papers*, 85 : 1—432
- van der Maesen LJG, Almeida SM, 1988. Two corrections to the nomenclature in the revision of *Pueraria* DC [J]. *Journal of the Bombay Natural History Society*, 85 (1) : 233—234
- van der Maesen LJG, 2002. *Pueraria*: Botanical characteristics in W. M. Keung (ed.), *Pueraria*, The genus *Pueraria* [M]. London and New York: Taylor & Francis, 1—28
- Makino T, 1901. *Polygonum reynoutria* (Houtt.) Makino [J]. *The Botanical Magazine* (Tokyo), 15 : 84
- McClain B, 1996. *Rhamnus* in Randall J. M. & J. Marinelli (eds.), *Invasive Plants: Weeds of the Global Garden*, 21st Century Gardening Series, Handbook # 149 [M]. Brooklyn: Brooklyn Botanic Garden
- McClain W, 2006. Distribution and reproductive potential of Kudzu (*Pueraria lobata*, Fabaceae) in Illinois, USA [J]. *Transactions of the Illinois State Academy of Science*, 99 (1/2) : 17—30
- McKean DR, 1997. *Frangula* [A]. In *European Garden Flora* [M]. London: Cambridge University Press, 5 : 192
- Melderis A, 1980. *Elymus* [A]. In *Flora Europaea* [M]. London: Cambridge University Press, 5 : 196
- Momiyama Y, 1977. On the correct name for a common Japanese species of *Ampelopsis* [J]. *The Journal of Japanese Botany*, 52 : 30
- Moore G, Glenn SD, Ma JS, 2009. Distribution of the native *Aralia spinosa* and non-native *Aralia elata* (Araliaceae) in the Northeastern United States [J]. *Rhodora*, 111 (946) : 145—154
- Mosyakin SL, 2003. *Kochia* [A]. In *Flora of North America: north of Mexico* [M]. New York and London: Oxford University Press, 4 : 310—311
- Mueller TC, Robinson DK, Beeler JE et al., 2003. *Dioscorea oppositifolia* L. phenotypic evaluations and comparison of control strategies [J]. *Weed Technology*, 17 (4) : 705—710
- Naithani HB, 1990. Flowering Plants of India, Nepal and Bhutan, Not Included in Sir. J. D. Hooker's Flora of British India [M]. Dehra Dun: Surya Publications, 124
- Nelson, E. C. 1989. *Humulus* [A]. In *European Garden Flora* [M]. London: Cambridge University Press, 3 : 101
- Ohba H, 1999. Vitaceae [A]. In *Flora of Japan* [M]. Tokyo: Kodansha, 2c : 121—122
- Page CN, Bennell FM, 1986. *Nephrolepis* [A]. In *European Garden Flora* [M]. London: Cambridge University Press, 1 : 61
- Piper CV, 1920. *Kudzu* USDA Department Circular, 89 : 1—7
- Porterfield WM, 1938. Ko, the Kudzu Vine, provides food, shade, clothes and medicine [J]. *Journal of the New York Botanical Garden*, 39 : 203
- Richardson JE, 1997. *Ampelopsis* [A]. In *European Garden Flora* [M]. London: Cambridge University Press, 5 : 201
- Rogers WE, 2003. Effects of simulated herbivory and resources on Chinese tallow tree (*Sapium sebiferum*, Euphorbiaceae) invasion of native coastal prairie [J]. *American Journal of Botany*, 90 (2) : 243—249
- Rogers WE, Evan S, 2002. Effects of simulated herbivory and resource availability on native and invasive exotic tree seedlings [J]. *Basic and Applied Ecology*, 3 (4) : 297—307
- Rominger JM, 1962. Taxonomy of *Setaria* (Gramineae) in North America. *Illinois Biological Monographs*, 29 : 1—108
- Ronse Decraene LP, Akeroyd JR, 1988. Generic limits in *Polygonum* and related genera (Polygonaceae) on the basis of floral characters [J]. *Botanical Journal of the Linnean Society*, 98 : 321—371
- Seiger LA, 1996. *Fallopia* [A]. In: Randall JM, Marinelli J (eds.), *Invasive Plants: Weeds of the Global Garden*, 21st Century Gardening Series, Handbook # 149 [M]. Brooklyn: Brooklyn Botanic Garden
- Shores M, 1997. The amazing story of Kudzu. University of Alabama Center for Public TV and Radio (<http://www.cptr.ua.edu/kudzu.htm>)
- Small E, 1978. A numerical and nomenclatural analysis of morpho-geographic taxa of *Humulus* [J]. *Systematic Botany*, 3 (1) : 68
- Sorrie BA, 2005. Alien vascular plants in Massachusetts [J]. *Rhodora*, 107 : 284—293
- Sun JH, Li ZC, Jewett DK et al., 2005. Genetic diversity of

- Pueraria lobata (kudzu) and closely related taxa as revealed by inter-simple sequence repeat analysis [J]. *Weed Research*, 45 : 255—260
- Synnott DM, 1984. *Sparganium* [A]. In European Garden Flora [M]. London: Cambridge University Press, 2 : 113
- Ting CT, Gilbert M, 2000. *Dioscorea* [A]. In Flora of China [M]. Beijing: Science Press, St. Louis: Missouri Botanical Garden, 24 : 276—296
- Tseng YC, 1997. *Sapium* [A]. In: Flora Reipublicae Popularis Sinicae [M]. Beijing : Science Press, 44 (3) : 12—23 (Chinese)
- Tutin TG, 1968. *Frangula* [A]. In Flora Europaea [M]. London. : Cambridge University Press, 2 : 245
- Wall DP, Darwin SP, 1999. Vegetation and elevational gradients within a bottomland hardwood forest of southeastern Louisiana [J]. *American Midland Naturalist*, 142 (1) : 17—30
- Wang WT, 1990. *Lithospermum* [A]. In Flora Reipublicae Popularis Sinicae [M]. Beijing: Science Press, 64 (2) : 36—38 (Chinese)
- Ward DB, 1998. *Pueraria montana*: The correct Scientific Name of the Kudzu [J]. *Castanea* , 63 (1) : 76—77
- Wen J, 2000. Revision of some problematic taxa of *Aralia* L. (Araliaceae) from China [J]. *Acta Phytotaxonomica Sinica* , 38 (1) : 1—9 (Chinese)
- Wu SH, 1999 Nephrolepidaceae [A]. In Flora Reipublicae Popularis Sinicae [M]. Beijing: Sciences Press, 6 (1) : 143—150 (Chinese)
- Wu TL, 1994. A study of Chinese *Pueraria* [J]. *Journal of Tropical and Subtropical Botany* , 2 (3) : 12—21 (Chinese)
- Wu TL, 1995. *Pueraria* [A]. In Flora Reipublicae Popularis Sinicae [M]. Beijing: Science Press, 41 : 219—230 (Chinese)
- nese)
- Xu LR, 1998. *Alhagi* [A]. In Flora Reipublicae Popularis Sinicae [M]. Beijing: Science Press , 42 (2) : 163 (Chinese)
- Yakovlev GP, 1979. Notes on the taxonomy of the genus *Alhagi* Gagnep. (Fabaceae) [J]. *Botanicheskii Zhurnal* , 64 (12) : 1794—1799 (Russian)
- Yakovlev GP, Sytin AK, Yu R et al. , 1996. Legumes of Northern Eurasia-A Checklist [M]. Kew: Royal Botanic Gardens , 449—450
- Zebryk TM, 2003. *Aralia spinosa*: An introduced woody species with invasive potential in Massachusetts [J]. *Rhodora* 105 , no. 924 , 395—398
- Zheng HY, Wu JQ, Ding D et al. , 2004. Invasive plants of Asian origin established in the United States and their natural enemies [R] , 1 : 1—147. USDA FS FHTET 2004 - 2005
- Zheng HY, Wu JQ, Ding D et al. , 2005. Invasive plants established in the United States that are found in Asia and their associated natural enemies [R] , 2 : 1—174. USDA FS FHTET 2005 - 15
- Zhou YP, Su XL, Cheng B et al. , 1995. Comparative study on pharmacological effects of various species of *Pueraria* [J]. *China Journal of Chinese Materia Medica* , 20 (10) : 619—621 (Chinese)
- Zhu GL, 1979 , Chenopodiaceae [A]. In Flora Reipublicae Popularis Sinicae [M]. Beijing : Sciences Press , 25 (2) : 98 (Chinese)
- Zhu GL, Riedl H, Kamelin RV, 1995. *Lithospermum* in Flora of China [M]. Beijing : Science Press , St. Louis: Missouri Botanical Garden , 16 : 343
- Zhu YP, Zhang HM, Zeng M, 2002. *Pueraria* (Ge) in traditional Chinese herbal Medicine [A]. In: Keung WM (ed.) , *Pueraria* , The genus *Pueraria* [M]. London and New York : Taylor & Francis , 59 —69