

## 毛梗希莶的化学成分\*

马云保 熊 江 许云龙<sup>+</sup>

(中国科学院昆明植物研究所植物化学开放实验室, 昆明 650204)

**摘要** 从毛梗希莶(*Siegesbeckia glabrescens*)的乙醇提取物中分到胡萝卜甙和3个二萜类成分, 根据光谱和化学证据, 3个二萜的化学结构被分别确定为: 对映-16 $\beta$ ,17-二羟基贝壳杉烷-19-酸(1), 腺梗希莶甙(2)和希莶甙(3)。对映二羟基-16 $\beta$ ,17-贝壳杉烷-酸和腺梗希莶甙系首次从毛梗希莶中到。

**关键词** 菊科, 毛梗希莶, 对映-16 $\beta$ ,17-二羟基贝壳杉烷-19-酸(1), 腺梗希莶甙(2), 希莶甙(3) **分类号** Q946.83 **Q949.783.5**

**The Constituents of Siegesbeckia glabrescens**MA Yun-Bao XIONG Jiang XU Yun-Long<sup>+</sup>

(Laboratory of Phytochemistry, Kunming Institute of Botany, The Chinese Academy of Sciences, Kunming 650204)

**Abstract** Three diterpenoids, compound A (1), B (2) and C (3), have been isolated together with daucosterol (4) from the ethanol extract of *Siegesbeckia glabrescens*. Their chemical structures have been elucidated as ent-16 $\beta$ ,17-dihydroxykauran-19-oic acid (1), siegesbeckioside (2), darutoside (3), on the basis of chemical and spectral evidences. Compounds 1 and 2 are isolated for the first time from *Siegesbeckia glabrescens*.

**Key words** Compositae, *Siegesbeckia glabrescens*, ent-16 $\beta$ ,17-dihydroxykauran-19-oic acid (1), Siegesbeckioside (2), Darutoside (3)

Plants of the genus *Siegesbeckia* are annual herbs widely distributed in tropical and temperate zones, and they have been used as a traditional medicine to treat rheumatic arthritis, hypertension, malaria, neurasthenia and snake-bite in China. Modern pharmacological experiments show that the extracts and constituents of *Siegesbeckia* exhibit analgesic, antiinflammatory(Yamatomo *et al.*, 1987), antihypertensive(Kim *et al.*, 1980), antioxidative(Su *et al.*, 1986), immuno-inhibitory, and infertile activities(Dong *et al.*, 1989; Ynag *et al.*, 1976). A series of ent-kaurane and ent-pimarane diterpenoids(Xiong *et al.*, 1992, 1997; Liu *et al.*, 1991; Kim *et al.*, 1979), sesquiterpene lactones, and flavonoids from *Siegesbeckia* have been reported(Zdero *et al.*, 1991). In our continuing search for biologically active constituents from *Siegesbeckia* plants, five new diterpenoids have been reported previously(Xiong *et al.*, 1992, 1997). The present paper describes the isolation, structural elucidation and identification of the other three diterpenoids from *Siegesbeckia glabrescens*.


\*Projects supported by the Natural Science Foundation of Yunnan

<sup>+</sup>Author to whom correspondence should be addressed

Received: 1997-04-21, Revised: 1997-06-12

## RESULTS AND DISCUSSION

**Compound A (1)**  $C_{20}H_{32}O_4$ , M 336, was obtained as colourless plates. Its IR spectrum revealed that hydroxyl ( $3420, 3250, 1027 \text{ cm}^{-1}$ ) and carboxyl ( $1690 \text{ cm}^{-1}$ ) were present as functional groups. 1 showed the presence of two methyl groups, ten methylene groups, three methine groups, four quaternary carbons, and one carboxyl group in the  $^{13}\text{C}$  NMR spectrum (Table 1). The above data and two tertiary methyl signals at  $\delta$ 1.19, 1.35 ppm and 5 unsaturation degrees suggested that 1 has a typical ent-kaurane nucleus as basic skeleton (Xiong *et al.*, 1992). In the  $^{13}\text{C}$  NMR spectrum of 1, two singlets ( $\delta$ 44.03, 180.19 ppm) and one quartet ( $\delta$ 29.43 ppm) are reasonably assigned to C-4, C-19 and C-18. The signals at (4.13 and 4.04 (each 1H, d, 10.8Hz) and at  $\delta$ 46.02 (d), 54.01 (t), 81.73 (s) and 66.53 (t), assigning to C-13, C-15, C-16 and C-17, indicated the presence of two-substituted 16 $\alpha$ ,17-glycol system. Therefore, the chemical structure of 1 can be represented as ent-16 $\beta$ ,17-dihydroxy-kauran-19-oic acid (1).



**Compound B (2)**  $C_{26}H_{44}O_8$ , M 484, was obtained as colourless needles. Its IR spectrum ( $3575, 3510, 3380, 1080, 1049, 1020 \text{ cm}^{-1}$ ) revealed the presence of hydroxyl groups. 2 showed the presence of two methyl groups, eleven methylene groups, three methine groups, four quaternary carbons, and one glucose moiety in the  $^{13}\text{C}$  NMR spectrum (Table 1). The above data and two tertiary methyl signals at (1.00, 0.79 ppm and 5 unsaturation degrees suggested that 2 has a typical ent-kaurane nucleus as basic skeleton (Xiong *et al.*, 1992). In the  $^{13}\text{C}$  NMR spectrum of 2, one singlet ( $\delta$ 37.65 ppm), one quartet ( $\delta$ 17.97 ppm) and one extreme downfield triplet

( $\delta$ 79.47 ppm) are reasonably assigned to C-4, C-19 and C-18. This suggestion is supported by the signals at 3.75, 3.42 (each 1H, d, 9.52 Hz). The signals at  $\delta$ 4.13, 4.04 (each 1H, d, 10.8 Hz) and at  $\delta$ 46.13 (d), 54.15 (t), 81.64 (s) and 66.52 (t), assigning to C-13, C-15, C-16 and C-17, indicated the presence of two-substituted 16 $\alpha$ ,17-glycol system. The signals at  $\delta$ 4.80 (1H, d, 7.76 Hz) and  $\delta$ 105.58 (d) were assignable to C-1 position of glucose, thus suggesting the  $\beta$ -configuration at the anomeric carbon of the glucoside. Other signals of 2 at  $\delta$ 4.61 (1H, dd, 11.64, 1.88 Hz), 4.46 (1H, dd, 11.64, 5.20 Hz), 4.30~4.23 (2H, m), 4.09~4.00 (2H, m) and at (75.26 (d), 78.53 (d), 71.86 (d), 78.67 (d), 62.96 (t) were in agreement with those of the  $\beta$ -D-glucoside. Furthermore, the signal assignable to C-18 ( $\delta$ 79.47 t) of 2 was unchanged in comparison

with that of the pentaacetate **2a**. Accordingly, the chemical structure of **2** can be determined as *ent*- $\alpha$ , $\beta$ , $17,18$ -trihydroxykauran- $18$ -O- $\beta$ -D-glucopyranoside, namely siegesbeckioside (**2**).

Table 1  $^{13}\text{C}$  NMR chemical shifts of **1**, **2**, **2a**, **3** and **3a** in  $\text{C}_5\text{D}_5\text{N}$ 

Carbon	<b>1</b>	<b>2</b>	<b>2a</b>	<b>3</b>	<b>3a</b>
1	41.18 t	40.00 t	40.00 t	37.14 t	36.75 t
2	19.92 t	18.36 t	18.18 t	24.17 t	23.87 t
3	38.86 t	36.42 t	36.01 t	85.32 d	86.17 d
4	44.03 s	37.65 s	37.36 s	38.77 s	38.73 s
5	57.18 d	49.28 d	49.66 d	55.08 d	54.99 d
6	23.03 t	20.74 t	20.87 t	22.73 t	22.67 t
7	42.88 t	42.02 t	42.11 t	36.46 t	36.27 t
8	45.06 s	44.88 s	44.95 s	138.39 s	140.76 s
9	56.46 d	56.92 d	57.20 d	50.90 d	50.71 d
10	41.15 s	39.45 s	39.41 s	38.21 s	38.36 s
11	19.07 t	18.73 t	18.30 t	18.87 t	18.86 t
12	26.84 t	26.89 t	26.76 t	32.93 t	32.74 t
13	46.02 d	46.13 d	46.57 d	38.10 s	37.43 s
14	37.87 t	37.90 t	37.70 t	129.53 d	126.93 d
15	54.01 t	54.15 t	54.07 t	76.82 d	74.97 d
16	81.73 s	81.64 s	78.99 s	64.04 t	64.11 t
17	66.53 t	66.52 t	69.32 t	23.40 q	23.50 q
18	29.43 q	79.47 t	79.26 t	29.02 q	28.82 q
19	180.19 s	17.97 q	17.73 q	14.97 q	14.84 q
20	16.12 q	18.51 q	18.30 q	17.29 q	16.85 q
Glc-1'		105.58 d	101.38 d	102.45 d	99.20 d
-2'		75.26 d	72.30 d	75.22 d	72.35 d
-3'		78.53 d	73.60 d	78.20 d	73.69 d
-4'		71.86 d	69.40 d	72.18 d	69.82 d
-5'		78.67 d	72.22 d	78.64 d	72.35 d
-6'		62.96 t	62.54 t	63.34 t	62.79 t
OAc		171.17 s			170.79 s
		170.48 s			170.71 s
		170.29 s			170.58 s
		169.81 s			170.48 s
		169.42 s			169.98 s
		20.87 q			169.59 s
		20.66 q			20.98 q
		20.58 q			20.76 q
		20.45 q			20.76 q
		20.45 q			20.76 q
					20.62 q
					20.62 q

**Compound C (3)**  $\text{C}_{26}\text{H}_{44}\text{O}_8$ , M 484; white amorphous powder. Its IR spectrum revealed that hydroxyl ( $3400\sim 3360, 1072, 1025, 1010 \text{ cm}^{-1}$ ) and double bond ( $1630 \text{ cm}^{-1}$ ) were present as functional groups. **3** showed the presence of four methyl groups, seven methylene groups, four methine groups, three quaternary carbons, two olefinic carbons and one glucose moiety in the  $^{13}\text{C}$  NMR spectrum (Table 1). The above data and four tertiary methyl signals at  $\delta$  1.19, 1.14, 0.88, 0.67 ppm and 5 unsaturation degrees suggested that **3** has a typical *ent-pimarane* nucleus as basic skeleton (Dong *et al.*, 1989). In the  $^{13}\text{C}$  NMR spectrum of **3**, one singlet ( $\delta$  38.77 ppm), one Extreme downfield doublet ( $\delta$  85.32 ppm), and two quartets ( $\delta$  29.02, 14.97 ppm) are reasonably assigned to C-4, C-3, C-18 and C-19. The glucose is linked to 3 $\alpha$  position based on the

above data and the signal at 3.52 (dd, 11.58, 3.50 Hz). The signals at  $\delta$ 38.10 (s), 76.82 (d), 64.04 (t) and 23.40 (q) assigning to C-13, C-15, C-16 and C-17, indicated the presence of one-substituted 15,16-glycol system. The signals at  $\delta$ 4.84 (1H, d, 7.68 Hz) and  $\delta$ 102.45 (d) were assignable to C-1 position of glucose, thus suggesting the  $\beta$ -configuration at the anomeric carbon of the glucoside. Other signals of 3 at  $\delta$ 4.51 (1H, dd, 9.84, 1.88 Hz), 4.51~4.33 (2H, m), 4.33 (1H, dd, 11.32, 5.24 Hz), 4.21~3.91 (2H, m) at  $\delta$ 75.22 (d), 78.20 (d), 72.18 (d), 78.64 (d), 63.34 (t) were in agreement with those of the  $\beta$ -D-glucoside. Furthermore, the signal assignable to C-3 ( $\delta$ 85.32, d) of 3 was unchanged in comparison with that of the hexaacetate **3a**. Accordingly, the chemical structure of 3 can be determined as *ent*- $\beta$ ,15,16-trihydroxypimaran-3-O- $\beta$ -D-glucopyranoside, namely, darutoside (3).

## EXPERIMENT

**General** Kofler melting points were uncorrected; Optical rotations were taken on a Jasco-20C digital polarimeter. IR were recorded on KBr discs with a Perkin-Elmer 577 spectrometer. UV were obtained in EtOH on a UV-210A spectrometer. EIMS (positive) were measured on a VG Auto Spec-3000 spectrometer with direct inlet 70 or 20 eV. NMR were run on a Brucker AM-400 spectrometer using TMS as internal standard; chemical shift values are reported in (ppm) units (pyridine-d5). Coupling constants ( $J$ ) were expressed in Hz.

**Plant Material** *Siegesbeckia glabrescens* was collected in Fumin County, Yunnan, China in Sept, 1992 and identified by Prof. Yanhui Li. A voucher specimen was deposited in the Herbarium of Kunming Institute of Botany, Academia Sinica.

**Extraction and isolation** Dried and powdered herbs (7.76 kg) were repeatedly soaked with warm EtOH for 2 days  $\times$  4 and then concd. to crude residue. The residue was suspended in H<sub>2</sub>O and shaken, in order, in EtOAc ( $\times$  3), and n-BuOH ( $\times$  4) saturated with H<sub>2</sub>O. The EtOAc soln was evapd in vacuum to obtain a residue (229 g) which was decoloured with activated charcoal in MeOH, filtered and evapd to yield 196 g brown syrup. The n-BuOH soln were also evapd in vacuum to yield 25 g yellow gums. The EtOAc fraction (166 g) was mixed with silica gel (180 g, 60~200 mesh) and subjected to CC over silica gel (1243 g, 200~300 mesh) eluting with CHCl<sub>3</sub> and increasing proportions of MeOH-CHCl<sub>3</sub> to obtain 1 (80 mg, 0.00103%), 4 (791 mg, 0.0102%), 3 (790 mg, 0.0102%), 2 (50 mg, 0.00064%). Some components were further purified by recrystallization and prep. TLC (silica gel).

**ent-16 $\beta$ ,17-Dihydroxykauran-19-oic acid (1)** C<sub>20</sub>H<sub>32</sub>O<sub>4</sub>, M 336; colourless plates (MeOH-CHCl<sub>3</sub>), mp. 266~268°C;  $[\alpha]_D^{25}$ -88° (c 0.651, C<sub>5</sub>H<sub>5</sub>N); no UV absorption; IR  $\nu_{max}^{KBr\text{cm}^{-1}}$ : 3420, 3250, 1690, 1225, 1027; EIMS (20eV) m/z (%): 318[M-H<sub>2</sub>O]<sup>+</sup>(23), 305[M-CH<sub>2</sub>OH]<sup>+</sup>(100), 287[M-H<sub>2</sub>O-CH<sub>2</sub>OH]<sup>+</sup>(25), 259[M-CH<sub>2</sub>O-H-HCOOH]<sup>+</sup>(50), 121(68), 109(72), 95(56), 81(56), 43(57); <sup>1</sup>H NMR (C<sub>5</sub>D<sub>5</sub>N): 4.13 and 4.04 (each 1H, ABd, J=10.8 Hz, 17-H<sub>2</sub>), 1.35 (3H, s, 18-Me), 1.19 (3H, s, 20-Me). The mp, mmp,  $[\alpha]_D$ , IR, and R<sub>f</sub> value (TLC) of 1 are in agreement with those of authentic sample[6]. <sup>13</sup>C NMR data see Table 1.

**siegesbeckioside (2)** C<sub>26</sub>H<sub>44</sub>O<sub>8</sub>, M 484; colourless needles (MeOH), mp. 276.5~277.5°C;  $[\alpha]_D^{25}$ -29.21° (c 0.290, C<sub>5</sub>H<sub>5</sub>N); no UV absorption; IR  $\nu_{max}^{KBr\text{cm}^{-1}}$ : 3575, 3510, 3380, 2930, 2920, 1465, 1440, 1380, 1190, 1163, 1080, 1049, 1020, 921, 875; EIMS (20eV) m/z (%): 484[M+ no appearance], 466[M-H<sub>2</sub>O]<sup>+</sup>, 453[M-CH<sub>2</sub>OH]<sup>+</sup>, 448[M-2H<sub>2</sub>O]<sup>+</sup>, 435[M-H<sub>2</sub>O-CH<sub>2</sub>OH]<sup>+</sup>, 430[M-3H<sub>2</sub>O]<sup>+</sup>, 417[M-CH<sub>2</sub>OH-2H<sub>2</sub>O]<sup>+</sup>, 412[M-4H<sub>2</sub>O]<sup>+</sup>, 405, 399[M-CH<sub>2</sub>OH-3H<sub>2</sub>O]<sup>+</sup>, 394[M-5H<sub>2</sub>O]<sup>+</sup>, 377[M-5H<sub>2</sub>O-OH]<sup>+</sup>, 333(2), 315(3),

304[M-Glucose]<sup>+(5)</sup>, 291(33), 287[304-OH]<sup>+(22)</sup>, 273[304-CH<sub>2</sub>OH]<sup>+(46)</sup>, 269[304-H<sub>2</sub>O-OH]<sup>+(13)</sup>, 255[304-CH<sub>2</sub>OH-H<sub>2</sub>O]<sup>+(11)</sup>, 229(2), 43(100); <sup>1</sup>H NMR (C<sub>5</sub>D<sub>5</sub>N): 4.80 (1H, d, J=7.76 Hz, Glc-1-H), 4.61 (1H, dd, J=11.64, 1.88 Hz, Glc-6-H), 4.46 (1H, dd, J=11.64, 5.20 Hz, Glc-6-H), 4.30-4.23 (2H, m, Glc-H<sub>2</sub>), 4.09-4.00 (2H, m, Glc-H<sub>2</sub>), 4.13, 4.04 (each 1H, ABd, J=10.8 Hz, 17-H<sub>2</sub>), 3.75, 3.42 (each 1H, ABd, J=9.52 Hz, 18-H<sub>2</sub>), 1.00 (3H, s, 19-Me), 0.79 (3H, s, 20-Me). The Above-mentioned data of 2 are in agreement with those of authentic sample[6]. <sup>13</sup>C NMR data see Table 1.

**pentaacetate of siegesbeckioside (2a)** C<sub>36</sub>H<sub>54</sub>O<sub>13</sub>, M 694; clubbed crystals (MeOH), mp. 171~172°C, [α]<sub>D</sub><sup>23</sup>-61.14° (c 0.240, CHCl<sub>3</sub>); no UV absorption; IR ν<sub>max</sub><sup>KBr</sup> cm<sup>-1</sup>: 3555, 3500, 1745, 1443, 1380, 1370, 1250, 1225, 1040, 905, 620; EIMS (20eV) m/z (%): 694[M<sup>+</sup>, no appearance], 677[M-OH]<sup>+(7)</sup>, 635[M-OAc]<sup>+(2)</sup>, 634[M-HOAc]<sup>+(3)</sup>, 621[M-CH<sub>2</sub>OAc]<sup>+(28)</sup>, 617[M-OAc-H<sub>2</sub>O]<sup>+(10)</sup>, 616[M-H<sub>2</sub>O-HOAc]<sup>+(25)</sup>, 579[621-Ketene]<sup>+(4)</sup>, 578[M-CH<sub>2</sub>OAc-CH<sub>3</sub>CO]<sup>+(12)</sup>, 556[M-H<sub>2</sub>O-2HOAc]<sup>+(5)</sup>, 514[556-Ketene]<sup>+(6)</sup>, 472[514-Ketene]<sup>+(4)</sup>, 412[472-HOAc]<sup>+(2)</sup>, 399[472-CH<sub>2</sub>OAc]<sup>+(6)</sup>, 346[M-Glc(Ac)<sub>4</sub>]<sup>+</sup>, 331[346-CH<sub>3</sub>]<sup>+</sup>, 315[346-CH<sub>2</sub>OH]<sup>+</sup>, 286[346-HOAc]<sup>+</sup>, 255[286-CH<sub>2</sub>OH]<sup>+(12)</sup>, 229(5), 169(30), 43(100); <sup>1</sup>H NMR (C<sub>5</sub>D<sub>5</sub>N): 5.71 (1H, t, J=9.56 Hz, Glc-H), 5.49, 5.44 (each 1H, ABd, J=9.88 Hz, Glc-6-H<sub>2</sub>), 4.84 (1H, d, J=7.92 Hz, Glc-1-H), 4.88, 4.61 (each 1H, ABd, J=11.16 Hz, 17-H<sub>2</sub>), 4.64~4.40 (2H, m, Glc-H<sub>2</sub>), 4.12 (1H, dd, J=8.22, 2.92 Hz, Glc-H), 3.57, 3.31 (each 1H, ABd, J=9.22 Hz, 18-H<sub>2</sub>), 2.14 (3H, s, OAc), 2.06 (6H, s, 2×OAc), 2.02 (3H, s, OAc), 2.01 (3H, s, OAc), 0.99 (3H, s, 19-Me), 0.77 (3H, s, 20-Me). The mp, mmp, [α]<sub>D</sub>, IR, and R<sub>f</sub>value (TLC) of 2a are in agreement with those of authentic sample(Xiong et al, 1989). <sup>13</sup>C NMR data see Table 1.

**darutoside (3)** C<sub>26</sub>H<sub>44</sub>O<sub>8</sub>, M 484; white amorphous powder (MeOH-CHCl<sub>3</sub>), mp. 234~237°C; no UV absorption; IR ν<sub>max</sub><sup>KBr</sup> cm<sup>-1</sup>: 3400~3360, 2940, 2870, 2850, 1630, 1450, 1375, 1160, 1072, 1025, 1010, 880; EIMS (70eV) m/z (%): 484[M<sup>+</sup>, no appearance], 440, 423[M-CH(OH)CH<sub>2</sub>OH]<sup>+(2)</sup>, 346(100), 331, 316(13), 301(18), 271(16), 243[M-Glucose-CH(OH)CH<sub>2</sub>OH]<sup>+(9)</sup>, 229(22), 217, 205(14), 189, 128, 115, 105, 91, 77, 43(39); <sup>1</sup>H NMR (C<sub>5</sub>D<sub>5</sub>N): 5.39 (1H, br s, 14-H), 4.84 (1H, d, J=7.68 Hz, Glc-1-H), 4.51 (1H, dd, J=9.84, 1.88 Hz, Glc-H), 4.51~4.33 (5H, m, 15-H, 16-H<sub>2</sub>, Glc-H<sub>2</sub>), 4.33 (1H, dd, J=11.32, 5.24 Hz, Glc-H), 4.21~3.91 (2H, m, Glc-H<sub>2</sub>), 3.52 (1H, dd, J=11.58, 3.50 Hz, 3-H), 1.19 (3H, s, 17-Me), 1.14 (3H, s, 18-Me), 0.88 (3H, s, 20-Me), 0.67 (3H, s, 19-Me). The Above-mentioned data of 3 are in agreement with those of authentic sample(Dong et al, 1989). <sup>13</sup>C NMR data see Table 1.

**hexaacetate of darutoside (3a)** C<sub>38</sub>H<sub>56</sub>O<sub>14</sub>, M 736; colourless needles (MeOH), mp. 88.5~89.5°C; no UV absorption; IR ν<sub>max</sub><sup>KBr</sup> cm<sup>-1</sup>: 1750, 1640, 1370, 1250, 1225, 1090, 1040, 910, 870, 755, 630, 605; <sup>1</sup>H NMR (C<sub>5</sub>D<sub>5</sub>N): 5.72 (1H, t, J=9.54 Hz, Glc-H), 5.47~5.35 (3H, m, 16-H, Glc-6-H<sub>2</sub>), 5.27 (1H, br s, 14-H), 4.93 (1H, d, J=7.96 Hz, Glc-1-H), 4.63~4.40 (3H, m, 16-H', Glc-H<sub>2</sub>), 4.30 (1H, dd, J=11.62, 9.14 Hz, 15-H), 4.09 (1H, dd, J=9.94, 4.76 Hz, Glc-H), 3.40 (1H, dd, J=11.70, 3.86 Hz, 3-H), 2.13 (6H, s, 2×OAc), 2.04 (3H, s, OAc), 2.03 (3H, s, OAc), 2.00 (3H, s, OAc), 1.98 (3H, s, OAc), 1.13 (3H, s, 17-Me), 1.03 (3H, s, 18-Me), 0.85 (3H, s, 20-Me), 0.84 (3H, s, 19-Me). <sup>13</sup>C NMR data see Table 1.

**daucosterol (4)** C<sub>35</sub>H<sub>60</sub>O<sub>6</sub>; white amorphous powder (MeOH-CHCl<sub>3</sub>), mp. 276°C (dec.); no UV absorption; IR ν<sub>max</sub><sup>KBr</sup> cm<sup>-1</sup>: 3450~3360, 2930, 2867, 1457, 1435, 1374, 1362, 1162, 1104, 1070, 1020; The mp, mmp, [α]<sub>D</sub>, IR, and R<sub>f</sub>value (TLC) of 4 are in agreement with those of authentic sample(Xiong et al, 1992).

## REFERENCES

Dong X Y, Chen M, Jin W et al, 1989. Studies on antifertility constituents of Siegesbeckia glabrescens Mak. *Acta*

*Pharmaceutica Sinica*, 24 (11), 833~836

Kim J H, Han K D, Kazuo Yamasaki et al, 1979. Darutoside, A aiterpenoid from *Siegesbeckia pubescens* and its structure revision. *Phytochemistry*, 18 (5), 894~895

Kim J H, Yu J C, Chang I M et al, 1980. Antihypertensive activities of diterpenoid (16,17-dihydroxy-16-(—)-kaurane-19-oic acid) from *Siegesbeckia pubescens* against okamoto-spontaneously hypertensive rats. *Soul Taehakkyo Saengyak Yonguso Ojukjip*, 19, 73~77

Liu K, Roder E. 1991. Diterpenes from *Siegesbeckia glabrescens*. *Planta Med*, 57 (4), 395~396

Su J D, Osawa T, Namiki M. Screening for antioxidative activity of crude drugs. *Agric Biol Chem*, 50 (1), 199~203

Xiong J, Ma Y B, Xu Y L, 1992. Diterpenoids from *Siegesbeckia pubescens*. *Phytochemistry*, 31(3), 917~921

Xiong J, Ma Y B, Xu Y L, 1997. The constituents of *Siegesbeckia orientalis*. *Natural Product Sciences*, 3(1), 14~18

Yamamoto F, Yokota M, Kosuge T, 1987. New sesquiterpenes from *Siegesbeckia pubescens* as analgesics and antiinflammatories. *Jpn Kokai Tokkyo Koho*, 175, 476 [87,175,476]

Yang TH, Liu S C, Chang S S et al, 1976. Studies on the constituents of *Siegesbeckia orientalis*. *Proc Natl Sci Coun*, Part 2, 9: 149~154

Zdero C, Bohlmann F, King R M et al, 1991. Sesquiterpene lactones and other constituents from *Siegesbeckia orientalis* and *Guizotia scabra*. *Phytochemistry*, 30 (5), 1579~1584

\* \* \* \* \*

云 南 植 物 研 究 1998, 20(2): 238~240  
Acta Botanica Yunnanica

(20)

## 238 蒙绣菊素的结构

谢海辉 魏孝义 韦璧瑜

(中国科学院华南植物研究所, 广州 510650)

### The Structure of Spiaramongolin

XIE Hai-Hui WEI Xiao-Yi WEI Bi-Yu

South China Institute of Botany, The Chinese Academy of Sciences, Guangzhou 510650

Key words *Spiraea*, *S. mongolica*, Spiaramongolin

关键词 绣线菊属, 蒙古绣线菊, 蒙绣菊素 化学结构 蔷薇科

分类号 Q946

R284.1 Q 949.751·8

蒙古绣线菊 (*Spiraea mongolica* Maxim.) 系蔷薇科绣线菊属植物, 分布于我国西北地区, 是一种民间药。性温, 味微甘, 有生津止渴、治腹水的功效 (青海省生物研究所等, 1972)。化学成分未见报道。为探讨其入药的化学基础, 我们研究了它的化学成分, 从中分离出  $\beta$ -谷甾醇、白桦酯醇、白桦脂酸、白桦脂酸-3, 5-二羟基肉桂酸酯、胡萝卜甙 (谢海辉等, 1994) 及一个具有三元结构的新化合物—蒙绣菊素 (V)。本文主要报道 V 的结构鉴定。

1997-08-14 收稿, 1998-01-21 接受发表