

中华青莢叶的一个新果糖酯^{*}

来国防^{1,2}, 赵沛基¹, 倪志伟¹, 许云龙¹,
王梅², 罗士德¹, 王易芬^{1**}

(1 中国科学院昆明植物研究所植物化学与西部植物资源持续利用国家重点实验室, 云南 昆明 650204;

2 云南省食品药品检验所, 云南 昆明 650011)

摘要: 从山茱萸科中华青莢叶 (*Helwingia chinensis*) 的乙醇提取物中分离得到一个新果糖酯和十个已知化合物。通过现代波谱技术分别鉴定为: 2-O-(E)-咖啡酰-3-O-(3, 5-二甲氧基香豆酰)-D-呋喃果糖甙 (1), 2-O-D-呋喃果糖基-D-异吡喃糖酯 (2), 甘草甜素 (3), 4-羟基-7-O-葡萄糖-2, 3-二羟黄酮甙 (4), 黄豆甙 (5), 5-葡萄糖芹菜甙 (6), 7-O-葡萄糖芹菜甙 (7), 4-O-葡萄糖香豆酸 (8), 葡萄糖咖啡酸 (9), 3-赤杨醇 (10), 薯蓣皂甙 3-O-{ -L-鼠李糖吡喃糖基 (1 2)-[-L-阿拉伯呋喃糖基 (1 3)]-D-葡萄糖吡喃糖} (11)。

关键词: 中华青莢叶; 山茱萸科; 呋喃果糖; 黄酮甙; 抗细菌活性

中图分类号: Q 946

文献标识码: A

文章编号: 0253- 2700 (2008) 01- 115- 06

A New Fructofuranoside from *Helwingia chinensis* (Cornaceae)

LAI Guo-Fang^{1,2}, ZHAO Pei-Ji¹, NI Zhi-Wei¹, XU Yun-Long¹,
WANG Mei², LUO Shi-De¹, WANG Yi-Fen^{1**}

(1 State Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany,

Chinese Academy of Sciences, Kunming 650204, China;

2 Yunnan Institute for Food and Drug Control, Kunming 650011, China)

Abstract: A new fructofuranoside, together with ten known compounds were isolated from the EtOH extraction of the aerial part of *Helwingia chinensis*. Their structures were elucidated to be 2-O-(E)-caffeyl-3-O-(3, 5-dimethoxylcoumaroyl)-D-fructofuranose (1), 2-O-D-fructofuranosyl-D-allopyranose (2), liquiritigenin (3), 4-hydroxy-2, 3-dihydro-flavonone-7-O-D-glucoside (4), daidzin (5), apigenin 5-O-glucopyranoside (6), apigenin-7-O-D-glucoside (7), coumaric acid 4-O-D-glucopyranoside (8), glucopyranosylcaffeic acid (9), 3-hydroxyglutin-5-ene (10), diosgenin 3-O-L-rhamnopyranosyl (1 2)-[-L-arabinofuranosyl (1 3)]-D-glucopyranoside (11) by intensive interpretation of spectral data.

Key words: *Helwingia chinensis*; Cornaceae; Fructofuranose; Flavonoids; Antibacterial activity

Helwingia chinensis Batal (Cornaceae) is distributed in the western and southern regions of China, for example, Gansu, Guizhou, Hubei, Hunan, Shanxi, Sichuan, Tibet and Yunnan (Song *et al.*, 1990). Its flowers and leaves were edible, and a decoction of the

leaves and the bark are ingested for treating skin inflammations (Zhonghua Bencao Bianweihui, 1999). The aerial part of this plant has long been used to treat dysentery, hematochezia, swelling, etc (How, 1997). Our previous investigation of the plant reported the iso-

* Foundation item: This work was financially supported by Yunnan Natural Science Foundation (2006C0045Q)

** Author for correspondence

Received date: 2007-08-09, Accepted date: 2007-11-02

作者简介: 来国防 (1972-) 男, 博士, 主管药师, 主要从事植物化学及药品质量标准的研究。

lation and characterization of three new triterpenoids (Lai *et al.*, 2003) and secoiridoid glycosides (Lai *et al.*, 2006). In continuation of our work on this genus, a new fructofuranose and ten known compounds were isolated from the EtOH extraction of the aerial part of *Helwingia chinensis*. Their structures were elucidated by intensive interpretation of spectral data.

Results and Discussion

Compound **1** was isolated as an amorphous powder, showed the molecular formula C₂₆H₂₈O₁₃ based on the [M-H]⁺ ion peak at *m/z* 547 in the negative FAB mass spectrum (MS) and NMR spectral data, which was confirmed by negative high resolution (HR)-FAB mass experiment at *m/z* 547.1438 ([M-H]⁺, C₂₆H₂₇O₁₃, cald. 547.1643). 13 unsaturated degrees and its ¹³C-NMR data showed the presence of two phenyl groups in the structure of **1**. The ¹³C-NMR and DEPT experiments indicated the presence of two methoxyls,

two methylenes, twelve methines and ten quaternary carbons. In the ¹H-NMR spectrum (see Table 1) of **1**, a typical ABX aromatic proton system at _H 7.14 (1H, d, *J* = 8.4 Hz), 7.23 (1H, dd, *J* = 8.4, 1.4 Hz) and 7.43 (1H, d, *J* = 1.4 Hz) and one *trans*-, -unsaturated olefin at _H 6.87, 8.01 (each 1H, d, *J* = 15.9 Hz) were observed, which suggested the presence of a (*E*)-caffeooyl moiety. This was confirmed by the HMBC experiments based on the correlations of between at _H 8.01 and _c 123.6 (s, C-1), 111.4 (d, C-2), 115.4 (d, C-8), 167.9 (s, C=O) (see Figure 2). Moreover, its ¹H-NMR spectrum contained another set of signals of a *trans*-, -unsaturated olefin at _H 6.71, 8.12 (each 1H, d, *J* = 15.8 Hz) and two aromatic proton signals at _H 7.13 (2H, s). The carbon signals at _c 166.9 (s), 149.3 (x2, s), 146.7 (d), 140.6 (s), 125.2 (s), 115.3 (d), 107.0 (x2, d), 56.5 (x2, q) showed the presence of 3, 5-dimethoxylcoumaroyl group. This was confirmed




Fig. 1 The structures of compounds **1-11**




Fig. 2 Key HMBC Correlations of compound 1

by the HMQC and HMBC experiments based on the correlations at δ 3.86 (6H, s) with τ 149.3 ($\times 2$, s), δ 6.71 with τ 125.2 (s) (see Figure). Except the above mentioned evidences, six carbon signals were present at δ 105.0 (s), 84.9 (d), 79.6 (d), 74.2 (d), 65.9 (t), 63.5 (t). The presence of -D-fructofuranose was identified by compared them with that of literature (Leslie *et al.*, 1980).

Long-rang correlations through 3J in the HMBC spectrum (See Figure 2) at δ 6.50 (1H, d, J = 7.7 Hz, H-3) with τ 166.9 (s, C=O) indicated that 3, 5-dimethoxylcoumaroyl group located at C-3. The weak correlations between at δ 6.87 (1H, d, J = 15.9 Hz) with τ 105.0 (s) showed that (E)-caffeyl moiety was located at C-2. Based on above mentioned evidence, the structure of 1 was identified to be 2-O-(E)-caffeyl-3-O-(3, 5-dimethoxyl)-(E)-coumaroyl-D-fructofuranose.

Compounds 2-11 were elucidated to be 2-O-D-fructofuranosyl-D-allopyranose (2) (Hongh *et al.*, 1986), liquiritigenin (3) (Tsutomu *et al.*, 1985), 4-hydroxy-2, 3-dihydro-flavonone 7-O-D-glucopyranoside (4) (Shul ts *et al.*, 2000), daidzin (5) (Park *et al.*, 1992), apigenin 5-O-D-glucopyranoside (6) (Martin *et al.*, 1991), apigenin-7-O-D-glucopyranoside (7) (Wang *et al.*, 1990), coumaric acid 4-O-D-glucopyranoside (8) (Cui *et al.*, 1992), glucopyranosylcaffeic acid (9) (Cui *et al.*, 1992), 3-hydroxyglutin-5-ene (10) (Antonio *et al.*, 1987), diosgenin 3-O-L-rhamnopyranosyl (1-2)-[L-arabinofuranosyl (1-3)]-D-glucopyranoside (11) (Xu *et al.*,

al., 1988) by NMR spectra data and compared with the report in the literatures. By the paper diffusion method (5 mm diameter) to assay their activities of against *Shigella flexneri*, *S. dysenteriae*, *S. sonnei*, *Mycobacterium tuberculosis*, -Hemolytic streptococcus and *Streptococcus pneumoniae*, compounds 1 and 6 showed weak activities against *S. flexneri* and *M. tuberculosis* in agar diffusion assay, respectively. In this paper, we describe the structures elucidation by various NMR spectra analysis.

Experimental

General experimental procedures Optical rotation was taken on a SEPA-300 polarimeter. FABMS data were obtained on a VG AutoSpec-3000 spectrometer. ^1H , ^{13}C , and 2D-NMR spectra were recorded on a Bruker AM 400 NMR and a DRX-500 spectrometer with TMS as internal standard. Silica gel (200-300 mesh, or silica gel H, 10-40 μm) for column chromatography (CC) and silica gel plate (GF254) for thin-layer chromatography (TLC) were obtained from the Qingdao Marine Chemical Factory, Qingdao, Shandong Province China.

Plant material The aerial parts of *Helwingia chinensis* were collected in Xishuangbanna, Yunnan, P. R. China in July 2002 (Lai *et al.*, 2003).

Extraction and Isolation The air-dried aerial parts (40 kg) were extracted twice with 95% EtOH at r. t. and the fractions were described as in the literature (Lai *et al.*, 2006). The fraction B was rechromatographed over silica gel (200-300 mesh), eluting with CHCl_3 -MeOH (99:1 to 10:1) to afford compound 10 (32 mg). The fraction D (3.1 g) was purified by repeated CC on silica gel (CHCl_3 -MeOH 50:1, 30:1, 20:1, 10:1) and then Sephadex LH-20 eluted by MeOH to give 1 (12 mg), 3 (16 mg) and 5 (24 mg). The fraction E (1.9 g) was purified by CC on silica gel; CHCl_3 -MeOH 20:1 to 4:1 and then Sephadex LH-20 (MeOH) to give 4 (28 mg), 6 (38 mg) and 7 (57 mg). The BuOH extract (15 g) was fractionated by CC on silica gel (200-300 mesh), CHCl_3 -MeOH from 20:1 to 2:1 and then RP-18 (MeOH-H₂O 6:4 to 8:2) to afford the compounds 2 (1.2 g), 8 (19 mg), 9 (325 mg) and 11 (14 mg).

Antimicrobial activity assay Inhibitory activities of compounds 1-11 against *Shigella flexneri*, *S. dysenteriae*, *S. sonnei*, *Mycobacterium tuberculosis*, -Hemolytic streptococcus and *Streptococcus pneumoniae* were determined by the paper disk diffusion assay on agar plates as described (Wang *et al.*, 2003). Only compounds 1 and 6 showed weak activities (6 mm/disk) against *S. flexneri* and *M. tuberculosis*, respectively.

Table 1 The ^1H , ^{13}C -NMR and 2D NMR spectral data of **1** in $\text{C}_5\text{D}_5\text{N}$ (500 MHz and 125 MHz, in ppm)

No.	^1H (Multiplicity, J in Hz)	c	HMBC (H C)
1	4.32 (2H, s)	65.9 t	C-2, 3
2		105.0 s	
3	6.50 (1H, d, 7.7)	79.6 d	C-1, 2, 4, 9
4	5.46 (1H, t, 7.5)	74.2 d	C-2, 3, 5, 6
5	4.78 (1H, m)	84.9 d	C-4, 6
6	4.57 (1H, dd, 11.8, 3.9) 4.64 (1H, dd, 11.8, 4.8)	63.5 t	C-4, 5 C-4, 5
1		123.6 s	
2	7.43 (1H, s)	111.4 d	C-1, 3, 4, 6, 7
3		149.0 s	
4		151.1 s	
5	7.14 (1H, d, 8.4)	106.8 d	C-1, 3, 4
6	7.23 (1H, d, 8.4)	116.7 d	C-1, 2, 4, 5, 7
7	8.01 (1H, d, 15.9)	145.8 d	C-1, 2, 6, 8, 9
8	6.87 (1H, d, 15.9)	115.4 d	C-2, 1, 7, 9
9		167.9 s	
1		125.2 s	
2, 6	7.13 (2H, s)	107.0 d	C-1, 6 (2), 3, 4, 5, 7
3, 5		149.3 s	
4		140.6 s	
7	8.12 (1H, d, 15.8)	146.7 d	C-1, 2, 6, 8, 9
8	6.71 (1H, d, 15.8)	115.3 d	C-1, 7, 9
9		166.9 s	
OMe	3.86 (6H, s)	56.5 q	C-3, 5

Chinenicside B (1) : white powder, $\text{C}_{26}\text{H}_{28}\text{O}_{13}$, negative FAB-MS m/z (%) : 547 [$\text{M}+\text{H}]^+$ (100), 532 [$\text{M}+16]^+$ (23), 255 (46), 223 (18), 177 (17); negative HR-FAB-MS m/z : 547.1438 ($[\text{M}+\text{H}]^+$, $\text{C}_{26}\text{H}_{27}\text{O}_{13}$, cald. 547.1643); ^1H -NMR and ^{13}C -NMR (see table 1).

2-O-D-fructofuranosyl-D-allopyranose (2) : white powder, $\text{C}_{12}\text{H}_{22}\text{O}_{11}$, negative FAB-MS m/z (%) : 341 [$\text{M}+\text{H}]^+$ (100), 293 (7), 127 (12); ^1H -NMR (CD_3OD , 400 MHz) : 5.20 (1H, d, $J=10.6$ Hz, H-1), 5.07 (1H, d, $J=10.6$ Hz, H-1), 5.15 (1H, d, $J=3.7$ Hz, H-1), 4.81 (1H, dd, $J=5.3, 11.9$ Hz, H-6), 4.42 (1H, dd, $J=2.0, 11.9$ Hz, H-6), 4.77 (1H, d, $J=10.2$ Hz, H-6), 4.52 (1H, d, $J=10.2$ Hz, H-6); ^{13}C -NMR (CD_3OD , 100 MHz) : 62.1 (t, C-1), 104.1 (s, C-2), 77.0 (d, C-3), 74.3 (d, C-4), 82.6 (d, C-5), 62.2 (t, C-6), 91.8 (d, C-1), 71.7 (d, C-2), 72.9 ($\times 2$, d, C-3, 5), 69.8 (d, C-4), 60.5 (t, C-6).

Liquiritigenin (3) : yellow powder, $\text{C}_{15}\text{H}_{12}\text{O}_4$, EIMS m/z (%) : 256 [$\text{M}]^+$ (100), 239 (7), 228 (12), 163 (17), 145 (22), 137 (73), 121 (95), 110 (41), 101 (55), 85 (40), 55 (70); ^1H -NMR (400 MHz, CD_3OD) : 2.75 (1H, dd, $J=2.8, 16.9$ Hz, H-3), 3.25 (1H, dd, $J=13.6, 16.5$ Hz, H-3), 5.55 (1H, dd, $J=2.8, 13.0$ Hz, H-2),

6.80 (1H, d, $J=2.0$ Hz, H-8), 6.88 (1H, dd, $J=2.0, 8.8$ Hz, H-6), 7.21 (2H, br. d, $J=8.6$ Hz, H-3, 5), 7.53 (2H, br. d, $J=8.7$ Hz, H-2, 6), 8.16 (1H, d, $J=8.8$ Hz, H-5); ^{13}C -NMR (100 MHz, CD_3OD) : 190.4 (s, C-4), 166.5 (s, C-7), 164.5 (s, C-9), 159.3 (s, C-4), 130.2 (s, C-1), 129.5 (d, C-5), 128.7 ($\times 2$, d, C-2, 6), 116.5 ($\times 2$, d, C-3, 5), 114.9 (s, C-10), 111.5 (d, C-6), 103.7 (d, C-8), 80.3 (d, C-2), 44.4 (t, C-3).

4-Hydroxy-2,3-dihydroflavonone 7-O-D-glucoside (4) : yellow powder, $\text{C}_{21}\text{H}_{22}\text{O}_9$, negative FAB-MS m/z (%) : 417 [$\text{M}+\text{H}]^+$ (64), 403 (16), 385 (5), 357 (8), 325 (100), 311 (978), 293 (12), 265 (17), 156 (4), 80 (14); ^1H -NMR (400 MHz, $\text{C}_5\text{D}_5\text{N}$) ppm: 5.54 (1H, dd, $J=2.8, 12.9$ Hz, H-2), 2.94 (1H, dd, $J=2.8, 16.6$ Hz, H-3), 3.22 (1H, dd, $J=13.0, 16.6$ Hz, H-3), 8.20 (1H, d, $J=8.7$ Hz, H-5), 6.94 (1H, dd, $J=2.0, 8.7$ Hz, H-6), 6.85 (1H, d, $J=2.2$ Hz, H-8), 7.52 (2H, d, $J=8.7$ Hz, H-2, 6), 7.40 (2H, d, $J=8.7$ Hz, H-3, 5), 5.70 (1H, d, $J=7.4$ Hz, H-1), 4.58 (1H, $J=11.8$ Hz, H-6), 4.34-4.44 (4H, m); ^{13}C -NMR (100 MHz, $\text{C}_5\text{D}_5\text{N}$) ppm: 190.2 (s, C-4), 166.7 (s, C-7), 164.5 (s, C-9), 158.9 (s, C-4), 133.4 (s, C-1), 129.5 (d, C-5), 128.3 (\times

2, d, C-2, 6), 117.5 (x2, d, C-3, 5), 114.5 (s, C-10), 111.8 (d, C-6), 80.2 (d, C-2), 44.8 (t, C-3), 102.2 (d, C-1), 78.9 (d, C-3), 78.6 (d, C-5), 74.9 (d, C-2), 71.5 (d, C-4), 62.5 (t, C-6).

Daidzin (5) : white powder, $C_{21}H_{20}O_9$, negative FAB-MS m/z (%) : 569 [M+154]⁺ (100), 415 [M-H]⁻ (15), 253 [M-162]⁻ (45); ¹H-NMR (500 MHz, CD₃OD) : 8.36 (1H, dd, $J=3.1, 8.2$ Hz, H-3, 5), 8.14 (1H, dd, $J=3.1, 8.2$ Hz, H-2, 6), 7.78 (1H, dd, $J=2.9, 8.1$ Hz, H-5), 7.43 (1H, s, H-2), 7.29 (1H, dd, $J=2.6, 8.2$ Hz, H-6), 5.82 (1H, d, $J=7.2$ Hz, H-1), 4.62 (2H, d, $J=11.9$ Hz, H-6), 4.27-4.45 (3H, m), 4.26 (1H, br s); ¹³C-NMR (125 MHz, CD₃OD) : 175.8 (s, C-4), 162.4 (s, C-7), 159.2 (s, C-9), 157.9 (s, C-4), 153.0 (d, C-2), 131.1 (d, C-2, 6), 127.9 (d, C-5), 125.2 (s, C-1), 119.3 (s, C-10), 116.4 (d, C-3, 5), 116.1 (d, C-6), 104.3 (d, C-8), 101.9 (d, C-1), 79.3 (d, C-3), 78.5 (d, C-5), 74.9 (d, C-2), 71.3 (d, C-4), 62.5 (t, C-6).

Apigenin 5-O- β -D-glucopyranoside (6) : white powder, $C_{21}H_{20}O_{10}$, negative FAB-MS m/z (%) : 431 [M-H]⁻ (100), 405 (8), 282 (13), 215 (17), 145 (24); ¹H-NMR (500 MHz, CD₃OD) ppm: 7.88 (1H, d, $J=8.6$ Hz, H-2, 6), 7.20 (1H, d, $J=6.1$ Hz, H-3, 5), 7.11 (1H, d, $J=1.2$ Hz, H-8), 6.91 (1H, s, H-3), 6.86 (1H, d, $J=1.5$ Hz, H-6), 5.86 (1H, d, $J=7.4$ Hz, H-1), 4.58 (2H, d, $J=10.6$ Hz, H-6), 4.44-4.34 (3H, m), 4.23 (1H, br s); ¹³C-NMR (125 MHz, CD₃OD) ppm: 182.9 (s, C-4), 165.0 (s, C-5), 164.1 (s, C-7), 162.9 (s, C-9), 162.6 (s, C-2), 157.9 (s, C-4), 129.0 (d, C-3, 5), 122.1 (s, C-1), 116.9 (d, C-2, 6), 106.6 (s, C-10), 104.1 (d, C-3), 100.7 (d, C-6), 95.4 (d, C-8), 101.8 (1d, C-1), 79.3 (d, C-3), 78.5 (d, C-5), 74.9 (d, C-2), 71.2 (d, C-4), 62.4 (t, C-6).

Apigenin 7-O- β -D-glucoside (7) : white powder, $C_{21}H_{20}O_{10}$, negative FAB-MS m/z (%) : 431 [M-H]⁻ (100), 415 (11), 269 [M-162]⁻ (70); ¹H-NMR (400 MHz, CD₃OD) : 8.11 (1H, s, H-2), 7.70 (2H, d, $J=8.5$ Hz, H-3, 5), 7.28 (2H, d, $J=8.5$ Hz, H-2, 6), 6.94 (1H, d, $J=2.7$ Hz, H-8), 6.86 (1H, d, $J=2.7$ Hz, H-6), 5.81 (1H, d, $J=7.3$ Hz, H-1), 4.57 (2H, d, $J=11.8$ Hz, H-6), 4.44-4.34 (3H, m), 4.20 (1H, br s); ¹³C-NMR (100 MHz, CD₃OD) : 181.4 (s, C-4), 164.4 (s, C-7), 159.4 (s, C-5), 158.1 (s, C-4), 163.0 (s, C-9), 153.9 (s, C-2), 131.0 (d, C-2, 6), 122.1 (s, C-1), 116.4 (d, C-3, 5), 107.3 (s, C-10), 100.7 (d, C-6), 95.2 (d, C-8), 100.7 (d, C-6), 101.7 (d, C-1), 79.3 (d, C-3), 78.5

(d, C-5), 74.8 (d, C-2), 71.2 (d, C-4), 62.4 (t, C-6).

Coumaric acid-4-O- β -D-glucopyranoside (8) : white powder, $C_{15}H_{18}O_8$, negative FAB-MS m/z (%) : 325 [M-H]⁻ (100), 163 (27); ¹H-NMR (400 MHz, C₅D₅N) : 7.30 (2H, d, $J=8.6$ Hz, H-2, 6), 6.71 (2H, d, $J=8.6$ Hz, H-3, 5), 7.75 (1H, d, $J=15.9$ Hz, H-), 6.30 (1H, d, $J=15.9$ Hz, H-), 5.64 (1H, d, $J=7.8$ Hz, H-1), 4.54 (1H, d, $J=10.5$ Hz, H-6), 4.42-4.26 (4H, m); ¹³C-NMR (100 MHz, C₅D₅N) : 170.5 (s, CO), 159.7 (s, C-4), 143.2 (d, C-), 130.5 (x2, d, C-2, 6), 127.5 (s, C-1), 117.8 (d, C-), 116.5 (x2, d, C-3, 5), 103.6 (d, C-1), 79.2 (d, C-3), 78.4 (d, C-5), 74.9 (d, C-2), 71.1 (d, C-4), 62.2 (t, C-6).

Glycopyanosylcaffeic acid (9) : white powder, $C_{15}H_{18}O_9$, negative FAB-MS m/z (%) : 341 [M-H]⁻ (100), 324 (11), 179 (26); ¹H-NMR (400 MHz, C₅D₅N) : 8.04 (1H, d, $J=15.9$ Hz, H-), 7.62 (1H, d, $J=1.7$ Hz, H-2), 7.53 (1H, d, $J=8.3$ Hz, H-5), 7.11 (1H, dd, $J=1.6, 8.3$ Hz, H-6), 6.87 (1H, d, $J=15.9$ Hz, H-), 5.62 (1H, d, $J=7.7$ Hz, H-1), 4.52 (1H, d, $J=10.2$ Hz, H-6), 4.42-4.26 (4H, m); ¹³C-NMR (100 MHz, C₅D₅N) : 169.7 (s, CO), 149.9 (s, C-3), 148.6 (s, C-4), 144.4 (d, C-), 130.9 (s, C-1), 120.9 (d, C-2), 118.9 (d, C-5), 116.5 (d, C-), 103.6 (d, C-1), 79.2 (d, C-3), 78.4 (d, C-5), 74.9 (d, C-2), 71.1 (d, C-4), 62.2 (t, C-6).

3-Hydroxyglutin-5-ene (10) : white powder, $C_{30}H_{50}O$, EFMS m/z (%) : 426 [M]⁺ (18), 408 (21), 393 (5), 274 (100), 259 (57), 245 (9), 205 (28), 173 (23), 119 (43), 109 (54), 95 (72), 69 (84), 55 (75); ¹H-NMR (400 MHz, CDCl₃) : 1.07, 1.04, 1.00, 0.95, 0.91, 0.90, 0.86, 0.82, 0.81, 0.76 (each, 3H, s), 3.37 (1H, br s, H-3), 5.52 (1H, d, $J=4.8$ Hz, H-5); ¹³C-NMR (100 MHz, CDCl₃) : 18.1 (t, C-1), 27.7 (t, C-2), 76.1 (d, C-3), 40.5 (s, C-4), 141.6 (s, C-5), 121.6 (d, C-6), 23.4 (t, C-7), 47.3 (d, C-8), 34.6 (s, C-9), 49.6 (d, C-10), 32.9 (t, C-11), 30.2 (C-12), 37.7 (s, C-13), 39.1 (s, C-14), 34.5 (t, C-15), 34.9 (t, C-16), 29.9 (s, C-17), 42.9 (d, C-18), 34.9 (t, C-19), 28.1 (s, C-20), 31.9 (t, C-21), 39.1 (t, C-22), 28.8 (q, C-23), 25.2 (q, C-24), 16.0 (q, C-25), 18.1 (q, C-26), 18.2 (q, C-27), 32.2 (q, C-28), 31.9 (q, C-29), 34.3 (q, C-30).

Diosgenin 3-O- β -L-rhamnopyranosyl(1 \rightarrow 2)-[β -L-arabinofuranosyl(1 \rightarrow 3)]- β -D-glucopyranoside (11) : white powder, $C_{44}H_{70}O_{16}$, negative FAB-MS m/z (%) : 853 [M-

H_2^+ (100), 721 (14), 691 (18), 479 (11), 447 (24), 411 (39); $^1\text{H-NMR}$ (400 MHz, $\text{C}_5\text{D}_5\text{N}$) : 6.30 (1H, br s, H-1), 5.31 (1H, d, $J = 4.7$ Hz, H-6), 4.98 (1H, d, $J = 3.7$ Hz, H-1''), 4.96 (1H, d, $J = 5.7$ Hz, H-1), 3.59 (1H, br d, $J = 10.7$ Hz, H-26), 3.51 (1H, dd, $J = 10.7, 10.8$ Hz, H-26), 4.63 (1H, dd, $J = 3.3, 9.3$ Hz, H-3), 1.78 (3H, d, $J = 6.3$ Hz, H-6), 1.14 (1H, d, $J = 6.9$ Hz, H-21), 1.05 (3H, s, H-19), 0.83 (3H, s, H-18), 0.70 (3H, d, $J = 5.0$ Hz, H-27); $^{13}\text{C-NMR}$ (100 MHz, $\text{C}_5\text{D}_5\text{N}$) : 37.6 (t, C-1), 31.9 (t, C-2), 77.8 (d, C-3), 39.9 (t, C-4), 140.8 (s, C-5), 121.9 (d, C-6), 32.3 (t, C-7), 31.7 (d, C-8), 50.3 (d, C-9), 37.2 (s, C-10), 21.2 (t, C-11), 39.0 (t, C-12), 40.5 (s, C-13), 56.7 (d, C-14), 30.2 (t, C-15), 81.2 (d, C-16), 62.9 (d, C-17), 19.5 (q, C-18), 16.4 (q, C-19), 42.0 (d, C-20), 15.1 (q, C-21), 109.3 (s, C-22), 32.4 (t, C-23), 29.3 (t, C-24), 30.7 (d, C-25), 66.9 (t, C-26), 17.4 (q, C-27), Gc: 100.2 (C-1), 78.1 (C-2), 77.9 (C-3), 76.8 (C-4), 77.0 (C-5), 62.6 (C-6), Rha: 102.0 (C-1), 72.6 (C-2), 72.9 (C-3), 74.2 (C-4), 69.6 (C-5), 18.7 (C-6), Ara: 109.7 (C-1''), 82.8 (C-2''), 77.5 (C-3''), 86.7 (C-4''), 61.4 (C-5'').

Acknowledgement: The authors are grateful to the analytical group of the Laboratory of Phytochemistry, Kunming Institute of Botany, Chinese Academy of Sciences for measuring NMR and MS data.

References :

- Antonio GG, Esteban EF, Angel GR, 1987. Triterpenes from *Maytenus horrida* [J]. *Phytochemistry*, **26**: 2785—2788
- Cui CB, Tezuka Y, Kikuchi T et al., 1992. Constituents of a fern, *Davallia mariesii* moore. IV. Isolation and structures of a novel norcarotane sesquiterpene glycoside, a chromone glucuronide, and two epicatechin glycosides [J]. *Chem Pharm Bull*, **40** (8) : 2035—2040
- Hough L, O'Brien E, 1986. Synthesis of 3-deoxy- and 3-deoxy-3-fluoro-sucrose, and α -D-allopyranosyl β -D-fructofuranoside [J]. *Carbohydr Res*, **147** (1) : 149—154
- How FC, 1997, A Dictionary of Traditional Chinese Drug [M]. Shanghai: Shanghai Science and Technology Press, 646
- Lai GF, Wang YF, Lu CH et al., 2003. Three novel triterpenoids from the aerial part of *Helwingia chinensis* [J]. *Helv Chim Acta*, **86**: 2136—2141
- Lai GF (来国防), Wang YF (王易芬), Wang M (王梅) et al., 2006. A new secoiridoid glycoside from *Helwingia chinensis* (Cornaceae) [J]. *Acta Bot Yunnan* (云南植物研究), **28**: 676—678
- Leslie H, Eugene O, 1980. α -D-Allopyranosyl β -D-fructofuranoside (α -D-sucrose) and its derivatives [J]. *Carbohydrate Res*, **84**: 95—102
- Martin G, Walter F, 1991. Flavonoid 5-glycosides from *Prunus cerasus* bark and their characteristic weak glycosidic bonding [J]. *Phytochemistry*, **30**: 1519—1521
- Park HH, Haramatsuka T, Noguchi H et al., 1992. Isoflavone glycosides exist as their 6-O-malonyl esters in *Pueraria lobata* and its cell suspension cultures [J]. *Chem Pharm Bull*, **40**: 1978—1980
- Shults EE, Petrova TN, Shakirov MM et al., 2000. Flavonoids of roots of *Glycyrrhiza uralensis* growing in Siberia [J]. *Chem Nat Compd*, **36**: 362—368
- Song ZP (宋滋圃), Hu WG (胡文光), 1990. Flora of China (中国植物志) [B]. **56**: 20—35
- Tseng CF, Iwakami S, Mikajiri A et al., 1992. Inhibition of in vitro prostaglandin and leukotriene biosyntheses by cinnamoyl- β -phenethylamine and N-acyldopamine derivatives [J]. *Chem Pharm Bull*, **40**: 396—400
- Tsutomo Nakanishi, Akira Inada, Kazuko Kambayashi et al., 1985. Flavonoid glycosides of the roots of *Glycyrrhiza uralensis* [J]. *Phytochemistry*, **24** (2) : 339—341
- Wang JX (王军宪), Hou GN (侯桂宁), 1990. Chemical study on the flower of *Ligusticum rumicinum* [J]. *Zhongguo Zhongyao Zaishi* (中国中药杂志), **3**: 168—170
- Wang YF, Lai GF, Lu CH et al., 2003. A new indolinizine from *Polygonatum kingianum* [J]. *Planta Med*, **69**: 1066—1068
- Xu XM (徐学民), Zhong ZC (钟炽昌), 1988. Chemical study on *Paris polyphylla* var. *chinensis* [J]. *Zhongcaoyao* (中草药), **19**: 194—198
- Zhonghua Bencaog Bianweihui (中华本草编委会), 1999. Zhonghua Bencaog (中华本草) [M], Vol. 15. Shanghai: Shanghai Science and Technology Press, 746—747